

For more information or to refer a patient, contact:

Aclan Dogan, MD

3303 SW Bond Avenue
Portland, OR 97239

(w) 503-494-2360

(c) 503-484-8614

(f) 503-494-7161

(e) dogana@ohsu.edu



OHSU Neurological Surgery

Mail code: CH8N

Center for Health & Healing

3303 S.W. Bond Avenue

Portland, OR 97239

tel 503 494-4314

fax 503 494-7161

www.ohsu.edu/neurosurgery

OHSU is an equal opportunity, affirmative action institution.



OHSU Neurological Surgery

Neurosurgical Case of the Month by Aclan Dogan, MD

June 2009:

Occipital artery to PICA bypass
for treatment of VA-PICA aneurysm



Occipital artery to PICA bybass for VA-PICA aneurysm

Patient history and diagnosis

An otherwise healthy 48-year-old male presented with: recent voice changes and hoarseness, vision problems, headache and increasing dizziness.

- was found to have a vertebral artery-PICA dissecting aneurysm
- has been taking coumadin for a while
- denies any difficulties with double vision, numbness or facial weakness, any history of seizures, extremity numbness or weakness, or gait difficulty.

Neurological Examination Results:

Mental status: Normal consciousness, orientation, affect and fluency

Cranial Nerves: II-XII intact on detailed examination

Motor: Normal strength, muscle bulk, and tone


Sensory: Intact to pinprick and light touch

Cerebellar: Normal finger-to-nose and rapid alternaing movements

Gait: Normal, Tandem and Romberg negative

Deep Tendon Reflexes: Present and normoactive

Pathologic Reflexes: Absent




Right AP vertebral artery angiogram: Shows dissecting aneurysm and minimal blood flow to the basilar artery.



Right lateral vertebral artery angiogram: Shows dissecting aneurysm, PICA is originating from the dome of the aneurysm and there is minimal blood flow to the basilar artery.





Left lateral vertebral artery angiogram: Shows hypoplastic vessel and very limited blood flow to the basilar artery.



Right lateral carotid artery angiogram: Shows good size occipital artery for possible by-pass surgery.

Plan and Surgical Treatment

Endovascular coiling of the PICA dissecting aneurysm presents significant risk for a PICA stroke as the origin is the aneurysm dome. Similarly, surgical clipping and reconstruction of this aneurysm presents the same risk. Therefore right occipital artery to right PICA by-pass anastomosis and trapping of the aneurysm is planned. A right far lateral craniotomy was performed and the right occipital artery harvested. Initially the occipital artery was anastomosed to the telelevelotonsillar segment of the PICA using 10/0 suture. The aneurysm was then trapped with clips between the proximal PICA and the distal vertebral artery.



Postoperative lateral right carotid angiogram: Shows patent and open anastomosis between occipital artery and PICA



Postoperative lateral right vertebral artery angiogram: Shows occlusion of VA and aneurysm without any residual filling.



Postoperative AP left vertebral artery angiogram: Shows no aneurysm filling but basilar artery flow has improved significantly



Right lateral carotid artery angiogram at 4 months post-op: Shows patent occipital artery to PICA anastomosis

Outcome

Postoperatively the patient was neurologically intact with no neurological deficit during postoperative follow up.