



Oregon Health & Science University
Hospital and Clinics Provider's Orders

PO7071



ADULT AMBULATORY INFUSION ORDER
Pentamidine (PENTAM) Infusion

Page 1 of 3

ACCOUNT NO.
MED. REC. NO.
NAME
BIRTHDATE

Patient Identification

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.

Weight: _____ kg Height: _____ cm

Allergies: _____

Diagnosis Code: _____

Treatment Start Date: _____ Patient to follow up with provider on date: _____

****This plan will expire after 365 days at which time a new order will need to be placed****

GUIDELINES FOR ORDERING

1. Send **FACE SHEET and H&P or most recent chart note.**
2. 12 Lead ECG should be completed prior to treatment with pentamidine. **Results MUST be faxed with this order set to be kept on record within the infusion pharmacy's electronic medical record.**
3. Avoid use in patients with diagnosed or suspected congenital long QT syndrome.
4. Use with caution in patients with pre-existing hypotension. Severe hypotension including fatalities, has been observed even after a single dose.
5. Use with caution in patients with pre-existing cardiovascular disease, diabetes mellitus, or hypocalcemia.
6. Use with caution in patients receiving nephrotoxic drugs such as aminoglycosides, amphotericin B, cisplatin, foscarnet, or vancomycin.

OTHER:

- 12 Lead ECG, routine, ONCE every _____ weeks

LABS:

- CMP (includes blood glucose), Routine, ONCE, every _____ (visit)(days)(weeks)(months) – Circle One
- Glucose (serum), Routine, ONCE, every _____ (visit)(days)(weeks)(months) – Circle One
- CBC with differential, Routine, ONCE, weekly during therapy
- CBC with differential, Routine, ONCE, every _____ (visit)(days)(weeks)(months) – Circle One
- Labs already drawn. Date: _____

MEDICATIONS:

pentamidine (PENTAM) in dextrose 5% 250 mL, intravenous, ONCE

- 300 mg
- 3 mg/kg = _____ mg
- 4 mg/kg = _____ mg

Infuse slowly over 1-2 hours. Vesicant. Flush line with D5W before and after infusion.

Interval: (must check one)

- Once
- Once daily x _____ doses
- _____ times per week x _____ doses
- Monthly x _____ doses



Oregon Health & Science University
Hospital and Clinics Provider's Orders

ADULT AMBULATORY INFUSION ORDER
Pentamidine (PENTAM) Infusion

Page 2 of 3

ACCOUNT NO.
MED. REC. NO.
NAME
BIRTHDATE

Patient Identification

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.

NURSING ORDERS:

1. Review patient's SCr, BUN, calcium, and blood glucose during each visit. Notify provider if laboratory values are abnormal.
2. VITAL SIGNS – Monitor patient's blood pressure for hypotension during and after infusion
3. Instruct patient to lie supine during the infusion. Patient should rise slowly after administration to avoid dizziness and other potentially severe hypotensive effects.
4. This medication is a vesicant. Avoid extravasation. Assess catheter position before and during infusion
5. Follow facility policies and/or protocols for vascular access maintenance with appropriate flush solution, dec clotting (alteplase), and/or dressing changes.

AS NEEDED MEDICATIONS:

1. prochlorperazine (COMPAZINE) tablet, 10mg, oral, AS NEEDED, x1 doses for nausea/vomiting
2. LORazepam (ATIVAN) tablet, 1 mg, oral, AS NEEDED, x1 dose for nausea/vomiting. Hold if patient does not have a driver

By signing below, I represent the following:

I am responsible for the care of the patient (*who is identified at the top of this form*);

I hold an active, unrestricted license to practice medicine in: Oregon _____ (*check box that corresponds with state where you provide care to patient and where you are currently licensed. Specify state if not Oregon*);

My physician license Number is # _____ (MUST BE COMPLETED TO BE A VALID PRESCRIPTION); and I am acting within my scope of practice and authorized by law to order Infusion of the medication described above for the patient identified on this form.

Provider signature: _____ **Date/Time:** _____

Printed Name: _____ **Phone:** _____ **Fax:** _____



Oregon Health & Science University
Hospital and Clinics Provider's Orders

ADULT AMBULATORY INFUSION ORDER
Pentamidine (PENTAM) Infusion

Page 3 of 3

ACCOUNT NO.
MED. REC. NO.
NAME
BIRTHDATE

Patient Identification

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.

Central Intake:

Phone: 971-262-9645 (providers only) Fax: 503-346-8058

Please check the appropriate box for the patient's preferred clinic location:

Beaverton

OHSU Knight Cancer Institute
15700 SW Greystone Court
Beaverton, OR 97006

Phone number: 971-262-9000

Fax number: 503-346-8058

NW Portland

Legacy Good Samaritan campus
Medical Office Building 3, Suite 150
1130 NW 22nd Ave.
Portland, OR 97210

Phone number: 971-262-9600

Fax number: 503-346-8058

Gresham

Legacy Mount Hood campus
Medical Office Building 3, Suite 140
24988 SE Stark
Gresham, OR 97030

Phone number: 971-262-9500

Fax number: 503-346-8058

Tualatin

Legacy Meridian Park campus
Medical Office Building 2, Suite 140
19260 SW 65th Ave.
Tualatin, OR 97062

Phone number: 971-262-9700

Fax number: 503-346-8058

Infusion orders located at: www.ohsuknight.com/infusionorders