
 <p>Oregon Health & Science University Hospital and Clinics Provider's Orders</p> <p>P07071 </p> <p>ADULT AMBULATORY INFUSION ORDER Cosyntropin (CORTROSYN) Stimulation Test</p> <p>Page 1 of 2</p>	<p>ACCOUNT NO. MED. REC. NO. NAME BIRTHDATE</p> <p style="text-align: right;"><i>Patient Identification</i></p>
ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.	

Weight: _____ kg Height: _____ cm

Allergies: _____

Diagnosis Code: _____

Treatment Start Date: _____ Patient to follow up with provider on date: _____

****This plan will expire after 365 days at which time a new order will need to be placed****

GUIDELINES FOR ORDERING

1. Send **FACE SHEET and H&P or most recent chart note.**
2. Patient should not receive corticosteroids or spironolactone within 24 hours prior to the cosyntropin test.
3. The Low Dose Protocol is not recommended in critically-ill patients.

LABS:

- ACTH Stimulation Test, Serum, Routine, ONCE, every ____ (visit)(days)(weeks)(months) – *Circle One*
- Cortisol, Serum Routine, ONCE, ONCE, every ____ (visit)(days)(weeks)(months) – *Circle One*
 - Draw baseline immediately before administration of Cosyntropin IVP
 - Draw 20 minutes after administration of Cosyntropin IVP (if cosyntropin 1 mcg test is ordered)
 - Draw 30 minutes after administration of Cosyntropin IVP
 - Draw 60 minutes after administration of Cosyntropin IVP

NURSING ORDERS:

1. Draw baseline ACTH and cortisol labs.
2. Administer Cosyntropin IVP over 2 minutes and flush with 5-6 mL normal saline flush.
3. Draw 30+ and 60+ Cortisol labs.
4. Only use a 22 gauge or larger needle.
5. Release labs as drawn so times are accurate. Do not release all labs at one time
6. Follow facility policies and/or protocols for vascular access maintenance with appropriate flush solution, dec clotting (alteplase), and/or dressing changes.

MEDICATIONS:

Cosyntropin (select one):

- cosyntropin (CORTROSYN) injection 1 mcg, intravenous, ONCE over 2 minutes
Low Dose Protocol. Diluted in sodium chloride 0.9%. Infuse over 2 minutes.
- cosyntropin (CORTROSYN) injection 0.25 mg, intravenous, ONCE over 2 minutes
Standard Dose Protocol. Diluted in sodium chloride 0.9%. Infuse over 2 minutes.



Oregon Health & Science University
Hospital and Clinics Provider's Orders

ADULT AMBULATORY INFUSION ORDER
**Cosyntropin (CORTROSYN)
Stimulation Test**

Page 2 of 2

ACCOUNT NO.
MED. REC. NO.
NAME
BIRTHDATE

Patient Identification

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.

By signing below, I represent the following:

I am responsible for the care of the patient (*who is identified at the top of this form*);

I hold an active, unrestricted license to practice medicine in: Oregon _____ (*check box that corresponds with state where you provide care to patient and where you are currently licensed. Specify state if not Oregon*);

My physician license Number is # _____ (MUST BE COMPLETED TO BE A VALID PRESCRIPTION); and I am acting within my scope of practice and authorized by law to order Infusion of the medication described above for the patient identified on this form.

Provider signature: _____ **Date/Time:** _____

Printed Name: _____ **Phone:** _____ **Fax:** _____

Central Intake:

Phone: 971-262-9645 (providers only) Fax: 503-346-8058

Please check the appropriate box for the patient's preferred clinic location:

Beaverton

OHSU Knight Cancer Institute
15700 SW Greystone Court
Beaverton, OR 97006
Phone number: 971-262-9000
Fax number: 503-346-8058

NW Portland

Legacy Good Samaritan campus
Medical Office Building 3, Suite 150
1130 NW 22nd Ave
Portland, OR 97210
Phone number: 971-262-9600
Fax number: 503-346-8058

Gresham

Legacy Mount Hood campus
Medical Office Building 3, Suite 140
24988 SE Stark
Gresham, OR 97030
Phone number: 971-262-9500
Fax number: 503-346-8058

Tualatin

Legacy Meridian Park campus
Medical Office Building 2, Suite 140
19260 SW 65th Ave
Tualatin, OR 97062
Phone number: 971-262-9700
Fax number: 503-346-8058

Infusion orders located at: www.ohsuknight.com/infusionorders