
 <p style="text-align: center;">Oregon Health & Science University Hospital and Clinics Provider's Orders</p> <div style="display: flex; align-items: center;"> <div style="writing-mode: vertical-rl; transform: rotate(180deg); font-size: small; margin-right: 5px;">PO7071</div>  </div> <p style="text-align: center; font-weight: bold;">ADULT AMBULATORY INFUSION ORDER Golimumab (SIMPONI ARIA) Infusion</p> <p style="text-align: center; font-size: small;">Page 1 of 3</p>	<p>ACCOUNT NO. MED. REC. NO. NAME BIRTHDATE</p> <p style="text-align: right; font-size: x-small;"><i>Patient Identification</i></p>
ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.	

Weight: _____ kg Height: _____ cm

Allergies: _____

Diagnosis Code: _____

Treatment Start Date: _____ Patient to follow up with provider on date: _____

****This plan will expire after 365 days at which time a new order will need to be placed****

GUIDELINES FOR ORDERING

1. Send **FACE SHEET and H&P or most recent chart note.**
2. Hepatitis B (Hep B surface antigen and core antibody total) screening must be completed prior to initiation of treatment and the patient should not be infected. Please send results with order.
3. A Tuberculin test must have been placed and read as negative prior to initiation of treatment (PPD or QuantiFERON Gold blood test). Please send results with order. If result is indeterminate, a follow up chest X-ray must be performed to rule out TB. Please send results with order.

PRE-SCREENING: (Results must be available prior to initiation of therapy):

- Hepatitis B surface antigen and core antibody total test results scanned with orders.
- Tuberculin skin test or QuantiFERON Gold blood test results scanned with orders.
- Chest X-Ray result scanned with orders if TB test result is indeterminate.

LABS:

1. CBC with differential, Routine, ONCE, every visit, If on methotrexate or leflunomide
2. CMP, Routine, ONCE, every visit, If on methotrexate or leflunomide

NURSING ORDERS:

1. TREATMENT PARAMETER – Hold treatment and contact provider if Hepatitis B surface antigen or core antibody total test result is positive, TB test result is positive, or if screening has not been performed.
2. VITAL SIGNS – Prior to infusion and at the completion of the infusion.
3. Infusions to be scheduled at weeks 0 and 4, then every 8 weeks thereafter.

MEDICATIONS: (check all that apply)

golimumab (SIMPONI ARIA) 2 mg/kg in sodium chloride 0.9%, intravenous, ONCE, over 30 minutes

- Initial doses:** Every 4 weeks for 2 treatments (week 0, 4)
- Maintenance doses:** Every 8 weeks thereafter (week 12 and beyond)

AS NEEDED MEDICATIONS:

1. acetaminophen (TYLENOL) tablet, 650 mg, oral, EVERY 4 HOURS AS NEEDED for headache, fever, chills or malaise
2. diphenhydramINE (BENADRYL) capsule, 50 mg, oral, EVERY 4 HOURS AS NEEDED for itching



Oregon Health & Science University
Hospital and Clinics Provider's Orders

ADULT AMBULATORY INFUSION ORDER
**Golimumab (SIMPONI ARIA)
Infusion**

Page 2 of 3

ACCOUNT NO.
MED. REC. NO.
NAME
BIRTHDATE

Patient Identification

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.

HYPERSENSITIVITY MEDICATIONS:

1. NURSING COMMUNICATION – If hypersensitivity or infusion reactions develop, temporarily hold the infusion and notify provider immediately. Administer emergency medications per the Treatment Algorithm for Acute Infusion Reaction (OHSU HC-PAT-133-GUD, HMC C-132). Refer to algorithm for symptom monitoring and continuously assess as grade of severity may progress.
2. diphenhydrAMINE (BENADRYL) injection, 25-50 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
3. EPINEPHrine HCl (ADRENALIN) injection, 0.3 mg, intramuscular, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
4. hydrocortisone sodium succinate (SOLU-CORTEF) injection, 100 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
5. famotidine (PEPCID) injection, 20 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction

By signing below, I represent the following:

I am responsible for the care of the patient (*who is identified at the top of this form*);

I hold an active, unrestricted license to practice medicine in: Oregon _____ (*check box that corresponds with state where you provide care to patient and where you are currently licensed. Specify state if not Oregon*);

My physician license Number is # _____ (MUST BE COMPLETED TO BE A VALID PRESCRIPTION); and I am acting within my scope of practice and authorized by law to order Infusion of the medication described above for the patient identified on this form.

Provider signature: _____ **Date/Time:** _____

Printed Name: _____ **Phone:** _____ **Fax:** _____



Oregon Health & Science University
Hospital and Clinics Provider's Orders

ADULT AMBULATORY INFUSION ORDER
**Golimumab (SIMPONI ARIA)
Infusion**

Page 3 of 3

ACCOUNT NO.
MED. REC. NO.
NAME
BIRTHDATE

Patient Identification

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.

Central Intake:

Phone: 971-262-9645 (providers only) Fax: 503-346-8058

Please check the appropriate box for the patient's preferred clinic location:

Beaverton

OHSU Knight Cancer Institute
15700 SW Greystone Court
Beaverton, OR 97006

Phone number: 971-262-9000

Fax number: 503-346-8058

NW Portland

Legacy Good Samaritan campus
Medical Office Building 3, Suite 150
1130 NW 22nd Ave
Portland, OR 97210

Phone number: 971-262-9600

Fax number: 503-346-8058

Gresham

Legacy Mount Hood campus
Medical Office Building 3, Suite 140
24988 SE Stark
Gresham, OR 97030

Phone number: 971-262-9500

Fax number: 503-346-8058

Tualatin

Legacy Meridian Park campus
Medical Office Building 2, Suite 140
19260 SW 65th Ave
Tualatin, OR 97062

Phone number: 971-262-9700

Fax number: 503-346-8058

Infusion orders located at: www.ohsuknight.com/infusionorders