



Oregon Health & Science University  
Hospital and Clinics Provider's Orders

PO7071



ADULT AMBULATORY INFUSION ORDER

**Lecanemab (LEQEMBI)**

Infusion

Page 1 of 3

ACCOUNT NO.

MED. REC. NO.

NAME

BIRTHDATE

*Patient Identification*

**ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.**

Weight: \_\_\_\_\_ kg      Height: \_\_\_\_\_ cm

Allergies: \_\_\_\_\_

Diagnosis Code: \_\_\_\_\_

Treatment Start Date: \_\_\_\_\_ Patient to follow up with provider on date: \_\_\_\_\_

**\*\*This plan will expire after 365 days at which time a new order will need to be placed\*\***

**GUIDELINES FOR ORDERING**

1. Send **FACE SHEET and H&P or most recent chart note.**
2. Confirm the presence of amyloid beta pathology prior to initiating treatment.
3. Obtain a recent (within one year) brain MRI prior to initiating treatment to evaluate for pre-existing Amyloid Related Imaging Abnormalities (ARIA).
4. Obtain an MRI prior to the 5th, 7th, and 14th infusions. If radiographically observed ARIA occurs, treatment recommendations are based on type, severity, and presence of symptoms.
5. Enhanced clinical vigilance for ARIA is recommended during the first 14 weeks of treatment with LEQEMBI. If patient experiences symptoms suggestive of ARIA, clinical evaluation should be performed, including MRI if indicated. If ARIA is observed on MRI, careful clinical evaluation should be performed prior to continuing treatment.

**NURSING ORDERS:**

1. Monitor for infusion reactions during infusion and observe for at least 1 hour after completion of first two infusions. The highest incidence of infusion reactions occurs with the first infusion.
2. Confirm an MRI was performed prior to the 5<sup>th</sup>, 7<sup>th</sup>, and 14<sup>th</sup> infusions.
3. Follow facility policies and/or protocols for vascular access maintenance with appropriate flush solution, dec clotting (alteplase), and/or dressing changes.

**PRE-MEDICATIONS:** (Administer 30 minutes prior to infusion)

**Note to provider: Please select which medications below, if any, you would like the patient to receive prior to treatment by checking the appropriate box(s)**

1. Acetaminophen (TYLENOL) tablet, 650 mg, oral, ONCE AS NEEDED IF PATIENT HAD PRIOR INFUSION REACTION, every visit
2. DiphenhydrAMINE (BENADRYL) capsule, 25 mg, oral, ONCE AS NEEDED IF PATIENT HAD PRIOR INFUSION REACTION every visit.  
**Give either diphenhydrAMINE or loratadine, not both.**
3. Loratadine (CLARITIN) tablet, 10 mg, oral, ONCE AS NEEDED IF PATIENT HAD PRIOR INFUSION REACTION and diphenhydrAMINE is not given, every visit.  
**Give either loratadine or diphenhydrAMINE, not both.**
4. Dexamethasone (DECADRON), 10 mg, intravenous, ONCE AS NEEDED IF PATIENT HAD PRIOR INFUSION REACTION, every visit.



Oregon Health & Science University  
Hospital and Clinics Provider's Orders

ADULT AMBULATORY INFUSION ORDER  
**Lecanemab (LEQEMBI) Infusion**

Page 2 of 3

ACCOUNT NO.  
MED. REC. NO.  
NAME  
BIRTHDATE

*Patient Identification*

**ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.**

**MEDICATIONS:**

- Lecanemab (LEQEMBI), 10 mg/kg in sodium chloride 0.9% 250 mL, intravenous, ONCE, over 1 hour, every 2 weeks.

**HYPERSENSITIVITY MEDICATIONS:**

1. NURSING COMMUNICATION – If hypersensitivity or infusion reactions develop, temporarily hold the infusion and notify provider immediately. Administer emergency medications per the Treatment Algorithm for Acute Infusion Reaction (OHSU HC-PAT-133-GUD, HMC C-132). Refer to algorithm for symptom monitoring and continuously assess as grade of severity may progress.
2. diphenhydrAMINE (BENADRYL) injection, 25-50 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
3. EPINEPHrine HCl (ADRENALIN) injection, 0.3 mg, intramuscular, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
4. hydrocortisone sodium succinate (SOLU-CORTEF) injection, 100 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
5. famotidine (PEPCID) injection, 20 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction

**By signing below, I represent the following:**

I am responsible for the care of the patient (*who is identified at the top of this form*);

I hold an active, unrestricted license to practice medicine in:  Oregon  \_\_\_\_\_ (*check box that corresponds with state where you provide care to patient and where you are currently licensed. Specify state if not Oregon*);

**My physician license Number is # \_\_\_\_\_ (MUST BE COMPLETED TO BE A VALID PRESCRIPTION);** and I am acting within my scope of practice and authorized by law to order Infusion of the medication described above for the patient identified on this form.

**Provider signature:** \_\_\_\_\_ **Date/Time:** \_\_\_\_\_

**Printed Name:** \_\_\_\_\_ **Phone:** \_\_\_\_\_ **Fax:** \_\_\_\_\_



Oregon Health & Science University  
Hospital and Clinics Provider's Orders

ADULT AMBULATORY INFUSION ORDER  
**Lecanemab (LEQEMBI) Infusion**

Page 3 of 3

ACCOUNT NO.  
MED. REC. NO.  
NAME  
BIRTHDATE

*Patient Identification*

**ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.**

OLC Central Intake Nurse:

Phone: 971-262-9645 (providers only) Fax: 503-346-8058

***Please check the appropriate box for the patient's preferred clinic location:***

**Beaverton**

OHSU Knight Cancer Institute  
15700 SW Greystone Court  
Beaverton, OR 97006

Phone number: 971-262-9000

Fax number: 503-346-8058

**NW Portland**

Legacy Good Samaritan campus  
Medical Office Building 3, Suite 150  
1130 NW 22nd Ave.

Portland, OR 97210

Phone number: 971-262-9600

Fax number: 503-346-8058

**Gresham**

Legacy Mount Hood campus  
Medical Office Building 3, Suite 140  
24988 SE Stark  
Gresham, OR 97030

Phone number: 971-262-9500

Fax number: 503-346-8058

**Tualatin**

Legacy Meridian Park campus  
Medical Office Building 2, Suite 140  
19260 SW 65th Ave.  
Tualatin, OR 97062

Phone number: 971-262-9700

Fax number: 503-346-8058

Infusion orders located at: [www.ohsuknight.com/infusionorders](http://www.ohsuknight.com/infusionorders)