
 <p style="text-align: center;">Oregon Health & Science University Hospital and Clinics Provider's Orders</p> <div style="display: flex; align-items: center;"> <div style="writing-mode: vertical-rl; transform: rotate(180deg); font-size: small; margin-right: 5px;">PO7071</div>  </div> <p style="text-align: center;">ADULT AMBULATORY INFUSION ORDER Ferric Derisomaltose (MONOFERRIC) Infusion Page 1 of 3</p>	<p>ACCOUNT NO. _____</p> <p>MED. REC. NO. _____</p> <p>NAME _____</p> <p>BIRTHDATE _____</p> <p style="text-align: right; font-size: small;"><i>Patient Identification</i></p>
ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.	

Weight: _____ kg **Height:** _____ cm

Allergies: _____

Diagnosis Code: _____

Treatment Start Date: _____ **Patient to follow up with provider on date:** _____

****This plan will expire after 365 days at which time a new order will need to be placed****

GUIDELINES FOR ORDERING

1. Send **FACE SHEET and H&P or most recent chart note.**
2. Ferritin must be obtained prior to start of treatment. Labs drawn date: _____

NURSING ORDERS:

1. TREATMENT PARAMETERS – Hold treatment and notify provider if Ferritin greater than 300 ng/mL.
2. Instruct patient to set follow up appointment with provider for follow up labs.
3. Monitor patient for signs and symptoms of hypersensitivity during the infusion and for at least 30 minutes after completion of the infusion.
4. Follow facility policies and/or protocols for vascular access maintenance with appropriate flush solution, declotting (alteplase), and/or dressing changes.

MEDICATIONS:

Ferric Derisomaltose (MONOFERRIC) dosing: (must check one)

- For weight greater than or equal to 50 kg:**
1,000 mg in sodium chloride 0.9% 100 mL, intravenous, ONCE, over 20 minutes
- For weight less than 50 kg:**
20 mg/kg in sodium chloride 0.9% 100 mL, intravenous, ONCE, over 20 minutes

AS NEEDED MEDICATIONS:

1. sodium chloride 0.9% bolus, intravenous, 500 mL, AS NEEDED x1 dose, for vein discomfort.
Give concurrently with ferric gluconate

HYPERSENSITIVITY MEDICATIONS:

1. NURSING COMMUNICATION – If hypersensitivity or infusion reactions develop, temporarily hold the infusion and notify provider immediately. Administer emergency medications per the Treatment Algorithm for Acute Infusion Reaction (OHSU HC-PAT-133-GUD, HMC C-132). Refer to algorithm for symptom monitoring and continuously assess as grade of severity may progress.
2. diphenhydrAMINE (BENADRYL) injection, 25-50 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
3. EPINEPHrine HCl (ADRENALIN) injection, 0.3 mg, intramuscular, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
4. hydrocortisone sodium succinate (SOLU-CORTEF) injection, 100 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction



Oregon Health & Science University
Hospital and Clinics Provider's Orders

ADULT AMBULATORY INFUSION ORDER

**Ferric Derisomaltose
(MONOFERRIC) Infusion**

Page 2 of 3

ACCOUNT NO.
MED. REC. NO.
NAME
BIRTHDATE

Patient Identification

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.

5. famotidine (PEPCID) injection, 20 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction

By signing below, I represent the following:

I am responsible for the care of the patient (*who is identified at the top of this form*);

I hold an active, unrestricted license to practice medicine in: Oregon _____ (*check box that corresponds with state where you provide care to patient and where you are currently licensed. Specify state if not Oregon*);

My physician license Number is # _____ (MUST BE COMPLETED TO BE A VALID PRESCRIPTION); and I am acting within my scope of practice and authorized by law to order Infusion of the medication described above for the patient identified on this form.

Provider signature: _____ **Date/Time:** _____

Printed Name: _____ **Phone:** _____ **Fax:** _____

Central Intake:

Phone: 971-262-9645 (providers only) Fax: 503-346-8058

Please check the appropriate box for the patient's preferred clinic location:

Beaverton

OHSU Knight Cancer Institute
15700 SW Greystone Court
Beaverton, OR 97006

Phone number: 971-262-9000

Fax number: 503-346-8058

NW Portland

Legacy Good Samaritan campus
Medical Office Building 3, Suite 150
1130 NW 22nd Ave.
Portland, OR 97210

Phone number: 971-262-9600

Fax number: 503-346-8058

Gresham

Legacy Mount Hood campus
Medical Office Building 3, Suite 140
24988 SE Stark
Gresham, OR 97030

Phone number: 971-262-9500

Fax number: 503-346-8058

Tualatin

Legacy Meridian Park campus
Medical Office Building 2, Suite 140
19260 SW 65th Ave.
Tualatin, OR 97062

Phone number: 971-262-9700

Fax number: 503-346-8058

Infusion orders located at: www.ohsuknight.com/infusionorders



**Oregon Health & Science University
Hospital and Clinics Provider's Orders**

ADULT AMBULATORY INFUSION ORDER

**Ferric Derisomaltose
(MONOFERRIC) Infusion**

Page 3 of 3

ACCOUNT NO.

MED. REC. NO.

NAME

BIRTHDATE

Patient Identification

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.