



Oregon Health & Science University
Hospital and Clinics Provider's Orders

P07071



ADULT AMBULATORY INFUSION ORDER
Natalizumab (TYSABRI)

Page 1 of 3

ACCOUNT NO.
MED. REC. NO.
NAME
BIRTHDATE

Patient Identification

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.

Weight: _____ kg Height: _____ cm

Allergies: _____

Diagnosis Code: _____

Treatment Start Date: _____ Patient to follow up with provider on date: _____

****This plan will expire after 365 days at which time a new order will need to be placed****

GUIDELINES FOR ORDERING

1. Send **FACE SHEET and H&P or most recent chart note.**
2. Natalizumab is restricted to credentialed prescribers only through the TOUCH™ Prescribing Program
 - a. Prescribers **MUST** be enrolled in the TOUCH™ Prescribing Program
 - b. Patients **MUST** be enrolled in the TOUCH™ Prescribing Program
 - c. Contact the TOUCH™ Prescribing Program at 1-800-456-2255 for details and enrollment
 - d. Notify Biogen Customer Service of any adverse reactions at 1-800-456-2255

LABS:

During first year of treatment:

- Basic Metabolic Set, Routine, ONCE, every _____ (visit)(days)(weeks)(months) – *Circle One*
- Basic Metabolic Set, Routine, ONCE, every 3 months (after first year of treatment)
- CBC with differential, Routine, ONCE, every _____ (visit)(days)(weeks)(months) – *Circle One*
- CBC with differential, Routine, ONCE, every 6 months (after first year of treatment)
- HCG Beta, PLASMA, Routine, ONCE, every _____ (visit)(days)(weeks)(months) – *Circle One*

After first year of treatment:

- Basic Metabolic Set, Routine, ONCE, every _____ (visit)(days)(weeks)(months) – *Circle One*
- Basic Metabolic Set, Routine, ONCE, every 3 months (after first year of treatment)
- CBC with differential, Routine, ONCE, every _____ (visit)(days)(weeks)(months) – *Circle One*
- CBC with differential, Routine, ONCE, every 6 months (after first year of treatment)
- HCG Beta, PLASMA, Routine, ONCE, every _____ (visit)(days)(weeks)(months) – *Circle One*

PRE-MEDICATIONS:

- sodium chloride 0.9% solution, 250 mL, intravenous, Infuse at rate necessary to keep vein open (KVO) until natalizumab is started and for 1 hour after infusion is complete, then discontinue

MEDICATIONS:

- Natalizumab (TYSABRI), 300 mg, intravenous, in NaCl 0.9% 100 mL, ONCE, over 60 minutes

Interval: (must check one)

- Once
- Every 4 weeks x _____ doses
- Every 4 weeks until discontinued



Oregon Health & Science University
Hospital and Clinics Provider's Orders

ADULT AMBULATORY INFUSION ORDER

Natalizumab (TYSABRI)

Page 2 of 3

ACCOUNT NO.
MED. REC. NO.
NAME
BIRTHDATE

Patient Identification

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.

NURSING ORDERS:

1. Follow facility policies and/or protocols for vascular access maintenance with appropriate flush solution, declotting (alteplase), and/or dressing changes.
2. Please indicate patient's TOUCH™ Prescribing Program authorization number from Biogen.
Authorization # is: _____
3. Review "Medication Guide" with patient. Review and complete TOUCH™ on-line checklist with patient. Proceed according to guidelines. **Note:** If patient answers "Yes" to any of the questions 1 through 3 in Step 3 of the checklist, **DO NOT** infuse natalizumab. Contact physician to obtain further orders regarding infusion.
4. Assess patient for signs of infection and notify prescriber if present.
5. Verify that urine HCG test is negative before starting Natalizumab infusion.
6. Do not have to wait for results of CBC with differential and/or BMP before starting natalizumab infusion.
7. VITAL SIGNS – Monitor and record vital signs, tolerance, and presence of infusion-related reactions prior to infusion, at end of infusion, 1 hour after infusion prior to discharge, and as clinically indicated.
8. Encourage patient to follow-up with the prescriber every 3 months.

HYPERSENSITIVITY MEDICATIONS:

1. NURSING COMMUNICATION – If hypersensitivity or infusion reactions develop, temporarily hold the infusion and notify provider immediately. Administer emergency medications per the Treatment Algorithm for Acute Infusion Reaction (Policy HC-PAT-133-GUD). Refer to algorithm for symptom monitoring and continuously assess as grade of severity may progress.
2. diphenhydramine (BENADRYL) injection, 25-50 mg, intravenous, AS NEEDED x1 dose for hypersensitivity reaction
3. EPINEPHrine HCl (ADRENALIN) injection, 0.3 mg, intramuscular, AS NEEDED x1 dose for hypersensitivity reaction
4. hydrocortisone sodium succinate (SOLU-CORTEF) injection, 100 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity reaction

AS NEEDED MEDICATIONS:

1. albuterol (PROVENTIL, VENTOLIN) 90 mcg/actuation inhaler, 2-4 puffs, every 10 Minutes AS NEEDED for bronchospasm
2. sodium chloride 0.9%, 1000 mL, intravenous, AS NEEDED, Infuse at 100-200 mL/hour, when natalizumab infusion is stopped for infusion reaction



Oregon Health & Science University
Hospital and Clinics Provider's Orders

ADULT AMBULATORY INFUSION ORDER

Natalizumab (TYSABRI)

Page 3 of 3

ACCOUNT NO.
MED. REC. NO.
NAME
BIRTHDATE

Patient Identification

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.

By signing below, I represent the following:

I am responsible for the care of the patient (*who is identified at the top of this form*);

I hold an active, unrestricted license to practice medicine in: Oregon _____ (*check box that corresponds with state where you provide care to patient and where you are currently licensed. Specify state if not Oregon*);

My physician license Number is # _____ (MUST BE COMPLETED TO BE A VALID PRESCRIPTION); and I am acting within my scope of practice and authorized by law to order Infusion of the medication described above for the patient identified on this form.

Provider signature: _____ **Date/Time:** _____

Printed Name: _____ **Phone:** _____ **Fax:** _____

OLC Central Intake Nurse:

Phone: 971-262-9645 (providers only) Fax: 503-346-8058

Please check the appropriate box for the patient's preferred clinic location:

Beaverton

OHSU Knight Cancer Institute
15700 SW Greystone Court
Beaverton, OR 97006
Phone number: 971-262-9000
Fax number: 503-346-8058

NW Portland

Legacy Good Samaritan campus
Medical Office Building 3, Suite 150
1130 NW 22nd Ave.
Portland, OR 97210
Phone number: 971-262-9600
Fax number: 503-346-8058

Gresham

Legacy Mount Hood campus
Medical Office Building 3, Suite 140
24988 SE Stark
Gresham, OR 97030
Phone number: 971-262-9500
Fax number: 503-346-8058

Tualatin

Legacy Meridian Park campus
Medical Office Building 2, Suite 140
19260 SW 65th Ave.
Tualatin, OR 97062
Phone number: 971-262-9700
Fax number: 503-346-8058

Infusion orders located at: www.ohsuknight.com/infusionorders