
 <p><b>Oregon Health &amp; Science University Hospital and Clinics Provider's Orders</b></p> <p style="font-size: small;">PO7071</p>  <p style="text-align: center;">ADULT AMBULATORY INFUSION ORDER <b>Ravulizumab-cwvz (ULTOMIRIS) Infusion</b></p> <p style="text-align: center; font-size: x-small;">Page 1 of 3</p>	<p>ACCOUNT NO. _____</p> <p>MED. REC. NO. _____</p> <p>NAME _____</p> <p>BIRTHDATE _____</p> <p style="text-align: right; font-size: x-small;"><i>Patient Identification</i></p>
<b>ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.</b>	

**Weight:** \_\_\_\_\_ kg      **Height:** \_\_\_\_\_ cm

**Allergies:** \_\_\_\_\_

**Diagnosis Code:** \_\_\_\_\_

**Treatment Start Date:** \_\_\_\_\_      **Patient to follow up with provider on date:** \_\_\_\_\_

**\*\*This plan will expire after 365 days at which time a new order will need to be placed\*\***

**GUIDELINES FOR ORDERING**

1. Send **FACE SHEET and H&P or most recent chart note.**
2. Ravulizumab-cwvz is part of FDA REMS Program
  - a. Providers **MUST** be enrolled in the Ultomiris REMS program.
  - b. Counsel patients using the Ultomiris patient safety card and patient safety brochure. Patients should carry the Ultomiris patient safety card at all times.
  - c. Please see reference links below for enrollment forms and additional help
    - i. <https://ultomirisrems.com/>
    - ii. [https://www.accessdata.fda.gov/drugsatfda\\_docs/rems/Ultomiris\\_2018\\_12\\_21\\_Prescriber\\_Enrollment\\_Form.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/rems/Ultomiris_2018_12_21_Prescriber_Enrollment_Form.pdf)
    - iii. [https://www.accessdata.fda.gov/drugsatfda\\_docs/rems/Ultomiris\\_2018\\_12\\_21\\_Prescriber\\_Safety\\_Brochure.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/rems/Ultomiris_2018_12_21_Prescriber_Safety_Brochure.pdf)
    - iv. [https://www.accessdata.fda.gov/drugsatfda\\_docs/rems/Ultomiris\\_2018\\_12\\_21\\_Patient\\_Safety\\_Brochure.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/rems/Ultomiris_2018_12_21_Patient_Safety_Brochure.pdf)
    - v. [https://www.accessdata.fda.gov/drugsatfda\\_docs/rems/Ultomiris\\_2018\\_12\\_21\\_Patient\\_Safety\\_Card.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/rems/Ultomiris_2018_12_21_Patient_Safety_Card.pdf)
3. Patients must receive meningococcal vaccines at least 2 weeks prior to treatment initiation; revaccinate according to current guidelines.
4. For patients switching from eculizumab to ravulizumab-cwvz, administer ravulizumab-cwvz loading dose 2 weeks after the last eculizumab infusion, and then administer maintenance doses once every 8 weeks, starting 2 weeks after loading dose administration.
5. Closely monitor patients for early signs and symptoms of meningococcal infections and evaluate immediately if infection is suspected. If ravulizumab-cwvz is administered to patients with active systemic infections, monitor for signs and symptoms of worsening infection.
6. Monitor patient after discontinuation for at least 16 weeks for signs and symptoms of hemolysis.
7. Consider penicillin prophylaxis for the duration of ravulizumab-cwvz therapy to potentially reduce the risk of meningococcal disease.

**PRE-SCREENING: (Results must be available prior to initiation of therapy):**

Meningococcal polysaccharide vaccines given on (dates) \_\_\_\_\_

**LABS:**

- CBC with differential, Routine, ONCE, every visit
- LDH Total, routine, ONCE, every visit
- Labs already drawn. Date: \_\_\_\_\_



Oregon Health & Science University  
Hospital and Clinics Provider's Orders

ADULT AMBULATORY INFUSION ORDER  
**Ravulizumab-cwvz (ULTOMIRIS)**  
Infusion

Page 2 of 3

ACCOUNT NO.  
MED. REC. NO.  
NAME  
BIRTHDATE

*Patient Identification*

**ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.**

**MEDICATION: Dose is based on weight at time of treatment (must check one)**

**Loading Dose:**

ravulizumab-cwvz (ULTOMIRIS) in sodium chloride 0.9%, intravenous, ONCE

- Patient weight 40-59.9 kg  **2400 mg over 2 hours**  
Patient weight 60-99.9 kg  **2700 mg over 2 hours**  
Patient weight 100 kg or greater  **3000 mg over 2 hours**

**Maintenance Doses:**

ravulizumab-cwvz (ULTOMIRIS) in sodium chloride 0.9%, intravenous, ONCE, every visit

- Patient weight 40-59.9 kg  **3000 mg over 2.5 hours**  
Patient weight 60-99.9 kg  **3300 mg over 2 hours**  
Patient weight 100 kg or greater  **3600 mg over 2.5 hours**

**Interval:**

- Every 8 weeks beginning 2 weeks after loading dose  
 Every 8 weeks beginning on date \_\_\_\_\_

**NURSING ORDERS:**

1. VITAL SIGNS – Monitor and record vital signs, tolerance, and presence of infusion-related reactions prior to infusion and every 15 minutes throughout infusion.
2. Observe for 1 hour after infusion complete (Unless the prescriber indicates this is not necessary).
3. Follow facility policies and/or protocols for vascular access maintenance with appropriate flush solution, declotting (alteplase), and/or dressing changes.

**HYPERSENSITIVITY MEDICATIONS:**

1. NURSING COMMUNICATION – If hypersensitivity or infusion reactions develop, temporarily hold the infusion and notify provider immediately. Administer emergency medications per the Treatment Algorithm for Acute Infusion Reaction (Policy HC-PAT-133-GUD). Refer to algorithm for symptom monitoring and continuously assess as grade of severity may progress.
2. diphenhydrAMINE (BENADRYL) injection, 25-50 mg, intravenous, AS NEEDED x1 dose for hypersensitivity reaction
3. EPINEPHrine HCl (ADRENALIN) injection, 0.3 mg, intramuscular, AS NEEDED x1 dose for hypersensitivity reaction
4. hydrocortisone sodium succinate (SOLU-CORTEF) injection, 100 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity reaction
5. famotidine (PEPCID) injection, 20 mg, intravenous, AS NEEDED x1 dose for hypersensitivity reaction



Oregon Health & Science University  
Hospital and Clinics Provider's Orders

ADULT AMBULATORY INFUSION ORDER  
**Ravulizumab-cwvz (ULTOMIRIS)**  
Infusion

Page 3 of 3

ACCOUNT NO.  
MED. REC. NO.  
NAME  
BIRTHDATE

*Patient Identification*

**ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.**

**By signing below, I represent the following:**

I am responsible for the care of the patient (*who is identified at the top of this form*);

I hold an active, unrestricted license to practice medicine in:  Oregon  \_\_\_\_\_ (*check box that corresponds with state where you provide care to patient and where you are currently licensed. Specify state if not Oregon*);

**My physician license Number is # \_\_\_\_\_ (MUST BE COMPLETED TO BE A VALID PRESCRIPTION);** and I am acting within my scope of practice and authorized by law to order Infusion of the medication described above for the patient identified on this form.

Provider signature: \_\_\_\_\_ Date/Time: \_\_\_\_\_

Printed Name: \_\_\_\_\_ Phone: \_\_\_\_\_ Fax: \_\_\_\_\_

OLC Central Intake Nurse:

Phone: 971-262-9645 (providers only) Fax: 503-346-8058

**Please check the appropriate box for the patient's preferred clinic location:**

**Beaverton**

OHSU Knight Cancer Institute  
15700 SW Greystone Court  
Beaverton, OR 97006  
Phone number: 971-262-9000  
Fax number: 503-346-8058

**NW Portland**

Legacy Good Samaritan campus  
Medical Office Building 3, Suite 150  
1130 NW 22nd Ave.  
Portland, OR 97210  
Phone number: 971-262-9600  
Fax number: 503-346-8058

**Gresham**

Legacy Mount Hood campus  
Medical Office Building 3, Suite 140  
24988 SE Stark  
Gresham, OR 97030  
Phone number: 971-262-9500  
Fax number: 503-346-8058

**Tualatin**

Legacy Meridian Park campus  
Medical Office Building 2, Suite 140  
19260 SW 65th Ave.  
Tualatin, OR 97062  
Phone number: 971-262-9700  
Fax number: 503-346-8058

Infusion orders located at: [www.ohsuknight.com/infusionorders](http://www.ohsuknight.com/infusionorders)