

General Screenings and Immunizations

For Males Ages 18-49* at Average Risk of Most Diseases

Screening Tests	When	Why
General Health:		
Full checkup, including height and weight 	Every 2-3 years. Discuss with your doctor or nurse.	To screen for diseases, assess risk for future problems, discuss lifestyle habits, and keep vaccinations up to date.
Thyroid Test (TSH)	Discuss with your doctor or nurse.	To identify an under- or overactive- thyroid, both of which are very treatable and either of which can lead to more serious conditions if left untreated.
Heart Health:		
Blood Pressure Test 	At least every 2 years.	To test for high blood pressure, which can lead to heart attack, stroke, heart failure, and kidney and eye problems.
Cholesterol Test	Every 5 years starting at age 35. If you smoke or are overweight, you may need to start at age 20. Discuss with doctor or nurse about when you should begin.	Treating cholesterol abnormalities can help reduce your risk of heart disease.

Screening Tests	When	Why
<p>Diabetes:</p>		
<p>Fasting Blood Sugar (Glucose) Test</p> 	<p>Discuss with your doctor or nurse.</p>	<p>To provide an early warning sign of high blood sugar levels, which could mean an increased risk for diabetes.</p>
<p>Reproductive Health:</p>		
<p>Testicular Exam</p>	<p>Starting at age 18, males should get a clinical testicular exam as part of their general checkup. Males should perform testicular self-exams once a month. Talk with your doctor or nurse.</p>	<p>To identify subtle changes in the testicles that could help detect testicular cancer early, when it's easy to treat.</p>
<p>Chlamydia Test</p> 	<p>Yearly until age 25 if sexually active. For ages 26 and over, get the test if you have new or multiple sexual partners, or at higher risk.</p>	<p>Prevents spread of chlamydia.</p>
<p>Sexually Transmitted Illnesses (STI) Tests</p>	<p>All sexually active males and their partners should be tested for STIs, including HIV, before beginning sexual activity.</p>	<p>Prevents spread of HIV and other STIs, many of which can only be detected through testing.</p>

Screening Tests	When	Why
Eye & Ear Health:		
<p data-bbox="272 296 440 331">Eye Exam</p> 	<p data-bbox="597 296 1008 638">At least once from ages 20-29; at least twice between ages 30-39. Get a baseline eye disease screening at age 40. Discuss your eye health with a doctor.</p>	<p data-bbox="1052 296 1479 533">To test your vision and screen for glaucoma and macular degeneration, two common, often age-related conditions.</p>
<p data-bbox="253 684 459 720">Hearing Test</p>	<p data-bbox="597 684 1008 768">Every 10 years from ages 18-50.</p>	<p data-bbox="1052 684 1474 768">To make sure you're hearing all life has to offer.</p>
Oral Health:		
<p data-bbox="253 884 459 919">Dental Exam</p> 	<p data-bbox="597 884 932 919">1-2 times every year.</p>	<p data-bbox="1052 884 1468 1121">To remove plaque and bacteria that could lead to tooth and gum disease; to check for tongue and mouth cancer.</p>
Skin Health:		
<p data-bbox="269 1251 443 1287">Skin Exam</p> 	<p data-bbox="597 1251 1024 1560">Males should do a monthly skin self-exam. Speak to your doctor or nurse about how often to have an additional skin exam by your doctor.</p>	<p data-bbox="1052 1251 1474 1497">To track worrisome moles, identify skin cancer early, and be able to report any changes to your doctor or nurse.</p>
Mental Health:		
<p data-bbox="159 1665 553 1701">Mental Health Screening</p> 	<p data-bbox="597 1665 987 1749">Discuss with your doctor or nurse.</p>	<p data-bbox="1052 1665 1458 1854">If you are feeling sad or anxious for a long time, it might be helpful to speak with a doctor.</p>

Screening Tests	When	Why
Immunizations:		
Influenza (Flu) Vaccine	Annually for everyone 6 months and older.	Protects against some flu viruses.
Human Papillomavirus Vaccine (HPV)	Males ages 11-12, or through age 26 if not previously vaccinated; 3 doses at 0-, 2-, and 6-month intervals; no booster necessary.	Protects against four common types of HPV, including those likely to cause cancers of the cervix, anus, vagina, penis, and throat.
Meningococcal Vaccine	College freshmen, military recruits, and other at-risk persons; discuss with your doctor or nurse.	Protects against some types of meningococcal disease (meningitis).
Tetanus, Diphtheria, Pertussis Booster Vaccine (Td/Tdap)	Every 10 years.	Protects against tetanus, diphtheria, and pertussis.
Varicella (Chickenpox) 	Given in 2 doses at 0- and 4- to 8-week intervals to those 19 or older who have not been vaccinated or had chickenpox.	Protects against chickenpox, a usually mild but highly contagious childhood disease which can be dangerous to infants and adults.

*** Please Note: These charts are guidelines only. Your doctor or nurse will personalize the timing of each test and immunization to best meet your health care needs.**