



# Similar Neural Pathways Control Foraging in Mosquitoes and Worms

Jogender Singh,<sup>a</sup> Alejandro Aballay<sup>a</sup>

<sup>a</sup>Department of Molecular Microbiology & Immunology, Oregon Health & Science University, Portland, Oregon, USA

**ABSTRACT** Female *Aedes aegypti* mosquitoes bite human hosts to obtain a blood meal and, in the process, act as vectors for many disease-causing viruses, including the dengue, chikungunya, yellow fever, and Zika viruses. After a complete meal, the female mosquitoes lose attraction to their hosts for several days. New research shows that pharmacological activation of neuropeptide Y-like receptor (NPYLR) signaling elicits host aversion in female mosquitoes. This behavior of mosquitoes shows remarkable similarities to a bacterial-aversion behavior of the nematode *Caenorhabditis elegans*. Feeding on pathogenic bacteria causes bloating of the gut in *C. elegans* that leads to activation of NPYLR signaling and bacterial aversion. Several studies suggest that this newly discovered mechanism underlying foraging may be conserved across a large number of species. A better understanding of the regulation of NPYLR signaling pathways could provide molecular targets for the control of eating behaviors in different animals, including human-disease vectors.

**KEYWORDS** *A. aegypti*, *C. elegans*, NPR-1, aversion behavior, bacterial colonization, feeding, neuropeptide Y

The *Aedes aegypti* mosquito is the primary vector for many disease-causing viruses, including the dengue, chikungunya, yellow fever, and Zika viruses. The female mosquitoes require blood from a host for the development of eggs and to complete their full reproductive life cycle (1). In the absence of a blood meal, the mosquitoes fail to develop eggs and do not reproduce. Therefore, the female mosquitoes possess a strong host-seeking behavior for a blood meal that depends primarily on sensing carbon dioxide (CO<sub>2</sub>) (1, 2). A single female goes through multiple blood-feeding and egg-laying cycles in her lifetime. Due to the cyclic nature of their feeding behavior, the female mosquitoes serve as effective carriers of disease-causing viruses from infected humans to healthy individuals. Thus, the suppression of the host-seeking behavior or elicitation of host aversion has the potential of being a practical method for reducing the spread of disease-causing viruses. In a recent study, Duvall et al. (3) reported that pharmacological activation of G protein-coupled neuropeptide Y (NPY)-related signaling elicits a host aversion behavior.

Extensive studies have been carried out to understand the blood-feeding behavior of mosquitoes, and they showed that host aversion consists of at least two phases: an early phase involving gut distension from a blood meal (4) and a sustained phase that lasts until the female lays her eggs (5, 6). Previous studies showed that a blood meal to repletion led to enhanced levels of neuropeptides in the hemolymph (7). Injection of hemolymph from blood-fed females or high doses of synthetic peptides that activate NPY-like receptors were sufficient to elicit host aversion in non-blood-fed females (6, 7). These findings suggested that a humoral response involving neuropeptides and activation of an NPY-related pathway plays a role in sustained host aversion in fully fed *A. aegypti* mosquitoes.

NPY-related signaling pathways are evolutionarily conserved and have been impli-


**Citation** Singh J, Aballay A. 2019. Similar neural pathways control foraging in mosquitoes and worms. *mBio* 10:e00656-19. <https://doi.org/10.1128/mBio.00656-19>.

**Editor** Frederick M. Ausubel, Mass General Hospital

**Copyright** © 2019 Singh and Aballay. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Alejandro Aballay, [aballay@ohsu.edu](mailto:aballay@ohsu.edu).

**Published** 30 April 2019



**FIG 1** A blood meal to repletion and gut distension caused by bacterial colonization lead to activation of NPY-related signaling in female *A. aegypti* mosquitoes and *C. elegans*, respectively. The increased NPY-related signaling is responsible for host aversion behavior in mosquitoes and bacterial-aversion behavior in *C. elegans*.

cated in several biological processes, including foraging, neuronal excitability, stress response, and motivated feeding behavior (8). Small-molecule drugs that possess high affinity to NPY human receptors might bind to mosquito NPY-like receptors and serve as powerful tools to investigate the functional role of NPY-related signaling in host-seeking behavior. Duvall and coworkers (3) performed a targeted drug screen and found that agonists for NPY-related signaling elicit a host aversion behavior even in non-blood-fed mosquitoes. Conversely, an antagonist for the NPY-related signaling elicited host-seeking behavior even in blood-fed mosquitoes. From a screen of all 49 predicted *A. aegypti* peptide receptors, the authors identified NPY-like receptor 7 (NPYLR7) as the sole target of these drugs. In addition, the investigators identified 9 peptide ligands for NPYLR7, including several Phe-Met-Arg-Phe amides (FMRFamides). These results showed that modulating NPYLR7 activity might affect host-seeking behavior. To identify *A. aegypti*-specific agonists with no cross-reactivity to human NPY receptors, the authors challenged NPYLR7 with 265,211 unique small molecules. The investigators isolated six highly selective NPYLR7 agonists that inhibit attraction to humans in wild-type mosquitos but not in *NPYLR7* null mutants. Finally, the investigators showed that these drugs were capable of inhibiting biting and blood feeding on a live host and, therefore, suggested a novel approach to control infectious disease transmission by controlling mosquito behavior.

This newly described mechanism for induction of host aversion in mosquitoes shows striking similarities to a bacterial-aversion behavior in the model nematode *Caenorhabditis elegans* (Fig. 1). *C. elegans* lives in anthropomorphic environments rich

in rotting vegetation and decaying fruit, where it is in contact with soilborne microbes, including bacteria that can be used as a food source. A recent study shows that gut distension caused by bacterial ingestion activates a *C. elegans* NPY-like receptor signal, involving NPR-1 and its FMRF-like peptide ligands, leading to bacterial aversion (9). Inhibition of the NPR-1 signaling pathway elicits avoidance of low CO<sub>2</sub> and high O<sub>2</sub>, while higher activity of the pathway induces avoidance of high CO<sub>2</sub> and low O<sub>2</sub> (10–12). Because bacterial lawns have high CO<sub>2</sub> and low O<sub>2</sub> due to bacterial metabolism, NPR-1 signaling integrates this information and helps in the elicitation of bacterial avoidance behavior in *C. elegans* (10, 13, 14). While mosquitoes seek hosts by their CO<sub>2</sub> levels and NPY-related signaling induces host aversion, it remains to be studied whether increased NPY-related signaling in blood-fed mosquitoes induces CO<sub>2</sub> avoidance by a mechanism similar to that of *C. elegans* (Fig. 1).

An important question that remains unanswered is how NPY-related signaling is activated in mosquitoes after a complete blood meal. In *C. elegans*, the activation of NPR-1 signaling was independent of colonization by live bacteria and depended on gut distension (9). Given the high similarities between the host aversion behavior of mosquitoes and bacterial-aversion behavior of *C. elegans*, it is possible that the NPY-related signaling is activated by similar mechanisms in mosquitoes. Indeed, gut distension is shown to be the reason for the early-phase induction of host aversion in mosquitoes. It will be interesting to study whether gut distension leads to the activation of the NPY-related signaling in mosquitoes.

Apart from mosquitoes and *C. elegans*, a variety of disparate species sense levels of CO<sub>2</sub> and/or O<sub>2</sub> in the environment and modulate their behavior accordingly (15). Other blood-feeding insects, such as black flies and tsetse flies, are attracted to CO<sub>2</sub> and use this signal to seek their human hosts. Similarly, the infective juveniles of the parasitic nematodes *Heterorhabditis bacteriophora* and *Steinernema carpocapsae* are attracted to CO<sub>2</sub> and use this cue for host seeking (16). CO<sub>2</sub> may also function as an alarm signal, as CO<sub>2</sub> emitted by stressed *Drosophila* acts as a signal for other *Drosophila* flies to flee (17). The hawkmoth *Manduca sexta* prefers flowers that emit a high level of CO<sub>2</sub>, suggesting that CO<sub>2</sub> acts as a proximal signal for nectar (18). Given that CO<sub>2</sub> and/or O<sub>2</sub> levels in the environment modulate the behavior of different organisms, it will be important to study whether the molecular mechanisms of sensing CO<sub>2</sub> and O<sub>2</sub> are conserved across species. The sensing mechanism of CO<sub>2</sub> and O<sub>2</sub> are relatively well studied in the model organisms *C. elegans* and *Drosophila* (15). This knowledge might act as a primer to expedite the understanding of the underlying molecular mechanisms in disease vectors and parasites.

## ACKNOWLEDGMENTS

The work in the Aballay laboratory is supported by grants from the National Institutes of Health (GM0709077 and AI117911).

## REFERENCES

1. Takken W, Verhulst NO. 2013. Host preferences of blood-feeding mosquitoes. *Annu Rev Entomol* 58:433–453. <https://doi.org/10.1146/annurev-ento-120811-153618>.
2. McMeniman CJ, Corfas RA, Matthews BJ, Ritchie SA, Vosshall LB. 2014. Multimodal integration of carbon dioxide and other sensory cues drives mosquito attraction to humans. *Cell* 156:1060–1071. <https://doi.org/10.1016/j.cell.2013.12.044>.
3. Duvall LB, Ramos-Espiritu L, Barsoum KE, Glickman JF, Vosshall LB. 2019. Small-molecule agonists of *Ae. aegypti* neuropeptide Y receptor block mosquito biting. *Cell* 176:687–701. <https://doi.org/10.1016/j.cell.2018.12.004>.
4. Klowden MJ, Lea AO. 1979. Abdominal distention terminates subsequent host-seeking behaviour of *Aedes aegypti* following a blood meal. *J Insect Physiol* 25:583–585. [https://doi.org/10.1016/0022-1910\(79\)90073-8](https://doi.org/10.1016/0022-1910(79)90073-8).
5. Klowden MJ. 1981. Initiation and termination of host-seeking inhibition in *Aedes aegypti* during oocyte maturation. *J Insect Physiol* 27:799–803. [https://doi.org/10.1016/0022-1910\(81\)90071-8](https://doi.org/10.1016/0022-1910(81)90071-8).
6. Liesch J, Bellani LL, Vosshall LB. 2013. Functional and genetic characterization of neuropeptide Y-like receptors in *Aedes aegypti*. *PLoS Negl Trop Dis* 7:e2486. <https://doi.org/10.1371/journal.pntd.0002486>.
7. Brown MR, Klowden MJ, Crim JW, Young L, Shrouder LA, Lea AO. 1994. Endogenous regulation of mosquito host-seeking behavior by a neuropeptide. *J Insect Physiol* 40:399–406. [https://doi.org/10.1016/0022-1910\(94\)90158-9](https://doi.org/10.1016/0022-1910(94)90158-9).
8. Brothers SP, Wahlestedt C. 2010. Therapeutic potential of neuropeptide Y (NPY) receptor ligands. *EMBO Mol Med* 2:429–439. <https://doi.org/10.1002/emmm.201000100>.
9. Singh J, Aballay A. 28 February 2019. Microbial colonization activates an immune fight-and-flight response via neuroendocrine signaling. *Dev Cell* <https://doi.org/10.1016/j.devcel.2019.02.001>.
10. Carrillo MA, Guillermin ML, Rengarajan S, Okubo RP, Hallem EA. 2013.

- O<sub>2</sub>-sensing neurons control CO<sub>2</sub> response in *C. elegans*. *J Neurosci* 33:9675–9683. <https://doi.org/10.1523/JNEUROSCI.4541-12.2013>.
11. Hallem EA, Sternberg PW. 2008. Acute carbon dioxide avoidance in *Caenorhabditis elegans*. *Proc Natl Acad Sci U S A* 105:8038–8043. <https://doi.org/10.1073/pnas.0707469105>.
  12. Bretscher AJ, Busch KE, de Bono M. 2008. A carbon dioxide avoidance behavior is integrated with responses to ambient oxygen and food in *Caenorhabditis elegans*. *Proc Natl Acad Sci U S A* 105:8044–8049. <https://doi.org/10.1073/pnas.0707607105>.
  13. Brandt JP, Ringstad N. 2015. Toll-like receptor signaling promotes development and function of sensory neurons required for a *C. elegans* pathogen-avoidance behavior. *Curr Biol* 25:2228–2237. <https://doi.org/10.1016/j.cub.2015.07.037>.
  14. Reddy KC, Hunter RC, Bhatla N, Newman DK, Kim DH. 2011. *Caenorhabditis elegans* NPR-1-mediated behaviors are suppressed in the presence of mucoïd bacteria. *Proc Natl Acad Sci U S A* 108:12887–12892. <https://doi.org/10.1073/pnas.1108265108>.
  15. Scott K. 2011. Out of thin air: sensory detection of oxygen and carbon dioxide. *Neuron* 69:194–202. <https://doi.org/10.1016/j.neuron.2010.12.018>.
  16. Hallem EA, Dillman AR, Hong AV, Zhang Y, Yano JM, Demarco SF, Sternberg PW. 2011. A sensory code for host seeking in parasitic nematodes. *Curr Biol* 21:377–383. <https://doi.org/10.1016/j.cub.2011.01.048>.
  17. Suh GSB, Wong AM, Hergarden AC, Wang JW, Simon AF, Benzer S, Axel R, Anderson DJ. 2004. A single population of olfactory sensory neurons mediates an innate avoidance behaviour in *Drosophila*. *Nature* 431:855–859. <https://doi.org/10.1038/nature02980>.
  18. Thom C, Guerenstein PG, Mechaber WL, Hildebrand JG. 2004. Floral CO<sub>2</sub> reveals flower profitability to moths. *J Chem Ecol* 30:1285–1288. <https://doi.org/10.1023/B:JOEC.0000030298.77377.7d>.