Oregon Health & Science University Hospital and Clinics Provider's Orders	ACCOUNT NO. MED. REC. NO. NAME BIRTHDATE			
Page 1 of 4				
ALL ORDERS MUST BE MARKED	IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.			
Weight:kg Height: Allergies:				
Diagnosis Code:				
Treatment Start Date: Patient to follow up with provider on date:				

This plan will expire after 365 days at which time a new order will need to be placed

GUIDELINES FOR ORDERING

1. Send FACE SHEET and H&P or most recent chart note.

LABS:

- □ CBC with differential, Routine, ONCE, every (visit)(days)(weeks)(months) Circle One
- □ CMP, Routine, ONCE, every_____(visit)(days)(weeks)(months) Circle One
- Labs already drawn. Date: _____

NURSING ORDERS:

1. Follow facility policies and/or protocols for vascular access maintenance with appropriate flush solution, declotting (alteplase), and/or dressing changes.

MEDICATIONS:

Penicillins:

ampicillin

- 1000 mg in sodium chloride 0.9%, 50 mL, intravenous, ONCE over 15-20 minutes
- □ 2000 mg in sodium chloride 0.9%, 100 mL, intravenous, ONCE over 15-20 minutes

nafcillin

- 1 gram in sodium chloride 0.9%, 100 mL, intravenous, ONCE over 30-60 minutes
- □ 2 grams in sodium chloride 0.9%, 100 mL, intravenous, ONCE over 30-60 minutes
- □ 12 grams over one day in sodium chloride 0.9%, 100.8 mL, intravenous, ONCE over 24 hours, continuous infusion via CADD (OHSU only)

oxacillin

- □ 1 gram in sodium chloride 0.9%, 100 mL, intravenous, ONCE over 30-60 minutes
- 2 grams in sodium chloride 0.9%, 100 mL, intravenous, ONCE over 30-60 minutes
- □ 12 grams over one day in sodium chloride 0.9%,100.8 mL, intravenous, ONCE over 24 hours, continuous infusion via CADD (OHSU only)

2			
	Oregon Health & Science University Hospital and Clinics Provider's Orders		
×		ACCOUNT NO.	
	ADULT AMBULATORY INFUSION ORDER Antibiotic Therapy (Penicillins and Carbapenems)	MED. REC. NO.	
		NAME	
		BIRTHDATE	
	Page 2 of 4	Patient Identification	
ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.			

penicillin G potassium (PFIZERPEN) intravenous

- 1 million units in sodium chloride 0.9%, 100 mL, ONCE over 1-2 hours
- □ 2 million units in sodium chloride 0.9%, 100 mL, ONCE over 1-2 hours
- □ 2.5 million units in sodium chloride 0.9%, 100 mL, ONCE over 1-2 hours
- □ 3 million units in sodium chloride 0.9%, 100 mL, ONCE over 1-2 hours
- □ 4 million units in sodium chloride 0.9%, 100 mL, ONCE over 1-2 hours
- □ 5 million units in sodium chloride 0.9%, 100 mL, ONCE over 1-2 hours
- 20 million over 1 day in sodium chloride 0.9%, 100.8 mL, ONCE over 24 hours, continuous infusion via CADD (OHSU only)

penicillin G benzathine (BICILLIN L-A) intramuscular

- □ 600,000 units as a single dose
- □ 1.2 million units as a single dose
- \Box 2.4 million units as a single dose

piperacillin/tazobactam (ZOSYN)

- □ 2.25 grams in sodium chloride 0.9%, 100 mL, intravenous, ONCE over 30 minutes
- □ 3.375 grams in sodium chloride 0.9%, 100 mL, intravenous, ONCE over 30 minutes
- 4.5 grams in sodium chloride 0.9%, 100 mL, intravenous, ONCE over 30 minutes
- 10.125 grams over one day in sodium chloride 0.9%, 112.8 mL intravenous, ONCE over 24 hours, continuous infusion via CADD (OHSU only)
- □ 13.5 grams over one day in sodium chloride 0.9%, 151.2 mL intravenous, ONCE over 24 hours, continuous infusion via CADD (OHSU only)
- □ 18 grams over one day in sodium chloride 0.9%, 201.6 mL intravenous, ONCE over 24 hours, continuous infusion via CADD (OHSU only)

Interval: (must check one)

□ Daily x _____ doses

Carbapenems:

ertapenem (IVANZ)

1 gram in sodium chloride 0.9%, 100 mL, intravenous, ONCE over 30 minutes

meropenem (MERREM)

- □ 500 mg in sodium chloride 0.9%, 100 mL, intravenous, ONCE over 15-30 minutes
- □ 1 gram in sodium chloride 0.9%, 100 mL, intravenous, ONCE over 15-30 minutes

Interval: (must check one)

- □ ONCE
- Daily x _____ doses

FOR InfuSystem[™] AMBULATORY PUMP USE (OHSU only; hook up at infusion location):

Duration:

□ _____ days

ONLINE 02/2023 [supersedes 10/2020]

Ø	Oregon Health & Science University Hospital and Clinics Provider's Orders	
OHSU Health	-	ACCOUNT NO.
	ADULT AMBULATORY INFUSION ORDER Antibiotic Therapy (Penicillins and Carbapenems)	MED. REC. NO.
		NAME
		BIRTHDATE
	Page 3 of 4	Patient Identification

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.

HYPERSENSITIVITY MEDICATIONS:

- NURSING COMMUNICATION If hypersensitivity or infusion reactions develop, temporarily hold the infusion and notify provider immediately. Administer emergency medications per the Treatment Algorithm for Acute Infusion Reaction (OHSU HC-PAT-133-GUD, HMC C-132). Refer to algorithm for symptom monitoring and continuously assess as grade of severity may progress.
- 2. diphenhydrAMINE (BENADRYL) injection, 25-50 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
- 3. EPINEPHrine HCI (ADRENALIN) injection, 0.3 mg, intramuscular, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
- 4. hydrocortisone sodium succinate (SOLU-CORTEF) injection, 100 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
- 5. famotidine (PEPCID) injection, 20 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction

By signing below, I represent the following:

I am responsible for the care of the patient (*who is identified at the top of this form*); I hold an active, unrestricted license to practice medicine in: *Oregon* (*check box that corresponds with state where you provide care to patient and where you are currently licensed. Specify state if not Oregon*);

My physician license Number is # ______ (MUST BE COMPLETED TO BE A VALID

PRESCRIPTION; and I am acting within my scope of practice and authorized by law to order Infusion of the medication described above for the patient identified on this form.

Provider signature:	Date/Time:	
Printed Name:	Phone:	Fax:

ACCOUNT NO.	
MED. REC. NO.	
NAME	
BIRTHDATE	

Page 4 of 4

(Penicillins and Carbapenems)

Patient Identification ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.

Central Intake:

Phone: 971-262-9645 (providers only) Fax: 503-346-8058

Please check the appropriate box for the patient's preferred clinic location:

□ Beaverton

□ Gresham

24988 SE Stark

Gresham, OR 97030

OHSU Knight Cancer Institute 15700 SW Greystone Court Beaverton, OR 97006 Phone number: 971-262-9000 Fax number: 503-346-8058

Legacy Mount Hood campus

Phone number: 971-262-9500

Fax number: 503-346-8058

Medical Office Building 3, Suite 140

□ NW Portland

Legacy Good Samaritan campus Medical Office Building 3, Suite 150 1130 NW 22nd Ave Portland, OR 97210 Phone number: 971-262-9600 Fax number: 503-346-8058

□ Tualatin

Legacy Meridian Park campus Medical Office Building 2, Suite 140 19260 SW 65th Ave Tualatin, OR 97062 Phone number: 971-262-9700 Fax number: 503-346-8058

Infusion orders located at: www.ohsuknight.com/infusionorders