



Oregon Health & Science University  
Hospital and Clinics Provider's Orders

PO7071



ADULT AMBULATORY INFUSION ORDER  
**Certolizumab (CIMZIA) Injection**

Page 1 of 3

ACCOUNT NO.  
MED. REC. NO.  
NAME  
BIRTHDATE

*Patient Identification*

**ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.**

Weight: \_\_\_\_\_ kg      Height: \_\_\_\_\_ cm

Allergies: \_\_\_\_\_

Diagnosis Code: \_\_\_\_\_

Treatment Start Date: \_\_\_\_\_ Patient to follow up with provider on date: \_\_\_\_\_

**\*\*This plan will expire after 365 days at which time a new order will need to be placed\*\***

**GUIDELINES FOR ORDERING**

1. Send **FACE SHEET and H&P or most recent chart note.**
2. Hepatitis B (Hep B surface antigen and core antibody total) screening must be completed prior to initiation of treatment and the patient should not be infected. Please send results with order.
3. A Tuberculin test must have been placed and read as negative prior to initiation of treatment (PPD or QuantiFERON Gold blood test). Please send results with order. If result is indeterminate, a follow up chest X-ray must be performed to rule out TB. Please send results with order.

**PRE-SCREENING: (Results must be available prior to initiation of therapy):**

- Hepatitis B **surface antigen** and **core antibody** test results included with orders
- Tuberculin skin test or QuantiFERON Gold test results included with orders
- Chest X-Ray result scanned with orders if TB test result is indeterminate.

**LABS:**

- CBC with differential, Routine, ONCE, every \_\_\_\_\_ (visit)(days)(weeks)(months) – *Circle One*

**NURSING ORDERS:**

1. TREATMENT PARAMETER – Hold treatment and contact provider if Hepatitis B surface antigen or core antibody total test result is positive, TB test result is positive, or if screening has not been performed.
2. Administer 400 mg dose as 2 divided doses subcutaneously using provided 23-gauge needles to separate sites on the abdomen or thigh. Rotate injection sites. Do not administer to areas where skin is tender, bruised, red, or hard
3. Follow facility policies and/or protocols for vascular access maintenance with appropriate flush solution, dec clotting (alteplase), and/or dressing changes.

**MEDICATIONS:**

**certolizumab (CIMZIA), subcutaneous, ONCE**

**Initial Dose:**

- 400 mg for 3 doses on week 0: \_\_\_\_\_, week 2: \_\_\_\_\_, week 4: \_\_\_\_\_

**Maintenance Doses:**

- 400 mg every 4 weeks beginning week 8: \_\_\_\_\_
- 200 mg every 2 weeks beginning week 6: \_\_\_\_\_



Oregon Health & Science University  
Hospital and Clinics Provider's Orders

ADULT AMBULATORY INFUSION ORDER  
**Certolizumab (CIMZIA) Injection**

Page 2 of 3

ACCOUNT NO.  
MED. REC. NO.  
NAME  
BIRTHDATE

*Patient Identification*

**ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.**

**HYPERSENSITIVITY MEDICATIONS:**

1. NURSING COMMUNICATION – If hypersensitivity or infusion reactions develop, temporarily hold the infusion and notify provider immediately. Administer emergency medications per the Treatment Algorithm for Acute Infusion Reaction (OHSU HC-PAT-133-GUD, HMC C-132). Refer to algorithm for symptom monitoring and continuously assess as grade of severity may progress.
2. diphenhydrAMINE (BENADRYL) injection, 25-50 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
3. EPINEPHrine HCl (ADRENALIN) injection, 0.3 mg, intramuscular, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
4. hydrocortisone sodium succinate (SOLU-CORTEF) injection, 100 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
5. famotidine (PEPCID) injection, 20 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction

**By signing below, I represent the following:**

I am responsible for the care of the patient (*who is identified at the top of this form*);

I hold an active, unrestricted license to practice medicine in:  Oregon  \_\_\_\_\_ (*check box that corresponds with state where you provide care to patient and where you are currently licensed. Specify state if not Oregon*);

**My physician license Number is # \_\_\_\_\_ (MUST BE COMPLETED TO BE A VALID PRESCRIPTION);** and I am acting within my scope of practice and authorized by law to order Infusion of the medication described above for the patient identified on this form.

Provider signature: \_\_\_\_\_ Date/Time: \_\_\_\_\_

Printed Name: \_\_\_\_\_ Phone: \_\_\_\_\_ Fax: \_\_\_\_\_



Oregon Health & Science University  
Hospital and Clinics Provider's Orders

ADULT AMBULATORY INFUSION ORDER  
**Certolizumab (CIMZIA) Injection**

Page 3 of 3

ACCOUNT NO.  
MED. REC. NO.  
NAME  
BIRTHDATE

*Patient Identification*

**ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.**

Central Intake:

Phone: 971-262-9645 (providers only) Fax: 503-346-8058

***Please check the appropriate box for the patient's preferred clinic location:***

**Beaverton**

OHSU Knight Cancer Institute  
15700 SW Greystone Court  
Beaverton, OR 97006

Phone number: 971-262-9000

Fax number: 503-346-8058

**NW Portland**

Legacy Good Samaritan campus  
Medical Office Building 3, Suite 150  
1130 NW 22nd Ave.  
Portland, OR 97210

Phone number: 971-262-9600

Fax number: 503-346-8058

**Gresham**

Legacy Mount Hood campus  
Medical Office Building 3, Suite 140  
24988 SE Stark  
Gresham, OR 97030

Phone number: 971-262-9500

Fax number: 503-346-8058

**Tualatin**

Legacy Meridian Park campus  
Medical Office Building 2, Suite 140  
19260 SW 65th Ave.  
Tualatin, OR 97062

Phone number: 971-262-9700

Fax number: 503-346-8058

Infusion orders located at: [www.ohsuknight.com/infusionorders](http://www.ohsuknight.com/infusionorders)