

Weight:

ADULT AMBULATORY INFUSION ORDER Iron Sucrose (VENOFER) Infusion

Page 1 of 3

Height:

ka

ACCOUNT NO.
MED. REC. NO.
NAME
BIRTHDATE

Patient Identification

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.

cm

Allergi	rgies:		
	nosis Code:		
Treatm	tment Start Date: Patient t	Patient to follow up with provider on date:	
This	nis plan will expire after 365 days at whic	h time a new order will need to be placed	
1.	DELINES FOR ORDERING 1. Send FACE SHEET and H&P or most 2. Provider must order and obtain a ferritin date:	recent chart note. prior to patient being scheduled for iron infusion. Labs drawn	
1. 2.	Instruct patient to obtain ferritin lab 30 d with provider.	atment and notify provider if Ferritin greater than 300 ng/mL. ays after infusion treatment and set up follow up appointment for vascular access maintenance with appropriate flush solution,	
MEDIO	DICATIONS:	ianges.	
iro	iron sucrose (VENOFER): (must check one))	
<u> </u>	<u> </u>	_, intravenous, ONCE, over 30 minutes, x 5 doses over 14 days _, intravenous, ONCE, over 1.5 hours, x 3 doses	

AS NEEDED MEDICATIONS:

1. sodium chloride 0.9%, 500 mL, intravenous, AS NEEDED x1 dose for vein discomfort. Give concurrently with iron sucrose

HYPERSENSITIVITY MEDICATIONS:

- NURSING COMMUNICATION If hypersensitivity or infusion reactions develop, temporarily hold the
 infusion and notify provider immediately. Administer emergency medications per the Treatment
 Algorithm for Acute Infusion Reaction (OHSU HC-PAT-133-GUD, HMC C-132). Refer to algorithm for
 symptom monitoring and continuously assess as grade of severity may progress.
- 2. diphenhydrAMINE (BENADRYL) injection, 25-50 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
- 3. EPINEPHrine HCI (ADRENALIN) injection, 0.3 mg, intramuscular, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
- 4. hydrocortisone sodium succinate (SOLU-CORTEF) injection, 100 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction



Oregon Health & Science University Hospital and Clinics Provider's Orders

OHSU ADULT AMBULATORY INFUSION ORDER Health Iron Sucrose (VENOFER) Infusion

Page 2 of 3

ACCOUNT NO.
MED. REC. NO.
NAME
BIRTHDATE

Patient Identification

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.

5. famotidine (PEPCID) injection, 20 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction

By signing below, I represent the following: I am responsible for the care of the patient (who is I hold an active, unrestricted license to practice methat corresponds with state where you provide can	edicine in: 🛭 Oreg	gon □ (check box
state if not Oregon);	re to patient and wir	ere you are currently licensed. Specify
My physician license Number is #	oe of practice and au	
Provider signature:	Dat	e/Time:
Printed Name:	Phone:	Fax:



Oregon Health & Science University Hospital and Clinics Provider's Orders

OHSU ADULT AMBULATORY INFUSION ORDER Health Iron Sucrose (VENOFER) Infusion

Page 3 of 3

ACCOUNT NO.
MED. REC. NO.
NAME
BIRTHDATE

Patient Identification

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.

Central Intake:

Phone: 971-262-9645 (providers only) Fax: 503-346-8058

Please check the appropriate box for the patient's preferred clinic location:

□ Beaverton

OHSU Knight Cancer Institute 15700 SW Greystone Court Beaverton, OR 97006

Phone number: 971-262-9000 Fax number: 503-346-8058

☐ Gresham

Legacy Mount Hood campus Medical Office Building 3, Suite 140 24988 SE Stark Gresham, OR 97030

Phone number: 971-262-9500 Fax number: 503-346-8058

□ NW Portland

Legacy Good Samaritan campus Medical Office Building 3, Suite 150 1130 NW 22nd Ave Portland, OR 97210

Phone number: 971-262-9600 Fax number: 503-346-8058

□ Tualatin

Legacy Meridian Park campus Medical Office Building 2, Suite 140 19260 SW 65th Ave Tualatin, OR 97062

Phone number: 971-262-9700 Fax number: 503-346-8058

Infusion orders located at: www.ohsuknight.com/infusionorders