

Oregon Health & Science University **Hospital and Clinics Provider's Orders**



ADULT AMBULATORY INFUSION ORDER Iron Sucrose (VENOFER) Infusion

Page 1 of 2

ACCOUNT NO. MED. REC. NO. NAME **BIRTHDATE**

Patient Identification

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.

ight:kg Height:cm
ergies:
gnosis Code:
atment Start Date: Patient to follow up with provider on date:
his plan will expire after 365 days at which time a new order will need to be placed**
 Send FACE SHEET and H&P or most recent chart note. Provider must order and obtain a ferritin prior to patient being scheduled for iron infusion. Labs drawn date:
 RSING ORDERS: TREATMENT PARAMETER – Hold treatment and notify provider if Ferritin greater than 300 ng/mL. Instruct patient to set follow up appointment with provider for follow up labs. Follow facility policies and/or protocols for vascular access maintenance with appropriate flush solution, declotting (alteplase), and/or dressing changes.
iron sucrose (VENOFER): (must check one)
 200 mg in sodium chloride 0.9% 100 mL, intravenous, ONCE, over 30 minutes, x 5 doses over 14 days 300 mg in sodium chloride 0.9% 250 mL, intravenous, ONCE, over 1.5 hours, x 3 doses (administered every 2 to 3 days)
NEEDED MEDICATIONS:

1. sodium chloride 0.9%, 500 mL, intravenous, AS NEEDED x1 dose for vein discomfort. Give concurrently with iron sucrose

HYPERSENSITIVITY MEDICATIONS:

- 1. NURSING COMMUNICATION If hypersensitivity or infusion reactions develop, temporarily hold the infusion and notify provider immediately. Administer emergency medications per the Treatment Algorithm for Acute Infusion Reaction (OHSU HC-PAT-133-GUD, HMC C-132). Refer to algorithm for symptom monitoring and continuously assess as grade of severity may progress.
- 2. diphenhydrAMINE (BENADRYL) injection, 25-50 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
- 3. EPINEPHrine HCI (ADRENALIN) injection, 0.3 mg, intramuscular, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
- 4. hydrocortisone sodium succinate (SOLU-CORTEF) injection, 100 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
- 5. famotidine (PEPCID) injection, 20 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction



Oregon Health & Science University Hospital and Clinics Provider's Orders

OHSU ADULT AMBULATORY INFUSION ORDER Health Iron Sucrose (VENOFER) Infusion

Page 2 of 2

ACCOUNT NO.
MED. REC. NO.
NAME
BIRTHDATE

Patient Identification

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.

By signing below, I represent the following: I am responsible for the care of the patient (who is identified at the top of this form); I hold an active, unrestricted license to practice medicine in: Oregon (check box that corresponds with state where you provide care to patient and where you are currently licensed. Specify state if not Oregon); My physician license Number is # (MUST BE COMPLETED TO BE A VALID PRESCRIPTION); and I am acting within my scope of practice and authorized by law to order Infusion of the medication described above for the patient identified on this form.				
Provider signature:	Date/Time:			
Printed Name:	Phone:	Fax:		
<u>Central Intake:</u> Phone: 971-262-9645 (providers only) Fax: 50 Please check the appropriate box for the p		cation:		
□ Beaverton OHSU Knight Cancer Institute 15700 SW Greystone Court Beaverton, OR 97006 Phone number: 971-262-9000 Fax number: 503-346-8058	□ NW Portland Legacy Good S	amaritan campus Building 3, Suite 150 Ave 7210 <u>971-262-9600</u>		
☐ Gresham Legacy Mount Hood campus Medical Office Building 3, Suite 140 24988 SE Stark Gresham, OR 97030 Phone number: 971-262-9500 Fax number: 503-346-8058	□ Tualatin Legacy Meridial Medical Office E 19260 SW 65th Tualatin, OR 97 Phone number: Fax number: 50	Building 2, Suite 140 Ave 1062 1971-262-9700		

Infusion orders located at: www.ohsuknight.com/infusionorders