



You and your simultaneous liver and kidney transplant

A handbook for patients and their families

<u>General information</u>	<u>3</u>
Important contacts	3
What to do once you get home.....	5
<u>Labs and blood tests</u>	<u>9</u>
<u>Diet</u>	<u>15</u>
<u>Caring for your health</u>	<u>37</u>
Activity.....	37
Long-term health.....	40
Preventing infectious disease	43
Dental care	46
Drugs, tobacco and alcohol.....	46
Pets at home.....	47
<u>Complications</u>	<u>49</u>
<u>Writing to your donor's family</u>	<u>55</u>
<u>References</u>	<u>57</u>
Prep for your Kidney Transplant Biopsy	57
Diagnostic tests	59
OTC drugs safe to take	61
Magnesium in food.....	62
Phosphorus in food.....	63
Glossary.....	64
<u>Forms</u>	<u>79</u>
Lab Results	79
Vital Signs	85
<u>Medication Information Handout</u>	<u>Separate Handout</u>

This Page Intentionally Left Blank

General Information

Important contacts

My post-transplant liver coordinator: _____	503-494-8500 800-452-1369 ext. _____
My post-transplant kidney coordinator: _____	503-494- _____ 800-452-1369 ext. _____
OHSU transplant office <i>Weekdays 8 a.m. - 5 p.m.</i>	503-494-8500 800-452-1369 ext. 8500 Fax: 503-494-4492
Urgent issues DURING open hours <i>Weekdays 8 a.m. - 5 p.m.</i>	503-494-8500 <i>For the first 3 months after your transplant ask to speak with a post-transplant liver coordinator. Starting - _____ call and speak with your kidney coordinator.</i>
Urgent issues AFTER hours <i>Nights, weekends, holidays, etc.</i> EMERGENCIES ONLY	503-494-8311 800-452-1369 <i>Tell the operator you are a liver and kidney transplant patient and that you need to speak with the liver transplant provider on call. Please remember, the on call provider is for EMERGENCIES ONLY.</i>
OHSU Pharmacy	503-346-3370
OHSU Specialty Pharmacy	503-418-8228 866-770-2666
OHSU Mail Order Pharmacy	503-346-3370 503-748-7600
OHSU Laboratory	503-494-7383
OHSU MyChart	503-494-5252
Patient Financial Coordinator	503-494-5539
Diabetes help DURING open hours Harold Schnitzer Diabetes Center <i>For help with blood sugars/insulin Management</i>	503-494-3273 <i>If you need to leave a message, your message will be returned soon.</i>
Diabetes help AFTER office hours <i>Nights, weekends, holidays, etc.</i> For URGENT help with high or low blood sugars	503-494-8311 800-452-1369 <i>Ask to speak to the person on call for transplant</i>

Mailing address

OHSU Clinical Transplant Services

Liver Transplant Program

Mail code: L590

3181 S.W. Sam Jackson Park Road

Portland, Oregon 97239-3098

What to do once you get home

You may be worried about all the new things you need to know about your health. It is a normal feeling, and we don't expect you to know it all right away. That's why we created this guide. You can read through it, write down any questions and discuss them with your transplant team. Please **ALWAYS ask us questions if you do not understand something**. We are here to help you return to an active life in family, work and play.

Start following these 5 steps as soon as you get home from the hospital.

1. **Every day, write down your vital signs** in your book.

Vital sign	When	How	Call
Weight	Every morning after you empty your bladder.	Weigh yourself at around the same time, wearing the same amount of clothes, and using the same scale every day.	If you gain 4+ pounds in 1 day , call the kidney transplant nurse
Temperature	1+ times a day	If you feel like you have a fever or the chills, take your temperature more often.	For the first 3 months : call the transplant nurse if your fever is 100°F or more . 3+ months after transplant: call your local doctor if your fever is over 100°F .
Blood pressure	Once in the morning , and once in the early evening .	Do NOT take your blood pressure on the arm with a dialysis fistula or graft.	If your blood pressure is too high or too low .

2. **Take your medications:** The #1 reason why transplants fail is because patients do not follow their medication routine. It is very important to take all your medications as prescribed.

3. **Go to the lab on schedule** and learn what your lab results mean. Write down your lab results on the forms in the back of this handbook. As your kidney and liver function stabilizes, the frequency of blood tests will decrease. *You will be notified when your lab schedule changes.*

Routine lab schedule	
First month – all labs will be done the first month at OHSU	Monday, Thursday at OHSU
After 1 month	Monday, Thursday
After 3 months	Monday
After 6 months	Every other Monday
After 1 year	Every 3 months

4. **Go to your clinic on schedule.** For the first 4 weeks, you will go to the OHSU transplant clinic 1 time a week. You will go less often once you are stable. Bring this guide and any questions to your clinic visits. We will look over your surgical wound, remove your staples when ready and talk to you about any medication changes.
5. **Get your kidney surveillance:** Surveillance labs are routinely done at 3 and 12 months.

Transplant emergencies



Call 911 if you have any of these:

- **Chest pain**
- **Fainting** (loss of consciousness)
- **Bleeding** that won't stop
- **Signs of stroke**
- **Trouble breathing**

When to call the transplant nurse

Call if you have any of these:

- Fever of **over 100° F**
- Weight gain of **4 or more pounds in 1 day**
- **Creatinine** goes up by **0.2 or more**
- **Trouble breathing**, especially when laying down
- Problems with your **surgical cut area**
- **Ongoing diarrhea**
- **Trouble urinating**
- **Urinary infection symptoms**
 - Pain, burning or tenderness when you urinate
 - Needing to go more often
 - Cloudy or foul-smelling urine
- Nausea and vomiting — **if you can't keep your pills down**
- Pain that does not go away **after you take pain medicine**
- **Blood pressure** above 150/85
- Heartburn
- Tenderness or pain in **the area around your transplanted organs**

OHSU transplant

Weekdays

8 a.m. - 5 p.m.

503-494-8500

800-452-1369 ext.

8500

Emergencies

AFTER hours

503-494-8311

800-452-1369

Ask to speak to the provider on call for transplant.

This Page Intentionally Left Blank

Labs and blood tests

Blood tests are extremely important! They tell us if your liver or kidney is being rejected or if you are having side effects from the drugs.

6 things to know:

1. Have your blood drawn in the **morning** so you can get your results that day.
2. You must get your labs done **at OHSU for the first month** after transplant.
3. After the first month, you can start going to the lab of your choice. Our office will send lab orders to your new lab.
4. These orders will give the lab permission to release the results to you. They will also fax the results to our office.
5. You need to **keep track** of your blood test results.
6. Learn what the numbers mean and what is normal for you.

We will NOT call you if your transplant numbers are normal for you.

Drug level blood tests

If you take Tacrolimus or Cyclosporine*

On the day of your drug level blood test:

- Take your medication about **12 hours before** your blood draw. Your blood draw must be between 11 ½ and 12 ½ hours **AFTER** you take that dose.
- Do **NOT** take your next dose **until after** your blood draw.

If you take Rapamune*

On the day of your drug level blood test:

- Take your medication **20 - 24 hours before** your blood draw.
- Do **NOT** take your next dose **until after** your blood is draw.

*** If you cannot get your blood drawn within the right time frames:**

- ⊙ Do **NOT** get a drug level blood test, and
- **Tell your transplant coordinator.**

Labs

The OHSU lab

The lab is on the 3rd floor of OHSU Physicians Pavilion. There is also a lab at The Center for Health and Healing Building 2.

To get your test results:

- Call **503-494-7383 after 2 p.m.** on the day of your lab draw, or
- Sign up for OHSU **MyChart** to see your results online. Go to www.ohsu.edu/mychart.

Hours change on holidays and weekends, so be sure to check with the lab staff.

If you use a non-OHSU lab

You and your transplant team must watch your lab work closely.

Routine lab schedule	
First month – all labs will be done the first month at OHSU	Monday, Thursday at OHSU
After 1 month	Monday, Thursday
After 3 months	Monday
After 6 months	Every other Monday
After 1 year	Every 3 months

*All labs will be done at OHSU the first month post-transplant

*You will be notified when your lab schedule changes.

Normal blood test values

Every lab has slightly different normal ranges for blood tests. On this page, we've listed the normal values OHSU uses for patients who have not had a transplant. Please use them only as guidelines. Every transplant patient has their own normal.

We will help you understand your results and define what your baseline (starting point) is. Many times, labs will run other tests along with the ones we've listed.

Lab test	Normal range
Total Bilirubin (T Bili) <i>The level of bilirubin (yellowish substance that travels through your liver) in your blood. Too high could be a sign of an obstruction, rejection or infection.</i>	0.1 - 1.2
Alkaline Phosphatase (Alk Phos) <i>The level of an enzyme your liver makes. Could be a sign of the liver or other organs not working right.</i>	45 - 129
Alanine Transaminase (SGPT or ALT) <i>The level of an enzyme in your liver. Too high could be a sign of liver injury, rejection or obstruction.</i>	9 - 57
Aspartate Transaminase (SGOT or AST) <i>The level of an enzyme in many of your organs. Too high could be a sign of liver injury or rejection.</i>	14 - 44
Gamma Glutamyl Transferase (GGT) <i>The level of an enzyme found in your liver and other organs. Too high could be a sign of an obstruction, rejection or infection.</i>	5 - 59
Albumin (Alb) <i>The level of a protein made by your liver. Too low could be a sign of malnutrition.</i>	3.5 - 5.0
Total Protein (TP) <i>The level many proteins found in your blood. Too low could be a sign of your liver not working right or malnutrition</i>	6 - 8.4
Glucose (Blood Sugar) <i>The level of sugar (glucose) in your blood.</i>	71 - 109

Blood Urea Nitrogen (BUN) <i>The level of nitrogen (a waste product of protein use) in your blood.</i>	7 - 23
Creatinine (Cr) <i>The level of creatinine (a waste product of muscle metabolism) in your blood.</i>	0.6 - 1.1
Sodium (Na) <i>A mineral needed to keep body fluids in balance.</i>	131 - 142
Potassium (K) <i>A mineral needed for muscles to work properly. This includes the heart.</i>	3.5 - 5.0
Phosphorous (PO4) <i>A mineral that works with calcium to form bone. A small amount is found in the muscles.</i>	2.2 - 4.2
Magnesium (Mg) <i>A mineral needed for normal bodily function.</i>	2.0 - 2.6
Calcium (Ca++) <i>A mineral needed for properly working muscles, nerves, heart, blood clotting, strong bones and teeth.</i>	8.4 - 10.4
White Blood Count (WBC) <i>Special blood cells that help the body fight infection.</i>	4.4 - 10.8
Hematocrit (Hct) <i>The % of red blood cells in your blood. The red cells carry oxygen and carbon dioxide throughout the body.</i>	Men: 40 - 54% Women: 37 - 47%
Platelet (Plt) <i>Special blood cells that help your blood clot.</i>	150,000 - 350,000
Hemoglobin (Hgb) <i>Oxygen part of the red blood cell</i>	Men: 12 - 18 Women: 12 - 16

This Page Intentionally Left Blank

How to eat after your transplant

After a transplant, you have special nutritional needs. This handout will explain:

- What you should eat right after surgery
- Food safety, to keep you from getting sick
- What you should eat in the long-term.

What you should eat right after your transplant

Calories and protein

You need to eat extra calories and protein right after your transplant. Good nutrition helps your body:

- Heal your surgical wounds
- Keep muscle strength
- Fight infection

If you take prednisone, you need to eat more protein. This medication can cause muscle loss. Protein helps you build muscle. It is in foods like eggs, meat, fish, poultry, milk, cheese, yogurt, beans, nuts and seeds.


Right after transplant surgery:

- You should eat _____ grams of protein each day.
- You should eat _____ calories each day.

8 weeks after surgery:

- Lower the amount of protein you eat to _____ grams of protein each day.
- Your dietitian may change the amount of calories you should eat based on how much activity you get.

A healthy plate for the first 2 months after transplant



Use the plate method of eating, where $\frac{1}{2}$ your plate is protein, $\frac{1}{4}$ is non-starchy vegetables and $\frac{1}{4}$ is starchy food like whole grain bread, pasta, rice, cereal or starchy vegetables.

High-protein foods

Protein Food	Portion Size	Protein Grams
Meat, cooked:		
• Chicken breast	3 oz	26
• Hamburger, lean	3 oz, 1 patty	18
• Pork chop	3 oz	24
• Steak, lean	3 oz	26
Fish, cooked:		
• Fillet	3 oz	15
• Salmon patty	3 oz, 1 patty	15
Fish, canned:		
• Tuna or Salmon	1 oz	7
• Anchovies, sardines	1 oz	7
Shellfish, cooked:		
• Crab	1 oz	6
• Scallops, steamed	1 oz	7
• Shrimp	1 oz	6
Beans, cooked:		
• Kidney, black, garbanzo, pinto	½ cup	7
• Lentils, split peas	½ cup	8
Cheese:		
• Cottage cheese	¼ cup	6
• Soft (goat, mozzarella)	1 oz	5
• Hard (cheddar, Swiss)	1 oz	7

Protein Food	Portion Size	Protein Grams
• Ricotta	¼ cup	7
• String cheese	1 oz (1 stick)	7
Milk:		
• Whole, 2% 1%, skim or soy	1 cup	8
• Non-fat, powdered	¼ cup	10
Yogurt:		
• Greek	5.3 oz	12-18
• Regular	5.3 oz	5-6
Nuts and seeds:		
• Peanuts, almonds, sunflower	1 oz	6-7
• Cashews, flaxseed (ground), walnuts	1 oz	4-5
• Nut butters	2 Tablespoons	5-8
Protein drinks:		
• Ensure Clear, Glucerna	8 oz	8-10
• Isopure Plus	16 oz	20
• Boost High Protein, Boost Plus, Ensure Plus	8 oz	13-16
• Boost VHC, Nepro	8 oz	19-22
• Premier Protein Shakes	11 oz	30
• Carnation Instant Breakfast with 1 cup milk	8 oz	13

Protein Food	Portion Size	Protein Grams
Other:		
• Whey protein powder (protein depends on brand)	1 oz	12-20
• Egg	1 large	6
• Quinoa, cooked	1 cup	8
• Tempeh	1 oz	5
• Tofu (soft, firm, extra firm)	3 oz	5-9

How to eat more protein

- Eat 5-6 small meals and snacks during the day instead of 3 large meals. Try to have:
 - 20+ grams of protein at each meal
 - 8+ grams of protein at each snack
- Eat Greek yogurt instead of regular yogurt. Greek yogurt has 2 times the protein of regular yogurt.
- Add diced meat, beans, cheese, nuts or seeds to salads, casseroles or soups.
- Spread nut butters or hummus on low sodium crackers, toast, or raw fruits and vegetables.
- Keep ready-made protein-rich foods like hard-boiled eggs, cheese sticks, single- serving yogurts, unsalted nuts and trail mix on hand for easy snacks.
- Add a glass of milk or protein drink to your meals or as a snack.
- Add protein powder or powdered milk to casseroles, soups or oatmeal.
- Try cottage cheese or ricotta cheese with fruit, vegetables or mixed into casseroles.
- Make your own smoothie using milk, Greek yogurt, protein powders, nut butters or tofu.
- Eat whole grains such as quinoa and high protein cereals.

Fluids

How much to drink every day

- Your goal is to drink 2 liters of fluid every day. This equals 8-10 cups or 64-80 ounces.
- Please ask your doctor if they have a fluid goal for you.

How to drink more fluids

- Sip liquids slowly and pace yourself during the day.
- Drink even when you are not thirsty.
- Drink a small amount before, during and after meals and snacks.
- Carry a full water bottle wherever you go.
- Limit coffee, tea and soda that contain caffeine.
- ⊙ **Do NOT** drink energy drinks such as Red Bull, Monster or Rock Star.
- Set a timer or alarm to ring every 1 to 2 hours to remind you to drink fluids.
- Whenever you can, choose healthful drinks such as water, milk, unsweetened tea, low-sodium vegetable juices, smoothies, flavored or sparkling waters or nutritional supplement shakes.

Foods NOT to eat

Some foods may interfere with your immunosuppressant medications, especially tacrolimus (Prograf) and cyclosporine. For this reason, **do NOT eat**:

- ⊗ Grapefruit and grapefruit juice
- ⊗ Mandarin oranges (also satsumas, clementines, tangerines and blood oranges)
- ⊗ Pomegranate and drinks with pomegranate juice
- ⊗ Starfruit
- ⊗ Black licorice
- ⊗ Herbal dietary supplements

Carbohydrates

Some of the medications you are taking may raise blood glucose (blood sugar) levels. Carbohydrates in foods can also raise your blood glucose levels. If you have too much sugar in your blood for a long time, it can cause serious health problems such as heart disease, stroke, kidney disease, eye problems and nerve problems.

You can keep your blood glucose levels in control by learning which foods to include in your meals and snacks. Also, limit simple sugars in your diet such as regular soda, juice, sugar, syrups and candy.

Foods with Carbohydrates	Foods with Little or No Carbohydrates
Fruit and fruit juice	Most vegetables
Starches – grains, dried beans and starchy vegetables like potatoes, corn and peas	Meat, poultry, fish, eggs, cheese and meat substitutes
Milk and yogurt	Fats (oil and butter)
Desserts, candy, and other sweets	

How much carbohydrates you should eat

Your needs may be different, but most men need 4 to 5 servings (60-75 grams) of carbohydrate foods per meal. Most women need 3 to 4 servings (45-60 grams) per meal. Each serving should have about 15 grams of carbohydrates.

Food with 1 serving of carbohydrates

Fruits:

- 1 small piece of fresh fruit
- 4 oz of juice
- ½ banana
- 2 tablespoons of dried fruit

Starches:

- 1 slice of bread
- 1/3 cup cooked rice or pasta
- ½ cup corn, peas or beans
- ½ cup hot cereal or ¾ cup dry cereal
- 1 small tortilla

Dairy:

- 1 cup of milk
- ¾ cup yogurt

Tips for good blood glucose control

- Try to eat well-balanced meals that include carbohydrates, protein and fat at the same times every day.
- You can keep your blood sugars in balance by eating the same number of servings of carbohydrate foods at the same times each day.
- Try to eat something every 4 to 5 hours to keep you from getting too hungry between meals. If you will not be able to eat a meal for 5+ hours, eat a small snack in between.
- Fill half of your plate with non-starchy vegetables. These do not contain a lot of carbohydrates, but provide good nutrients.
- If you eat starchy foods, choose high fiber choices such as whole grain bread, brown rice and whole-wheat pasta.
- Limit or avoid foods high in added sugars. Read food labels to find out how many grams of added sugars are in the foods you commonly eat.

Minerals in food

We may ask you to change the amount of some minerals you eat. This will depend on how well your kidneys are working and how you are responding to your medications.

Potassium

Potassium is a mineral that keeps muscles and nerves working well. Too much or too little potassium can affect your heart.

Some medications, such as tacrolimus and cyclosporine, can make you have too much potassium in your blood. If this happens, or if your new kidney does not work well right away, you may need to eat less high-potassium foods.

Once your kidney works better or your blood levels of potassium are stable, you should go back to eating potassium-rich foods. High potassium foods have many health benefits.

Foods High in Potassium

- Artichokes
- Avocado
- Banana
- Beet greens
- Broccoli, cooked
- Dairy (milk, yogurt)
- Dried fruit
- Melons (cantaloupe, honeydew)
- Mango
- Kale
- Kiwi
- Nectarines
- Nutritional supplements (i.e. Ensure, Premier Protein shakes)
- Oranges and orange juice
- Parsnips
- Potatoes (baked, french fries, chips)
- Prunes and prune juice
- Some sports drinks
- Soy beans, cooked
- Spinach, cooked
- Tomatoes (including tomato sauce and juice)
- White beans, cooked
- Winter squash (acorn, butternut, pumpkin)

Magnesium

Magnesium helps your muscles work, reduces cramping, helps control blood pressure and blood glucose, and keeps your heartbeat steady.

Tacrolimus and cyclosporine may cause you to need more magnesium. You may need to take magnesium supplements to keep

your levels in a healthy range. But, if you eat magnesium-rich foods each day, you may not need supplements. Foods high in magnesium often also have protein, fiber and phosphorus.

Tips for eating more magnesium

- Eat many kinds of high-magnesium foods every day.
- Crush magnesium tablets if they are hard to swallow. Crushing may help your body absorb it better.
- Some people may get diarrhea when they take high-dose magnesium supplements. Tell your provider if that happens to you. A different form of magnesium may work better for you.

Foods high in magnesium

- Fish and other seafood
- Beans, peas and lentils
- Chocolate and chocolate milk
- Coffee
- Fruits such as bananas, avocado
- Nuts and seeds, nut butters
- Quinoa
- Soybeans and soymilk
- Vegetables such as potatoes, corn and spinach
- Whole grains such as brown rice and wild rice
- Whole grain cereals and bran

Phosphorus

Phosphorus helps our kidneys and muscles to work. It also sends signals to our nerves and helps our bodies use the nutrients we eat.

Phosphorus levels can be high after a kidney transplant, but this is usually a short-term problem. It does not mean you need to lower the amount of phosphorus you eat. Your dietitian will tell you if you need to follow a low phosphorus diet after your transplant.

If your phosphorus levels become low after your transplant, you need to eat high-phosphorus foods each day. You may need to take phosphorus supplements if you do not get enough from your food.

Foods high in phosphorus

- Dairy products like milk, yogurt and cheese
- Bran cereals
- Dried beans and peas
- Lentils
- Fish
- Beef, pork, turkey, chicken
- Nuts and nut butters
- Pancakes made from a mix
- Some cola drinks (diet preferred if you have elevated blood sugar levels)

Calcium

You are more likely to get bone disease because you have had kidney disease. Plus, you could get bone and joint weakness as a possible long-term side effect of taking prednisone. To help avoid osteopenia (low bone mass) and osteoporosis (brittle bones), you must eat enough calcium, phosphorus and vitamin D each day.

Eat a few servings of calcium-rich food in your diet each day. We may ask that you take calcium and vitamin D supplements when you leave the hospital.

Foods high in calcium

- Milk
- Yogurt
- Cheese
- Salmon
- Orange juice with added calcium
- Soy or almond milk with added calcium
- Tofu, firm (processed with calcium)

- Sesame seeds
- Bok choy, boiled or steamed
- Broccoli, boiled or steamed
- Collard greens, boiled or steamed
- Kale, boiled or steamed

Food safety

You are more likely to get infections now that you are on medicine that suppresses your immune system. You'll need to practice food safety to help stop foodborne illnesses.

Food that looks fine can still contain pathogens (disease-causing bacteria, viruses or parasites) that can make you sick. You should never taste a food to determine if it is safe to eat.

How to prepare food safely

Keep your hands and kitchen surfaces clean and free of bacteria by following these rules:

- Cleaning
 - Wash hands in warm soapy water for at least 20 seconds before and after handling food, or when you change tasks (like loading the dishwasher and then chopping vegetables).
 - If you can't always wash your hands, pack hand sanitizer or moist towelettes to use before eating.
 - Clean the tops of canned goods before opening them. Clean the can opener after each use.
- Fruits and vegetables
 - Wash fresh fruits and vegetables under running tap water, even those with skins and rinds that you will not eat.

- Wash fresh foods that are hard to clean (like leafy greens, berries and mushrooms) by soaking them in water first, then rinsing with fresh water.
- Lunch meat
 - **Do NOT** eat deli or lunch meat directly from the package. Instead, you can:
 - Heat lunch meat in the microwave or stovetop until it is steaming hot, before putting it into a cold sandwich.
 - Make a hot grilled sandwich.
- Raw meat and food
 - Use different dishes, utensils and cutting boards for raw and cooked foods.
 - Wash cutting boards, dishes, utensils and counter tops with hot soapy water after they have had raw meat, poultry or seafood on them.
 - Keep raw meat, poultry, seafood and eggs away from other foods in your grocery cart and in your refrigerator.
 - ⊗ **NEVER** place cooked food on an unwashed dish that held raw meat, poultry, seafood or raw eggs.
 - ⊗ **Do NOT** reuse marinades that you used on raw foods, unless you boil them first.
 - Marinate food in the refrigerator, not at room temperature.
 - Defrost food in the refrigerator, not at room temperature. If you defrost in the microwave, cook the food right after defrosting it.
 - ⊗ **Do NOT** taste raw or partly-cooked meat, poultry, eggs, fish or shellfish. This includes cake batter and cookie dough that contains raw eggs.
 - ⊗ **Do NOT** eat undercooked eggs, meat, poultry or fish.

Use a meat thermometer and cook to these internal temperatures:

Food	Internal Temperature or State
Beef, pork veal, lamb (chops, roasts or steaks)	145° (let meat rest for 3 minutes after you remove it from heat)
Ground meat	160°
Fully cooked ham	145°
Poultry (ground, whole, parts, stuffing)	165°
Eggs	Cook until yolk and egg white is firm
Egg dishes	160°
Fin fish	145° or meat is opaque (not clear) and flakes easily with a fork
Shrimp, crab or lobster	Cook until the meat is pearly and opaque (not clear)
Clams, oysters or mussels	Cook until the shells open during cooking
Scallops	the meat is milky-white or opaque (not clear) and firm
Leftovers, casseroles	165°

If you do not have a meat thermometer:

- Cook steaks to medium-well done.
- Cook ground meat, fish and poultry to well done.

Refrigeration and storage tips

- Refrigerate or freeze meat, poultry, seafood, eggs and other dishes within 2 hours after cooking (within 1 hour if the room temperature is above 90°F).
- Break up large amounts of food into small containers for quicker cooling in the refrigerator.
- Keep your refrigerator temperature just below 40°F (4.5°C).
- Keep your freezer temperature at 0°F (-18°C).
- ⊗ **Do NOT** eat food that has been left out at room temperature for 2+ hours. This includes meat, potato salad and other dishes at barbecues or picnics.
- Keep leftover seafood, stuffing and foods in broth or gravy in the refrigerator no longer than 1-2 days.
- Keep other leftovers in the refrigerator for no longer than 3-5 days (when in doubt, throw it out).

Eating at restaurants

- ⊗ **Do NOT** eat from salad bars or at buffet-style restaurants.
- ⊗ **Do NOT** eat sushi (raw or cooked), poke or ceviche.
- Choose hot sandwiches. All sliced meats must be heated.
- Ask the server how foods are prepared.
- ⊗ **Do NOT** eat food made from raw eggs, such as mayonnaise, salad dressings or sauces. These foods should be identified on the menu.

Cleaning

- Wash kitchen surfaces and tools with hot soapy water or a bleach solution after each use.
- To make your own bleach solution, mix 1 teaspoon of bleach with 1 quart of water.
- Use paper towels to clean up kitchen surfaces. If you use cloth towels, change them often and wash them in hot water in the washing machine.
- Throw away any cracked dishes, utensils, pots, pans and cutting boards. They are hard to clean well.


Raw and unpasteurized foods

- ⊗ **Do NOT** eat raw seed and vegetable sprouts.
 - Eat only pasteurized dairy products.
 - Wash all bagged “pre-washed” vegetables.
- ⊗ **Do NOT** eat unroasted nuts and nuts in the shell.
 - Drink only pasteurized juices and ciders.
- ⊗ **Do NOT** drink kombucha.

How to eat 2+ months after surgery

About 2 months after your transplant surgery, you will need fewer calories and less protein. This is a good time to work on long-term nutritional goals.

A healthy long-term diet after transplant



Use the plate method of eating, where $\frac{1}{2}$ your plate is non-starchy vegetables, $\frac{1}{4}$ is protein and $\frac{1}{4}$ is starchy food like whole grain bread, pasta, rice, cereal or starchy vegetables. You should also eat low-fat dairy products and fruit.

Heart health

Patients who have had kidney and liver disease are at increased risk of heart disease. Some of your new medications may raise your cholesterol and triglyceride levels. This can also increase your risk. Follow a heart healthy lifestyle to lower your risk.

Exercise

Start exercising as soon as your doctor says it is okay. Try to exercise for 30-60 minutes, 5+ days a week. Do heart-healthy exercise like:

- Walking
- Biking
- Aerobics
- Gardening
- Housework
- Strength training
- Swimming
- Running
- Yoga

Heart healthy eating

- Choose lean sources of protein, such as fish, chicken, turkey, beans, nuts or tofu.
- Add more heart-healthy fats to your diet. Unsaturated fats and omega-3 fatty acids may lower cholesterol and triglycerides:
 - Heart-healthy fats are in olive oil, canola oil, avocado, ground flaxseed, chia seeds, hempseeds and walnuts.
 - Eat 2 or 3 servings of fish every week. Salmon, tuna and mackerel are good sources of healthy fats.
- Lower the amount of saturated fat you eat:
 - ⊗ Do NOT eat red meat more than 2 times a week. When you do eat red meat, choose lean cuts such as loins or rounds.
 - ⊗ Try not to eat fried foods and butter.
 - Choose low-fat or non-fat dairy products.

- Eat no more than 2000 mg of sodium (salt) a day by following these tips:
 - ⊙ Do NOT add salt when cooking or at the table.
 - Season foods with spices, herbs, lemon juice and vinegars.
 - Cook at home. Use fresh meats and produce.
 - Try not to eat fast food or processed food.
 - Choose products with labels that say “Without Added Salt”, “No Salt Added” or “Unsalted”.
 - Read the “Nutrition Facts” label to find the amount of sodium in packaged foods.
 - For meals, avoid food with more than 600 mg of sodium per serving.
 - For a side dish or condiment, avoid food with more than 140 mg sodium per serving.
 - Eat 25 to 35 grams of fiber a day. Fiber helps you absorb less cholesterol and fat. It may also help you feel full so you don’t gain weight. Good sources of fiber include whole grains, beans, fruits and vegetables.

Heart health diets

If you want more information about heart healthy eating, we recommend two well-researched diets: The Mediterranean Diet and the DASH Diet (DASH = Dietary Approaches to Stop Hypertension).

Mediterranean Diet: Suggested books and resources

- The New Mediterranean Diet Cookbook: A Delicious Alternative for Lifelong Health by Nancy Harmon Jenkins and Marian Nestle. New York: A Bantam Book, 2009
- The Mediterranean Diet Plan: Heart Healthy Recipes and Meal Plans for Every Eater by Susan Zogneib and Phillip Anderson III. Berkley, CA: Rockridge Press, 2016

- Good Food and Great Medicine: A Mediterranean Diet and Lifestyle Guide by Miles Hassell and Mea Hassell. Hillsboro, OR: Lithtex, 4th edition 2018
- Harvard School of Public Health Diet Review: Mediterranean Diet: www.hsph.harvard.edu/nutritionsource/healthy-weight/diet-reviews/mediterranean-diet/

DASH Diet: Suggested books and resources

- The DASH Diet Action Plan: Based on the National Institutes of Health Research: Dietary Approaches to Stop Hypertension by Marla Heller. Deerfield, IL: Amadon Press, 2007
- Your Guide to Lowering Your Blood Pressure with DASH www.nhlbi.nih.gov/files/docs/public/heart/new_dash.pdf

Weight control

It is best to be at a healthy weight after your transplant and to stay that way. Being at a healthy weight will:

- Reduce your risk of diabetes, or improve your blood glucose control if you have diabetes.
- Reduce your risk of heart disease.
- Improve your blood pressure.
- Ease joint and back pain.
- Improve your energy level.

Watch for weight gain

After your transplant, you will be feeling better and your appetite may increase. You may also have fewer limits on your diet than you did before transplant. Medicines such as prednisone may increase your appetite, which can cause you to overeat and gain weight.

Tips for Keeping a Healthy Weight

- Eat from a smaller plate or bowl. This can make your mind feel like you are eating more than you are.

- Include regular exercise as part of your lifestyle after your transplant. Exercise helps build muscle and burn calories.
- Eat 3 smaller meals and 1 snack each day.
- Wait 20 minutes before taking a second helping. It takes that long for your body to feel full after eating.
- Chew on gum instead of eating between meals.
- Limit sweets and sugars.
- Limit sweetened beverages. They have calories but don't make you feel less hungry.
- Drink plenty of water.

If you find yourself gaining weight, try some of these internet resources:

- Aim for a Healthy Weight from the National Heart, Lung and Blood Institute
www.nhlbi.nih.gov/health/educational/lose_wt/index.htm
- USDA Nutrition
www.nutrition.gov/healthy-weight
- Academy of Nutrition and Dietetics
www.eatright.org/health#weight-loss

Caring for your health

Activity

Limits

After your transplant, we want you to get back to an active life.

There are only a few limits during the first 3 months:

- **Lifting**
 - For the first 6 weeks, do NOT lift 10+ pounds.
 - For the next 6-12 weeks after transplant, do NOT lift 20+ pounds.
 - 3+ months after transplant, there are no lifting limits.
- **Running**
 - For the first 3 months, do NOT jog or run on hard surfaces, such as cement or asphalt.
- **Bouncing**
 - For the first 3 months, do NOT do things that cause you to “bounce,” such as horseback riding, snowmobiling, and trail or cross country motorcycling.
- **Soaking**
 - Until your surgical cut is healed (6-7 weeks), NO tub baths or swimming.

Building muscle

Your muscles may feel weak from not using them and as a side effect of prednisone. You can make your leg muscles stronger by walking and bike riding. After 3 months, you can do sit-ups and other abdominal exercises to tone of your stomach muscles.

Returning to work

We encourage you to return to work as soon as you can. Most people can go back to work in 4 - 6 weeks, unless they do manual labor. You may be eligible for up to 12 weeks of job protected leave through OFLA or FMLA — check with your employer.

Play

Use common sense as your guide to any activity. As you get stronger, you will increase your amount of physical activity. If you are unsure if you should do an activity, please call the transplant office.

Sexual Activity

Wait 6 weeks before having sex to allow your surgical cut and muscles to heal. Your new organs are well protected. You will not harm them by having sex. Always practice safe sex since you are less able to fight sexually transmitted diseases (STDs).

Men's Fertility

Some men are able to father children, desire sex and function again — even if they couldn't during liver/kidney failure. Your sexual function may change depending on how your liver and kidney are doing and due to certain medications. If you notice a change in your sexual ability, ask your provider for advice. Talk to your hepatologist before you decide to have a child. Men should NOT father children while on certain immunosuppressive medications.

Women's fertility

Many women are fertile after transplant. Be sure to use a reliable form of birth control right away. You should start getting your periods again within several months of transplant. You may still be ovulating even if you do not have a regular period. Many women get pregnant and deliver healthy children after a transplant. If you are on Cellcept (mycophenolate) or Myfortic (mycophenolate delayed release), talk to your transplant team if you are pregnant or want to be pregnant. These drugs increase the risk of birth defects.

If you want to get pregnant, it is best to:

- Wait until at least 1 year after transplant.
- Not have other medical problems that might add to the risk of the pregnancy.
- Get lab tests more often.
- Adjust your immunosuppression drugs.
- Get proper prenatal care.

You will most likely have a successful pregnancy. But, there are some special risks to mother and child:


- **3 out of 10 mothers** will get high blood pressure and protein in their urine. In 10% of these cases, their kidney function will decrease.




- **3 out of 10 babies** will be born premature.



- **5 out of 100 babies** will have a major birth defect if the mother has had a transplant.



- **4 out of 100 babies** will have a major birth defect if the father has had a transplant.



Make sure to get your yearly well-woman check-up, with a Pap smear and a breast exam.

Long-term health

Transplant patients should follow these guidelines to care for the long-term health of your whole body.

Heart attack and stroke

Prevent heart attack and stroke by doing the following:

- Exercise daily
- Eat a low-fat, balanced diet with plenty of fruits and vegetables
- Limit weight gain
- Control blood pressure
- Watch blood sugars — people with diabetes should get help to control blood sugar
- Check cholesterol
- Do NOT smoke or use tobacco
- See your doctor every year

Diabetes

Post-transplant diabetes is a known risk for transplant patients.

Watch your blood sugar and start treating it with diet and exercise.

Many patients need insulin after transplant.

Travel

Contact a travel clinic before you go to another country. They will tell you which vaccines you need to travel there and what your risks of infection are. Please check with your transplant team with any questions. Remember, after transplant you should **NEVER get LIVE virus vaccines.**

Bone disease

All transplant patients get some bone disease. You should have a bone density study at least once every 2 years. Ask your primary care provider how to prevent bone disease with diet and exercise.

Regular cancer screenings

Contact the transplant office if you are ever told you have cancer. Due to the immunosuppression drugs you are on, you should have the following tests:

Women

- All women — pelvic exam and pap smear every year
- Age 40+ — yearly mammogram
- Age 30+ who has a female relative diagnosed with breast cancer before menopause — yearly mammogram

Men

- Age 45 if your father or brother has had prostate cancer — your first prostate-specific antigen test (PSA)
- Age 45 if you are African American — your first PSA test
- Age 50+ — yearly PSA test

Everyone ages 50+

- Colon cancer screening (with rectal exam), every 2 years
- Fecal occult blood test, every year
- Colonoscopy, every 5 years

Skin cancer

You have a **much higher risk of skin cancer** due to the medicine you take to help prevent rejection. The most common type of skin cancer we see after transplant is squamous cell carcinoma — can be up to **65 times more common** than in non-transplant patients! Patients usually get their first skin cancer 3-8 years after their transplant. You are at even higher risk if you are light-skinned, had much sun-exposure in your lifetime, are older, or have already had skin cancer. Squamous cell carcinoma may spread (metastasize) to your lymph nodes in 5-10% of cases.

How to lower your risk

- Avoid the sun from 11 a.m. to 3 p.m. — when the harmful rays are most intense.
- Wear sunscreen with both UVA/UVB protection of SPF 30+ every day.
- Use plenty of sunscreen on any exposed skin and reapply every 2 hours.
- Don't forget to protect your lips and ears.
- Wear a wide-brimmed hat and protective clothing, such as long sleeves, pants and sunglasses.
- See a dermatologist for skin exams after your transplant.



Call your dermatologist or primary care doctor right away if you:

- See any change in your moles, or
- Have a new sore that does not heal

Preventing infectious disease

1. **Wash your hands often.** It is the one best way to prevent infection!
2. **Wear masks and gloves** at these times:
 - First 3 months after transplant, when you are:
 - In the hospital
 - Around active construction
 - Near farming areas
 - In crowds
 - **Do NOT garden during the first 3 months after transplant**
 - From 3 months to 1 year after transplant, where a **mask and gloves** when you are gardening.
 - After 1 year, **always wear gloves** when you garden.
3. **Get the right vaccines.**
 - Get your flu shot every year.
 - If you get hurt but haven't had a tetanus booster within the last 5 years, call your primary care provider.
 - Transplant patients should **NEVER get LIVE vaccines**. Live vaccines have live organisms and you could get the actual disease since your immune system is less resistant to disease. Vaccines made from dead organisms are safe.
 - ⊗ **Do NOT get these vaccines:**
 - ⊗ Smallpox ⊗ Measles
 - ⊗ Mumps ⊗ Rubella
 - ⊗ Oral polio ⊗ Chicken pox
 - ✓ **You CAN get these vaccines:**
 - ✓ Injectable polio
 - ✓ Flu shot (Influenza A and B) - every year
 - ✓ Pneumovax - every 5 years
 - ✓ TB skin test
 - ✓ Diphtheria/Tetanus - every 10 years
 - ✓ Shingrix (a specific Shingles vaccine)
 - ✓ COVID

Avoiding infection

You need to always protect yourself from infection. Here's how to do that:

- **Wash your hands often** with soap and water.
 - This is the #1 way to protect yourself.
 - Wash your hands before and after changing the dressing (bandages) of any wounds.
 - Try to keep your hands away from your face and mouth.
- **Avoid sick people.**
 - Stay away from people who are ill.
 - Ask friends to visit only when they are well.
- **Stay away from crowds** — especially during cold and flu season.
 - In the first few months and when you are highly immunocompromised — NO crowded places like stores, movies, restaurants and churches.
 - After that, wear a surgical mask in large crowds until your prednisone dose is down to 10 mg a day.
- **Don't share items that touch spit** (saliva) or mucus.
 - ⊗ Do NOT share forks, spoons, cups, toothbrushes, etc.
 - ⊗ Do NOT share razors.
- **Do NOT work in the garden or yard** for the first 3 months after transplant.
 - After 3 months, wear gloves when touching soil.
 - Avoid compost piles, wet leaves and rotting organic matter. They can carry molds that cause infections.
- **Wear a surgical mask** if you are **near a construction site** until your prednisone dose is down to 10 mg a day.
- **Do NOT touch animal waste.**
 - Avoid contact with any animals that roam outside.
 - Do NOT clean birdcages, fish tanks, turtle tanks or cat litter boxes.

- When someone else changes the cat litter, they should first cover the litter box and take it out of your home before changing it.
- Stay away from reptiles, turtles, amphibians (frogs), hamsters, and guinea pigs. They can carry infections.
- **Be safe with vaccines.**
 - NO close contact with anyone who had the oral (by mouth) polio vaccine within 2 weeks. The virus may shed from them for up to 2 weeks after their vaccine.
 - Do NOT change the diapers of kids for the first 2 weeks after they get live immunizations including the chicken pox vaccine.
 - If you have an infant, make sure they get the Salk polio injection.
 - You CAN be near a child who recently got a MMR shot.

Dental care

Make sure to get a dental checkup every 6 months. Infections could be serious because of your lowered immunity. Be sure to:

- Wait until 3 months after transplant to get routine dental care.
- Let your dentist know you had a liver and kidney transplant and are on immunosuppressive medications.
- You do NOT need to take antibiotics before routine dental visits unless you have:
 - Had a heart transplant
 - Artificial heart valves
 - Any artificial material in your heart
 - Had a previous infection in your heart
 - Congenital heart defects.
- If your dentist is unsure about antibiotics before dental procedures, we follow the guidelines set by the American Heart Association and American Dental Association.
- Ask your dentist to call our transplant office if they have any questions.

Drugs, tobacco and alcohol

Tobacco

Do NOT use tobacco.

- Chewing tobacco can lead to neck and mouth cancers.
- Smoking raises your risk of:
 - Bad surgery results
 - Cancer
 - Atherosclerosis (fatty deposits in blood vessels)
 - Heart attack
 - Stroke

- Please contact your transplant nurse if you need help quitting tobacco.

Marijuana

We do NOT recommend that you use marijuana. It can cause lung and brain fungal infections.

Alcohol

Do NOT drink alcohol. Do NOT drink “nonalcoholic” drinks like “near beers” and wine coolers. Any type of alcohol can harm your liver and destroy its cells.

Pets at home

For the most part, transplant patients can still live with pets in their home. Wash your hands after handling or cleaning up after your pets, and follow the guidelines below.

Dogs and cats

- Keep vaccinations up to date.
- Treat regularly for flea and tick prevention.
- Transplant patients CANNOT clean out a cat's litter box.

Birds

- Transplant patients CANNOT clean out a bird cage or chicken coop.

Reptiles

- ⊗ Transplant patients should NOT touch reptiles. They may carry salmonella and should NOT be pets.

This Page Intentionally Left Blank

Complications

Rejection

Your body may treat your new liver and kidney as foreign objects and start attacking them. This is called “rejection” and it can happen at any time. Rejection does not mean loss of your organs. Often, we can treat it with medication. Some patients need more intense treatment. We watch for rejection by looking at your labs and biopsies of the organs.

- 10% to 30% of all transplant patients will have at least 1 rejection episode. The first one often happens within 3 months of surgery.
- We will watch your liver function tests (LFTs) to see what is happening to your liver.
- Kidney function is watched through labs.

Treatment

If your body starts to reject your liver or kidney, we can control that by changing the amount of your immunosuppressive medications or by adding a new one. If you have a severe rejection, we may need to admit you to the hospital for stronger IV medicine.

Risk

The risk of rejection goes down over time. But, it can still happen at any time. Lower your risk by taking good care of yourself, taking your medications as prescribed, and having your blood tests done.

Recurring Hepatitis C

A liver transplant does not cure Hepatitis C since the virus is found in the blood. If you had Hepatitis C before your transplant and did not get treatment or clear the virus, you will have it again (recur) at some point after your transplant. Most people with recurring Hepatitis C will get cirrhosis within 5 years of their transplant, if the virus is not treated. There are many new medications to treat Hepatitis C. Talk to your hepatologist about when you should start treatment.

Recurring kidney disease

Some kidney diseases that cause organ failure can happen again (recur) in your new kidney. If you have a disease that can recur, we would have already talked to you about this during your transplant work up. We watch for recurring disease by looking at your labs and doing a kidney biopsy, if needed.

Infection and viruses

You are at more risk of getting infections due to the anti-rejection medications you take. These infections can be serious, so let us know right away if you have any of these symptoms:

- Fever
- Chills
- New diarrhea
- New or unexplained pain
- Cough
- Tiredness for no reason
- Not feeling well
- Urine problems such as going too much, pain when urinating or cloudy urine

Also, let us know if you have a viral infection like Cytomegalovirus (CMV), Epstein Barr virus (EBV), or BK virus.

CMV

Cytomegalovirus (CMV) is a very common viral infection that usually has no symptoms in non-transplant patients. Transplant patients need antiviral medications to help prevent CMV infection. Symptoms of infection can be fever, extreme tiredness (fatigue), diarrhea, stomach (abdominal) pain, cough and low white blood cell count. We can do a lab test called a CMV PCR to see if you have the infection. If you are infected, your transplant team will let you know what treatment you need.

Diabetes

Your anti-rejection medications increase your risk of developing diabetes after transplant. If you had diabetes before transplant, you may need to take more diabetic medications. You may need to see a diabetic specialist.

Cancer

Your anti-rejection medications increase your risks of certain types of cancer. The most common cancers are skin cancers, cancers of the genitals and urinary system, and lymphoma (cancer of the white blood cells).

All patients should see a dermatologist once a year. Learn about your cancer risks, and tell your doctor right away about any new symptoms or concerns.

Other new health concerns

You may develop unrelated health problems after transplant. Work closely with your doctor on your overall wellness.

Anxiety and depression

You and your family face a new lifestyle after transplant. This may cause you to feel nervous, stressed or depressed. It is normal to feel anxiety and depression after transplant. Ask your social worker about counseling services that can help you and your family deal with these changes. They can help you address things like:

- Mood swings.
- Job planning.
- Rehabilitation.
- Family stresses — such as parent-child conflicts and marital conflict.
- Financial concerns — such as questions about Medicare, disability or insurance.

High cholesterol

Many immunosuppressant drugs can lead to high cholesterol. This can cause blood vessels — including the ones attached to the transplanted organ — to become clogged. This may slow blood flow, affect the success of your transplant, and may lead to heart disease. Talk to your doctor about how to reduce the risk factors of heart disease, including controlling your cholesterol.

High blood pressure

High blood pressure (hypertension) is common. If you had high blood pressure before transplant, you may still need to be treated for it. Even if you never had it before, high blood pressure can be a side effect of anti-rejections medications. You may also get high blood pressure if your kidneys are not working well. Your doctor may prescribe one or more medications to treat high blood pressure.

Diabetes

As mentioned before, you have an increased risk of diabetes after transplant. Diabetes is an increased level of sugar in your blood. Some of the immunosuppressive medications that you take may cause diabetes. If you develop diabetes, we will give you special teaching and ongoing follow up about how to deal with it.

- Symptoms of diabetes may include:
 - Needing to pee more often.
 - Increased thirst.
 - Blurred vision.
 - Confusion.
 - Extreme hunger.
 - Constant itching.
 - Irritability.
- How to lower your risk:
 - Lose weight if you are overweight.
 - Eat a balanced diet — talk to your dietitian, transplant coordinator and doctor about the best diet for you.
 - Exercise.
 - Reduce stress — talk to your social worker, transplant coordinator or doctor about any increased stress you are having.
- Uncontrolled diabetes can lead to:
 - Heart disease.
 - Stroke.
 - High blood pressure.
 - Blindness.
 - Kidney disease and failure.
 - Amputation (loss of a limb).

This Page Intentionally Left Blank

Writing to your donor's family

Some transplant patients choose to write to their donor family. This is strictly your choice. It is not expected and there are no requirements. But, if you would like to write the family, here are some tips.

What to write

We know the words can be hard to find, so here are some ideas to help you get started:

- Say “thank you.” It may not completely get across how much the gift of transplant means to you, but it is always welcome.
- Share a bit about yourself (job, hobbies, etc.), your family and how the transplant has changed your life (what you can now do or enjoy because of the transplant).
- Include photos of yourself or family.
- Avoid religion since you do not know the donor family. Instead, you can use general statements of “feeling blessed.”
- Do NOT pressure the donor family to meet in person or to write back to you.

What to keep private

Do NOT include:

- ⊗ Last name
- ⊗ City where you live, address or phone number
- ⊗ Hospital where you had the transplant
- ⊗ Names of your medical team

The process

After you write your donor family:

1. Place your letter or card in an unsealed envelope.
2. On a separate paper, write your full name, transplant center and date of transplant. This will help get the letter to your donor family.
3. Put the paper and unsealed letter together in another envelope and mail to:

OHSU Transplant Social Work
Mail Code: SJH 569
3181 S.W. Sam Jackson Park Rd.
Portland OR 97239

What happens next

We will call you to let you know we got the letter. We will send it to Cascade Life Alliance (CLA). CLA will forward the letter to your donor family if the family allows it. It could take many weeks for your letter to get to the family. We will let you know if your donor family decides not to accept letters or if the CLA cannot reach them.

How will I know if the donor family writes back?

If the donor family writes you back, your social worker will let you know and ask you if you would like to accept the letter.

What if I need help deciding if I should write or what to write?

You can always talk with your social worker. They will not decide for you, but they can help you process your feelings. This can help you make the right choice for you.

References

How to prepare for your kidney transplant biopsy

A kidney transplant biopsy is when we take a very small piece of tissue from the transplant kidney and look at it under a microscope for signs of rejection or renal disease. We get the tissue through a small hole in your skin (minimally invasive) and use ultrasound to guide a biopsy needle into your kidney.

How to prepare

- We will let you know the date and time of your biopsy.
- We will collect your blood and urine the day of the biopsy.
- ⊖ Do **NOT** eat or drink for the **2 hours before** your biopsy.
- You will need a **designated driver**. You and your driver should expect it to take 5-8 hours from the time you get to the hospital to when you can leave.

Your medications

- Take your morning blood pressure medication. If your blood pressure is too high the day of the biopsy, we will need to reschedule it.
- Bring your other medications with you to the biopsy.
- **STOP** taking blood thinners (anticoagulants) before your biopsy. Blood thinners will increase your risk of bleeding.

Blood thinner	STOP taking	START taking again
Coumadin (warfarin)	5 days before your biopsy	
Aspirin	7 days before your biopsy	
Plavix (clopidogrel)	7 days before your biopsy	
Eliquis (apixaban)	48 hours before your biopsy	
Xarelto (rivaroxaban)	48 hours before your biopsy	
Lovenox (enoxaparin)	36 hours before your biopsy	

Where to go on the day of your biopsy

1. 30 minutes before your biopsy, go to the outpatient lab on the 3rd floor of **OHSU Physicians Pavilion** (3270 SW Pavilion Loop, Portland, OR 97239) for your lab draw.
2. After your lab draw, go across the street and check in at Admitting on the 9th floor of **OHSU Hospital** (3181 SW Sam Jackson Park Road, Portland, OR 97239).

What happens during your biopsy

1. We take you to the Procedural Care Unit (PCU).
2. We give you a gown to change into.
3. We explain the procedure and ask you to sign a consent form.
4. We numb the area of your body where we will insert the needle.
5. We use ultrasound to see where the needle tip is and use the needle to take a tissue sample.
6. The procedure should take 30 - 60 minutes.
7. After the biopsy, we take you to the recovery area in the PCU for you to rest for a few hours. We watch you for signs of bleeding and other complications during this time.
8. Once it is time for you to leave, your designated driver picks you up. You are NOT allowed to drive yourself home.

After the biopsy

- Do NOT do any strenuous physical activity for the next 48 hours.
- **Go to your local emergency room if you have any of these signs of bleeding:**
 - Your heart feels like it is beating too hard, too fast, skipping a beat, or fluttering (palpitations)
 - You feel light-headed
 - Your pain gets worse
- You will get the biopsy results within 1 week.



Diagnostic tests

Biopsy

We use this procedure to see if your body is rejecting your new organs. We numb your skin and take a very small piece of your kidney or liver using a needle.

If you are on any blood thinners such as aspirin, Plavix or Coumadin, your coordinator will discuss a plan with you before scheduling your biopsy.

MRCP (Magnetic Resonance Cholangiopancreatography)

We use this special non-invasive MRI (magnetic resonance imaging) to look at the drainage system or ducts of the gallbladder, pancreas and liver. This test helps us find blockages, pancreatic cysts, bile duct stones, narrowing in the ducts or tumors.

PTC (Percutaneous Transhepatic Cholangiography)

This procedure x-rays the bile ducts. We numb the area and insert a thin needle in your skin, through the liver and into a bile duct. Then, we inject a dye so we can see the bile duct system. We may need to insert a thin tube to drain bile into a bag outside the body or into the small intestine. We can also remove stones and insert balloons to open up narrowing ducts.

ERCP (Endoscopic Retrograde Cholangiopancreatography)

For this test, we use an endoscope to look at and treat the GI tract. An endoscope is a thin, flexible tube with a tiny video camera and light on the end. The GI tract includes the stomach, intestine, liver, pancreas and gallbladder.

Doppler Ultrasound

This test uses sound waves to see the blood flow in your abdomen. It can help us find blockages in the blood flow to your liver, as well as other problems in the organs.

CT (Computed Tomography) Scan

We use a CT scan to get a 3-dimensional x-ray of your internal organs. It can help us find problems in the liver, spleen, colon, pancreas, kidneys and more. Sometimes we use a dye to see better. We usually give the dye by IV but sometimes by mouth.

Over-the-counter medicines that are safe to take

Symptoms	Safe over-the-counter choices
Fever	Acetaminophen (Tylenol). Call your nurse coordinator or on-call provider if your fever is over 101° F.
Head and body aches	Acetaminophen (Tylenol) Take NO more than 3,000mg in 24 hours.
Sneezing, itching or runny nose	Loratadine (Claritin), Cetirizine (Zyrtec), Fexofenadine (Allegra)
Stuffy nose (nasal and sinus congestion)	Nose sprays: Sodium chloride (Ocean), (SinuCleanse), Oxymetazoline (Afrin), Phenylephrine (Neo-Synephrine), Pseudoephedrine (Sudafed). Do NOT use Sudafed if you have high blood pressure.
Chest congestion	Guaifenesin (Robitussin), Coricidin HBP Chest Congestion & Cough
Wet cough (lots of phlegm)	Guaifenesin (Robitussin)
Dry cough (no phlegm)	Dextromethorphan (Delsym), Guaifenesin/ Dextromethorphan (Robitussin DM), Coricidin HBP Chest Congestion & Cough
Sore throat	Lozenges - Benzocaine, Halls
Constipation	Polyethylene glycol (MiraLAX), Psyllium (Metamucil), Docusate (Colace), Docusate with senna (Senokot-S), Bisacodyl (Dulcolax), Senna
Diarrhea	Loperamide (Imodium AD). Call your nurse coordinator during the work day if you have more than 3 loose stools (poop) in one day.
Trouble sleeping (insomnia)	Diphenhydramine (Benadryl), Doxylamine (Unisom), Melatonin
Stomach upset	Calcium carbonate (TUMS)
Heartburn	Ranitidine (Zantac), Famotidine (Pepcid AC), Nizatidine (Axid), Omeprazole (Prilosec),
Gas	Simethicone (Gas-X)
Dry, irritated eyes	Artificial Tears, Ketotifen (Zaditor)
Nausea and vomiting	Meclizine (Antivert) Call your nurse coordinator or on-call nurse if you cannot keep down your anti-rejection medications.
Joint pain	Capsaicin cream (Capzasin-P), Lidocaine (Aspercreme, Icy Hot), Camphor & Menthol (Bengay)
Skin irritation, insect bites, poison ivy	Hydrocortisone (Cortisone-10)

Magnesium in certain foods

	Milligrams (mg)	% Daily Value
Halibut – cooked, 3 ounces	90	20
Almonds – dry roasted, 1 ounce	80	20
Cashews – dry roasted, 1 ounce	75	20
Soybeans (mature) – cooked, ½ cup	75	20
Spinach (frozen) – cooked, ½ cup	75	20
Nuts (mixed) – dry roasted, 1 ounce	65	15
Shredded wheat cereal – 2 rectangular biscuits	55	15
Oatmeal (instant, fortified) – cooked with water, 1 cup	55	15
Potato (with skin) – baked, 1 medium	50	15
Peanuts – dry roasted, 1 ounce	50	15
Peanut butter (smooth) – 2 tablespoons	50	15
Wheat bran (crude) – 2 tablespoons	45	10
Black-eyed peas – cooked, ½ cup	45	10
Yogurt (plain, nonfat) – 8 fluid ounces	45	10
Bran flakes – ½ cup	40	10
Vegetarian baked beans – ½ cup	40	10
Brown rice (long-grained) – cooked, ½ cup	40	10
Lentils (mature seeds) – cooked, ½ cup	35	8
Avocado (California) – ½ cup pureed	35	8
Kidney beans (canned) – ½ cup	35	8
Pinto beans – cooked, ½ cup	35	8
Wheat germ (crude) – 2 tablespoons	35	8

Phosphorus in certain foods

	Serving size	Phosphorus (mg)
Mac and cheese (from mix)	1 cup	400
Liver and organ meats	3.5 ounces	400
Yogurt (plain, nonfat)	8 ounces	385
Pancakes (from mix)	3 4-inch pancakes	368
Yogurt (regular, no added probiotics)	1 cup	326
Mac and cheese (from scratch)	1 cup	322
Pudding (made with low fat milk)	½ cup	313
Dried beans and peas	1 cup, after boiling	266
Salmon	3 ounces, cooked*	252
Halibut	3 ounces, cooked*	242
Milk (nonfat)	8 ounces	247
Pizza (cheese and pepperoni)	1 slice	234
Ice cream (low fat)	1 cup	200
Peanut butter	3 tablespoons	180
Lentils**	½ cup, cooked	178
Beef and turkey	3 ounces, cooked*	173
Cheese (low fat)	1 ounce	171
Cream soup (made with low fat milk)	1 cup	160
Chicken	3 ounces, cooked*	155
Biscuit (from mix)	1	140
Almonds**	1 ounce (23 nuts)	134
Mozzarella cheese (part skim)	1 ounce	131
Peanuts**	1 ounce	107
Egg	1 large, cooked	104
Bread (whole wheat)	1 slice	57
Bread (enriched white)	1 slice	25

* A 3-ounce serving is about the size of a deck of cards.

** Your body is only able to absorb 50% (half) of the phosphorus from nuts, seeds and grains.

Glossary

Acute

Short and severe

Acute Tubular Necrosis (ATN)

When the new kidney doesn't work temporarily due to its storage time before transplant

Albumin

A protein made by the liver that helps your body maintain its balance of fluids

Alkaline Phosphatase (Alk Phos)

An enzyme made by the liver. High levels can mean your liver or other organs aren't working well.

Analgesic

Pain medicine

Anemia

A low number of red blood cells

Anesthetic

Medication that reduces pain

Antacid

A drug that protects your digestive system and relieves heartburn

Antibody

A part of the immune system that fights infection or foreign organisms or tissue

Antigen

A harmful substance that enters your body and starts an immune response to make antibodies

Arteriosclerosis

A hardening of the arteries that blocks blood flow to the kidneys

B Cells

A type of white blood cell that makes antibodies

Bacteria

Germs that can cause disease or infection

Bile

A fluid made by the liver that is stored in the gallbladder and released into the small intestine to help absorb the fats you eat

Bile Duct

The tubes that bile flows through

Bile Leak

A hole in a bile duct that causes bile to spill into the abdominal cavity

Biliary Stenosis

When a bile duct gets narrow

Biliary Tree

All bile ducts in and out of the liver that carry bile to the intestines

Bilirubin

A fluid made by the breakdown of red blood cells in the liver. High levels can lead to jaundice.

Biopsy

Procedure to see if your body is rejecting your liver or kidney, where we numb your skin and take a very small piece of your liver or kidney using a needle.

Bladder

The organ that receives and stores urine from the kidneys until it is urinated out of the body

Blood Urea Nitrogen

A waste product that your kidneys remove from your blood

Cadaveric Donor

An organ donor who has just died

Calcium

A mineral needed for bone growth, blood clotting, and for the heart and nerves to work

Catheter

A small rubber tube (commonly inserted into the bladder to drain urine)

CellCept

An immunosuppressive drug used to limit or reverse rejection

Cholangiogram

A test that looks for leaks or blockages in the liver's bile ducts

Cholangitis

A bacterial infection in the bile ducts

Cholestasis

A buildup of bile in the liver — caused by medicines, injury to the liver, liver disease, total parenteral nutrition (TPN)

Cholesterol

A kind of fat that your body needs, but too much can cause heart disease

Chronic

Something that lasts over a long period of time

Chronic Renal Insufficiency

Permanent damage to both kidneys, treated by dialysis or transplant

Cirrhosis

A disease that causes permanent scarring of the liver

CMV (Cytomegalovirus)

A common viral infection that can be harmful to transplant recipients if they get it after transplant

Coagulation

The process of blood clotting

Coagulopathy

Abnormal blood clotting

Complete Blood Count (CBC)

Blood test that measures what makes up blood, including hemoglobin (Hgb), hematocrit (Hct), platelets (Plt), and the types of white blood cells (WBC)

Corticosteroids

A group of immunosuppressive medications that includes prednisone and prednisolone

Creatinine

A waste product made by your muscles as part of normal activity. Normal kidneys filter out creatinine. The higher the creatinine level in your blood, the lower your kidney function.

Crossmatch

A test to see how compatible a donor's blood is with that of someone who could receive it

CT Scan

A 3-dimensional x-ray of internal organs

Cyclosporine

A powerful immunosuppressive drug

Cyst

A sac-like structure that contains fluid and matter

Cytomegalovirus (CMV)

A common viral infection that can be harmful to transplant recipients if they get it after transplant

Diabetes

A disease that makes your blood sugar higher than it should be, causing health problems

Diabetic Nephropathy

Kidney failure as a result of diabetes

Dialysis

A procedure where toxins are cleaned out of your blood

Diastolic Blood Pressure

The bottom number of a blood pressure measurement. It is the pressure on the arteries between heartbeats.

Echocardiogram

A test that uses beams of ultrasonic waves to measure the motion and position of the heart and nearby tissue

Edema

Swelling of a certain area of the body (such as the hands or legs) due to too much fluid

EGD (esophagogastroduodenoscopy)

A test where we use an endoscope to look at the lining of the esophagus, stomach and first part of the small intestine

Electrocardiogram

A test where we place electrodes on your chest to measure the heart rhythm and look for injury to heart tissue

Electrolyte

A dissolved mineral (such as magnesium or potassium) that helps the body function

Endoscope

A small telescope-like instrument that we use to look into your body to see the esophagus, stomach and small intestine

Endotracheal Tube

A tube we insert through the mouth and into the windpipe to help you breathe during surgery

End Stage Renal Disease

When your kidneys no longer work well enough to keep you alive (working at less than 10%)

Enzyme

A bodily protein that can break down other substances

ERCP (endoscopic retrograde cholangiopancreatogram)

A test where we use an endoscope to look at the drainage system or ducts of the gallbladder, pancreas, and liver (the biliary tree).

Fibrosis

When healthy tissue in your liver becomes scarred and can't work as well, leading to cirrhosis

Gallbladder

An organ that stores bile, which helps break down fat from food. It is attached to the liver and we remove it during transplant.

Gastroenterologist

A doctor who treats diseases of the digestive system, including the liver

Gastrointestinal (GI)

The tract between the mouth and the rectum, including the intestines and stomach

Glomerular Filtration Rate (GFR)

A test to measure the level of kidney function

Glucose

Sugar found in the blood or urine

Graft

A transplanted tissue or organ, such as a kidney or pancreas

Helper T-cell

The white blood cell that tells the immune system to fight infection or foreign substances, such as transplanted tissue

Hematocrit

A blood test that counts the number of red-blood cells in your blood

Hematoma

A bruise or swelling caused by a build-up of blood in tissue

Hemoglobin

Protein in red blood cells that carry oxygen from the lungs to the rest of your body.

Hepatic

Relating to the liver

Hepatitis

Liver inflammation, usually caused by a virus

Hepatologist

Doctor who treats liver disease

Hepatomegaly

Enlarged liver

Herpes

A family of viruses that can cause lip, genital sores, or other symptoms

Hirsutism

Excessive hair growth. A common side effect of cyclosporine in both men and women.

Human Leukocyte Antigens (HLA)

Genetic markers you get from your parents

Human Leukocyte Antigen (HLA) Compatibility

A test done on the donor and the potential recipient to see how the recipient's cells would attack the graft

Hypertension

High blood pressure

Hypotension

Low blood pressure

Immune System

Your cells, tissues and organs that work together to fight foreign organisms or tissues in your body

Immunosuppression

Making it hard for your body to fight a foreign object like a transplant or disease. You can do this by taking certain drugs to help prevent or control a transplant rejection.

Immunosuppressive Agents

Medications you take to prevent your body from rejecting an organ transplant

Insulin

A hormone made by your pancreas that controls blood sugar levels

Intravenous (IV)

Putting fluids or drugs directly into your vein with a needle or catheter

Jaundice

When your skin and eyes look yellow because your liver doesn't work as well as normal

Kidney

Organ that filters the waste and extra fluid from your blood that becomes urine.

Kidney Failure – Acute

When your kidneys suddenly stop working well, but it can be reversed

Kidney Failure – Chronic

When your kidneys slowly stop working over time, but it can't be reversed

Leukocyte

A white blood cell that helps fight infection

Liver Enzymes

Proteins that speed up chemical reactions in the body

Liver Function Tests (LFTs)

Blood tests to see how well the liver is working — measures ALT, AST, GGT, bilirubin and alkaline phosphatase.

Lymphocyte

White blood cells made by the lymph glands and defend your body against infection and foreign substances.

MRCP (Magnetic Resonance Cholangio-Pancreatography)

A special kind of MRI to look at the drainage system or ducts of the gallbladder, pancreas, and liver (the biliary tree). It is used to find blockages, pancreatic cysts, bile duct stones and tumors.

Nephrectomy

A surgery to remove one or both kidneys

Nephrologist

A physician who specializes in diagnosing and treating kidney disease

Neutrophil

A type of white blood cell

Noncompliance

When you fail to follow a treatment plan — like taking medications, getting tests on time, taking vital signs, etc. This often shortens the lifetime of the transplanted organ(s).

Orally

By mouth

Panel Reactive Antibody (PRA)

A blood test to find how likely your body is to reject an organ transplant from the general public. A score of 0/42 or 0% sensitized is best, a score of 42/42 or 100% sensitized is worst. You will tend to get a worse score if you've been pregnant, had blood transfusions or had previous transplants.

Phlebotomy

Removal of around 1 pint of blood through a vein

Platelet

A small blood cell you need for clotting

PJP (Pneumocystis Jiroveci Pneumonia)

An infection of the lungs found in people with weakened immune systems

Pneumocystis Carinii Pneumonia (PCP)

A type of pneumonia that people with suppressed immune systems tend to get

Polycystic Kidney Disease (PKD)

A genetic disease that causes cysts to grow in place of normal kidney tissue

PTC (Percutaneous Transhepatic Cholangiography)

A procedure that x-rays the bile ducts

PTLD (Post-Transplant Lymphoproliferative Disease)

A group of life-threatening disorders that can happen after transplant. It is treated by lowering immunosuppression and giving antiviral medications.

Potassium

A mineral your body needs but too much can harm your heart. High levels often happen when your kidneys don't work well.

Prophylactic Medication

Drugs that help prevent disease

Red Blood Cells

The part of the blood that carries oxygen to body tissues

Rejection

When the immune system attacks what it thinks is a foreign substance (such as a transplanted kidney)

Renal

Related to the kidneys

Renal scan

Procedure where we inject a small and safe amount of radioactive dye into your vein to see a picture of the kidney and if there are any changes in blood flow

Sepsis

A severe infection that has spread to the blood stream

Shingles

A herpes virus infection that causes a red, painful skin rash

Signs

Things you can measure to see how well your body is working — such as heart rate, blood pressure, breathing rate and temperature

Sleepy Kidney

An expression for the temporary delay in kidney function after a transplant (also known as ATN)

Sodium

The main salt that is found in blood

Stenosis

Narrowing of a passage in the body (also known as “stricture”)

Systolic Blood Pressure

The top number of a blood pressure measurement. It is the pressure on the arteries when the heart beats.

Symptoms

Things you feel, such as pain, dizziness or fatigue

T Cells

White blood cells that play a major part in rejection

T-Tube (Turcotte Tube)

A tube we place in the bile duct so that bile can drain into a bag outside the body

Thrombosis

A blood clot

Thrush

A fungal infection in the mouth

Tissue Typing

A blood test to compare a person's major antigens with a donor's to see if they are a good match

Toxins

Waste products in the blood that are bad for the body when there is too much

Turcotte Tube

A tube placed in the bile duct that allows bile to drain into a bag outside the body.

Ultrasound

Procedure that uses sound waves to see the size of an organ and if there is fluid around it or urinary blockage

Ureter

One of 2 tubes in your body that carries urine from the kidney to the bladder

Urethra

The tube in your body that carries urine out of the bladder

Urinary Catheter

Tube that we insert into the bladder to drain urine

Urinary Tract

The body system that makes, carries, stores and gets rid of urine. It is made up of the kidneys, ureters, bladder and urethra.

United Network for Organ Sharing (UNOS)

Organization that manages the national transplant waiting list to make sure it is fair, and collects data on transplant recipients and organ donors

Virus

A small germ that causes infection

Wean

To slowly withdraw or reduce. We may wean immunosuppression slowly over time if your transplanted liver is working well.

White Blood Cells

The part of the blood that fights infection

This Page Intentionally Left Blank

Forms

Lab results

Lab name: _____

Medical record number: _____

Lab phone: 503-494-7383 MyChart help desk: 503-494-5252

Date							
Glucose							
BUN							
Creatinine *warning if increase of 0.2 or more							
Potassium							
Calcium							
Phosphorus							
Weekly magnesium							
WBC *warning if less than 3.0							
Hematocrit							
Drug Level							
Target Range							

Lab results

Lab name: _____

Medical record number: _____

Lab phone: 503-494-7383 MyChart help desk: 503-494-5252

Date							
Glucose							
BUN							
Creatinine *warning if increase of 0.2 or more							
Potassium							
Calcium							
Phosphorus							
Weekly magnesium							
WBC *warning if less than 3.0							
Hematocrit							
Drug Level							
Target Range							

Lab results

Lab name: _____

Medical record number: _____

Lab phone: 503-494-7383 MyChart help desk: 503-494-5252

Date							
Glucose							
BUN							
Creatinine *warning if increase of 0.2 or more							
Potassium							
Calcium							
Phosphorus							
Weekly magnesium							
WBC *warning if less than 3.0							
Hematocrit							
Drug Level							
Target Range							

Lab results

Lab name: _____

Medical record number: _____

Lab phone: 503-494-7383 MyChart help desk: 503-494-5252

Date							
Glucose							
BUN							
Creatinine *warning if increase of 0.2 or more							
Potassium							
Calcium							
Phosphorus							
Weekly magnesium							
WBC *warning if less than 3.0							
Hematocrit							
Drug Level							
Target Range							

Lab results

Lab name: _____

Medical record number: _____

Lab phone: 503-494-7383 MyChart help desk: 503-494-5252

Date							
Glucose							
BUN							
Creatinine *warning if increase of 0.2 or more							
Potassium							
Calcium							
Phosphorus							
Weekly magnesium							
WBC *warning if less than 3.0							
Hematocrit							
Drug Level							
Target Range							

Lab results

Lab name: _____

Medical record number: _____

Lab phone: 503-494-7383 MyChart help desk: 503-494-5252

Date							
Glucose							
BUN							
Creatinine *warning if increase of 0.2 or more							
Potassium							
Calcium							
Phosphorus							
Weekly magnesium							
WBC *warning if less than 3.0							
Hematocrit							
Drug Level							
Target Range							

Vital signs

Date							
Weight							
Temperature							
Blood pressure (morning)							
Blood pressure (evening)							
Heart rate							

Date							
Weight							
Temperature							
Blood pressure (morning)							
Blood pressure (evening)							
Heart rate							

Vital signs

Date							
Weight							
Temperature							
Blood pressure (morning)							
Blood pressure (evening)							
Heart rate							

Date							
Weight							
Temperature							
Blood pressure (morning)							
Blood pressure (evening)							
Heart rate							

Vital signs

Date							
Weight							
Temperature							
Blood pressure (morning)							
Blood pressure (evening)							
Heart rate							

Date							
Weight							
Temperature							
Blood pressure (morning)							
Blood pressure (evening)							
Heart rate							

Vital signs

Date							
Weight							
Temperature							
Blood pressure (morning)							
Blood pressure (evening)							
Heart rate							

Date							
Weight							
Temperature							
Blood pressure (morning)							
Blood pressure (evening)							
Heart rate							

Vital signs

Date							
Weight							
Temperature							
Blood pressure (morning)							
Blood pressure (evening)							
Heart rate							

Date							
Weight							
Temperature							
Blood pressure (morning)							
Blood pressure (evening)							
Heart rate							

Vital signs

Date							
Weight							
Temperature							
Blood pressure (morning)							
Blood pressure (evening)							
Heart rate							

Date							
Weight							
Temperature							
Blood pressure (morning)							
Blood pressure (evening)							
Heart rate							