

Oregon Health & Science University Hospital and Clinics Provider's Orders

ADULT AMBULATORY INFUSION ORDER
Alteplase (t-PA) Infusion for Central
Venous Catheters

Page 1 of 2

ACCOUNT NO.
MED. REC. NO.
NAME
BIRTHDATE

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.

Patient Identification

Weight:	_kg	Height:	cm
Allergies:			
Diagnosis Code:			
Treatment Start Date:		Patient to fo	ollow up with provider on date:

This plan will expire after 365 days at which time a new order will need to be placed

GUIDELINES FOR ORDERING

- 1. Send FACE SHEET and H&P or most recent chart note.
- 2. This plan is intended for patients with line access challenges after all other delegation protocols have been attempted.

NURSING ORDERS:

- 1. Refer to nursing and IV therapy guidelines for care of central venous catheters.
- 2. Follow facility policies and/or protocols for vascular access maintenance with appropriate flush solution, declotting (alteplase), and/or dressing changes.

MEDICATIONS:

□ alteplase (ACTIVASE) 2 mg in sodium chloride 0.9% infusion, 100 mL, intracatheter, ONCE over 4 hours (Maximum of 4 mg total in all lumens)

HYPERSENSITIVITY MEDICATIONS:

- NURSING COMMUNICATION If hypersensitivity or infusion reactions develop, temporarily hold the
 infusion and notify provider immediately. Administer emergency medications per the Treatment
 Algorithm for Acute Infusion Reaction (OHSU HC-PAT-133-GUD, HMC C-132). Refer to algorithm for
 symptom monitoring and continuously assess as grade of severity may progress.
- 2. diphenhydrAMINE (BENADRYL) injection, 25-50 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
- 3. EPINEPHrine HCI (ADRENALIN) injection, 0.3 mg, intramuscular, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
- 4. hydrocortisone sodium succinate (SOLU-CORTEF) injection, 100 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
- 5. famotidine (PEPCID) injection, 20 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction



Oregon Health & Science University Hospital and Clinics Provider's Orders

ADULT AMBULATORY INFUSION ORDER
Alteplase (t-PA) Infusion for Central
Venous Catheters
Page 2 of 2

ACCOUNT NO.
MED. REC. NO.
NAME
BIRTHDATE

Patient Identification

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.

I hold an active, unrestricted licen that corresponds with state where state if not Oregon); My physician license Number is	e pa se to you	tient (who is identified at the top of practice medicine in: provide care to patient and where (MUST BE (u (check box you are currently licensed. Specify	
medication described above for the			onzed by law to order initiation of the	
Provider signature: Date/Time:				
-		Phone:	Fax:	
	☑ Please indicate the patient's preferred clinic location below			
Phone (providers only)		BEAVERTON OHSU Knight Cancer Institute	15700 SW Greystone Court Beaverton OR 97006	
(971) 262-9645		NW PORTLAND Legacy Good Samaritan campus	Medical Office Building 3 – Suite 150 1130 NW 22nd Ave, Portland OR 97210	
Fax completed orders to (503) 346-8058		GRESHAM Legacy Mount Hood campus	Medical Office Building 3 – Suite 140 24988 SE Stark, Gresham OR 97030	
Infusion orders located at: www.ohsuknight.com/infusionorders		TUALATIN Legacy Meridian Park campus	Medical Office Building 2 – Suite 140 19260 SW 65th Ave, Tualatin OR 97062	