

Oregon Health & Science University Hospital and Clinics Provider's Orders



ADULT AMBULATORY INFUSION ORDER
Efgartigimod-Hyaluronidase-qvfc
(VYVGART HYTRULO) for Chronic
Inflammatory Demyelinating
Polyneuropathy

ACCOUNT NO.
MED. REC. NO.
NAME
BIRTHDATE

Patient Identification

Page 1 of 3

. «ge							
ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.							
Treatment Start Date:kg	Height:	ergies:cm					
REQUIRED ITEMS for all orders – necessary for insurance approval, scheduling, and patient safety 1. FACE SHEET with complete INSURANCE information and patient CONTACT information 2. Recent VISIT NOTE to support treatment (if not available in Epic) 3. LAB RESULTS for any required prescreening (if not available in Epic) 4. DIAGNOSIS CODE 5. Patient NAME and DATE OF BIRTH on EVERY page faxed							

GUIDELINES FOR ORDERING

- 1. Send FACE SHEET and H&P or most recent chart note.
- 2. Patients should be up to date with all immunizations before initiating therapy. Avoid the use of live vaccines in patients during treatment
- 3. Efgartigimod Alfa-fcab may increase the risk of infection. Delay treatment in patients with an active infection until the infection is resolved. Monitor for infection during treatment and consider withholding treatment if infection develops.
- 4. Do **NOT** substitute efgartigimod alfa/hyaluronidase (for SUBQ use) and efgartigimod alfa (for IV administration); products have different dosing and are **NOT** interchangeable

NURSING ORDERS:

- 1. TREATMENT PARAMETER Hold infusion and contact provider if patient has signs or symptoms of infection.
- 2. Subcutaneous injection: Monitor patient for signs and symptoms of hypersensitivity reactions during infusion and for 30 minutes following completion of injection.
- 3. Administer using 12-inch tubing, PVC winged set. Choose an injection site on abdomen a minimum of 2 to 3 inches from the naval, avoiding areas with moles or scars, or where skin is red, bruised or hard. Rotate injection sites for subsequent injections. Administer over a period of 30 to 90 seconds.

MEDICATIONS:

☐ Efgartigimod-hyaluronidase-qvfc (VYVGART HYTRULO) subcutaneous injection, 1008 mg, subcutaneous, ONCE WEEKLY



Oregon Health & Science University Hospital and Clinics Provider's Orders

ADULT AMBULATORY INFUSION ORDER

Efgartigimod-Hyaluronidase-qvfc (VYVGART HYTRULO) for Chronic Inflammatory Demyelinating Polyneuropathy ACCOUNT NO.
MED. REC. NO.
NAME
BIRTHDATE

Patient Identification

Page 2 of 3

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.

HYPERSENSITIVITY MEDICATIONS:

- NURSING COMMUNICATION If hypersensitivity or infusion reactions develop, temporarily hold the
 infusion and notify provider immediately. Administer emergency medications per the Treatment
 Algorithm for Acute Infusion Reaction (Policy HC-PAT-133-GUD, HMC C-132). Refer to algorithm for
 symptom monitoring and continuously assess as grade of severity may progress.
- 2. diphenhydrAMINE (BENADRYL) injection, 25-50 mg, intravenous, AS NEEDED x1 dose for hypersensitivity or infusion reaction.
- 3. famotidine (PEPCID) injection, 20 mg, intravenous, AS NEEDED x1 dose for hypersensitivity or infusion reaction.
- 4. hydrocortisone sodium succinate (SOLU-CORTEF) injection, 100 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction.
- 5. EPINEPHrine HCI (ADRENALIN) injection, 0.3 mg, intramuscular, AS NEEDED x1 dose for hypersensitivity or infusion reaction.

By signing below, I represent the following:

- I am responsible for the care of the patient identified on this form
- I hold an active, unrestricted license to practice medicine
- I am acting within my scope of practice and authorized by law to order the medication described above for the patient identified on this form

ALL ITEMS BELOW MUST BE COMPLETED TO BE A VALID PRESCRIPTION							
Signature:	License #:	Date:					
Print Name:	Phone:	Fax:					
Plan will expire 1 year after signature date at which time a new order will need to be placed							



Oregon Health & Science University Hospital and Clinics Provider's Orders

ADULT AMBULATORY INFUSION ORDER

Efgartigimod-Hyaluronidase-qvfc (VYVGART HYTRULO) for Chronic Inflammatory Demyelinating Polyneuropathy ACCOUNT NO.
MED. REC. NO.
NAME
BIRTHDATE

Patient Identification

Page 3 of 3

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.

OHSU Infusion Locations							
Contact the Referral Team	$\overline{\mathbf{A}}$	Please	ease indicate the patient's preferred clinic location below				
directly for assistance at the centralized numbers below (do not contact individual clinics		BEAVERTON OHSU Knight Cancer Institute		15700 SW Greystone Court Beaverton OR 97006			
INFUSION REFERRAL TEAM	□ NW PO		ORTLAND Good Samaritan campus	Medical Office Building 3 – Suite 150 1130 NW 22nd Ave, Portland OR 97210			
Fax completed orders to (503) 346-8058		□ GRESHAM Legacy Mount Hood campus		Medical Office Building 3 – Suite 140 24988 SE Stark, Gresham OR 97030			
Phone (providers only) (971) 262-9645		TUALATIN Legacy Meridian Park campus		Medical Office Building 2 – Suite 140 19260 SW 65th Ave, Tualatin OR 97062			
Infusion orders located at: www.ohsuknight.com/infusionorders	Community providers only (no Legacy) EAST PORTLAND Adventist Health Portland campus		PORTLAND	Pavilion – 10000 SE Main St – Suite 350 Portland, Oregon 97216			
Referral team will consider other locations as appropriate (e.g. selected site not available, urgent treatment, patient preference)							
OHSU Partner Infusion Locations							
Please indicate the patient's preferred clinic location below Not all therapies are offered at every site, contact site for more information							
Community providers only (no Legacy) HILLSBORO MEDICAL CENTER Fax completed orders to (503) 681-4120			364 SE 8th Ave – Medical Plaza Suite 108B Hillsboro, OR 97123 Phone (providers only) (503) 681-4124				
Community providers only (no Legacy) ADVENTIST HEALTH – PORTLAND Fax completed orders to (503) 261-6756		Infusion Services – 10123 SE Market St Portland, OR 97216 Phone (providers only) (503) 261-6631					