



Oregon Health & Science University
Hospital and Clinics Provider's Orders

PO7071



ADULT AMBULATORY INFUSION ORDER
Iron Sucrose (VENOFER) Infusion

Page 1 of 2

ACCOUNT NO.
MED. REC. NO.
NAME
BIRTHDATE

Patient Identification

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.

Weight: _____ kg Height: _____ cm

Allergies: _____

Diagnosis Code: _____

Treatment Start Date: _____ Patient to follow up with provider on date: _____

****This plan will expire after 365 days at which time a new order will need to be placed****

GUIDELINES FOR ORDERING

1. Send **FACE SHEET and H&P or most recent chart note.**
2. If patient is pregnant, estimated due date is: _____.
3. Provider must order and obtain a ferritin prior to patient being scheduled for iron infusion.
Labs drawn date: _____
Copy of ferritin must be attached.
4. Many insurance providers require a ferritin result within 90 days. If ferritin is not within 90 days of signed date then patient's insurance may deny coverage for this treatment.

NURSING ORDERS:

1. TREATMENT PARAMETER – Hold treatment and notify provider if Ferritin greater than 300 ng/mL.
2. Instruct patient to set follow up appointment with provider for follow up labs.
3. Follow facility policies and/or protocols for vascular access maintenance with appropriate flush solution, declothing (alteplase), and/or dressing changes.

MEDICATIONS:

iron sucrose (VENOFER): (must check one)

- ☐ 100 mg, intravenous, ONCE, administer slow IV injection over 5 minutes, x 5 doses over 14 days
- ☐ 200 mg, intravenous, ONCE, administer slow IV injection over 5 minutes, x 5 doses over 14 days
- ☐ 300 mg in sodium chloride 0.9% 250 mL, intravenous, ONCE, over 1.5 hours, x 3 doses
(administered every 2 to 3 days)

AS NEEDED MEDICATIONS:

1. sodium chloride 0.9%, 500 mL, intravenous, AS NEEDED x1 dose for vein discomfort. Give concurrently with iron sucrose

HYPERSENSITIVITY MEDICATIONS:

1. NURSING COMMUNICATION – If hypersensitivity or infusion reactions develop, temporarily hold the infusion and notify provider immediately. Administer emergency medications per the Treatment Algorithm for Acute Infusion Reaction (OHSU HC-PAT-133-GUD, HMC C-132). Refer to algorithm for symptom monitoring and continuously assess as grade of severity may progress.
2. NURSING COMMUNICATION – Avoid intravenous or oral diphenhydramine, move to next option in the algorithm. Adverse effects of diphenhydramine may overlap with IV iron adverse effects such as flushing, hypotension, tachycardia.



Oregon Health & Science University
Hospital and Clinics Provider's Orders

ADULT AMBULATORY INFUSION ORDER
Iron Sucrose (VENOFER) Infusion

Page 2 of 2

ACCOUNT NO.
MED. REC. NO.
NAME
BIRTHDATE

Patient Identification

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.

3. EPINEPHrine HCl (ADRENALIN) injection, 0.3 mg, intramuscular, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
4. hydrocortisone sodium succinate (SOLU-CORTEF) injection, 100 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
5. famotidine (PEPCID) injection, 20 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction

By signing below, I represent the following:

I am responsible for the care of the patient (*who is identified at the top of this form*);

I hold an active, unrestricted license to practice medicine in: ☐ Oregon ☐ _____ (check box that corresponds with state where you provide care to patient and where you are currently licensed. Specify state if not Oregon);

My physician license Number is # _____ (MUST BE COMPLETED TO BE A VALID PRESCRIPTION); and I am acting within my scope of practice and authorized by law to order Infusion of the medication described above for the patient identified on this form.

Provider signature: _____ **Date/Time:** _____

Printed Name: _____ **Phone:** _____ **Fax:** _____

Central Intake:

Phone: 971-262-9645 (providers only) Fax: 503-346-8058

Please check the appropriate box for the patient's preferred clinic location:

☐ **Beaverton**

OHSU Knight Cancer Institute
15700 SW Greystone Court
Beaverton, OR 97006
Phone number: 971-262-9000
Fax number: 503-346-8058

☐ **NW Portland**

Legacy Good Samaritan campus
Medical Office Building 3, Suite 150
1130 NW 22nd Ave
Portland, OR 97210
Phone number: 971-262-9600
Fax number: 503-346-8058

☐ **Gresham**

Legacy Mount Hood campus
Medical Office Building 3, Suite 140
24988 SE Stark
Gresham, OR 97030
Phone number: 971-262-9500
Fax number: 503-346-8058

☐ **Tualatin**

Legacy Meridian Park campus
Medical Office Building 2, Suite 140
19260 SW 65th Ave
Tualatin, OR 97062
Phone number: 971-262-9700
Fax number: 503-346-8058

Infusion orders located at: www.ohsuknight.com/infusionorders