

Oregon Health & Science University Hospital and Clinics Provider's Orders



ADULT AMBULATORY INFUSION ORDER Rezafungin (REZZAYO) Infusion

Page 1 of 3

ACCOUNT NO.
MED. REC. NO.
NAME
BIRTHDATE

				Patient Identification
				RKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.
Weigh	nt:	kg	Height:	cm
Allerg	ies:			<u> </u>
Diagn	osis Code:			_
Treatment Start Date: Patient t		Patie	ent to follow up with provider on date:	
This	s plan will exp	oire afte	r 365 days at w	hich time a new order will need to be placed
1. 2. 3.	Infusion-rela REZZAYO r sources of L	SHEET ited reac nay caus JV radiat	and H&P or mo tions have been se photosensitivi ion.	ost recent chart note. In reported during and following rezafungin administration. Itiy. Advise patients to use protection from sun exposure and other en established beyond 4 weekly doses.
LABS	S: CMP Routin	e, ONCE	E, every visit	
1.	prior to infus Follow facilit	S - Moni ion and y policie:	at the end of info	ols for vascular access maintenance with appropriate flush solution,
MEDI	CATIONS: (m	nust che	ck one)	
	ONCE, o Loading aintenance d Rezafun	on day 1 dose alro ose(s): gin (REZ	eady given on _	ım chloride 0.9%, 200 mg, intravenous, administer over 1 hour,
PRN	MEDICATIO	NS:		
		n (ZOFR	, ·	tegrating tablet, 4 mg, oral, ONCE AS NEEDED, for nausea and



Oregon Health & Science University Hospital and Clinics Provider's Orders

OHSU ADULT AMBULATORY INFUSION ORDER Health Rezafungin (REZZAYO) Infusion

Page 2 of 3

ACCOUNT NO.
MED. REC. NO.
NAME
BIRTHDATE

Patient Identification

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.

HYPERSENSITIVITY MEDICATIONS:

- NURSING COMMUNICATION If hypersensitivity or infusion reactions develop, temporarily hold the
 infusion and notify provider immediately. Administer emergency medications per the Treatment
 Algorithm for Acute Infusion Reaction (OHSU HC-PAT-133-GUD, HMC C-132). Refer to algorithm for
 symptom monitoring and continuously assess as grade of severity may progress.
- 2. diphenhydrAMINE (BENADRYL) injection, 25-50 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
- 3. EPINEPHrine HCI (ADRENALIN) injection, 0.3 mg, intramuscular, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
- 4. hydrocortisone sodium succinate (SOLU-CORTEF) injection, 100 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
- 5. famotidine (PEPCID) injection, 20 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction

By signing below, I represent the following I am responsible for the care of the patient (O	his form);
I hold an active, unrestricted license to pract that corresponds with state where you provistate if not Oregon);		
My physician license Number is #	y scope of practice and authori	
Provider signature:	Date/Tim	ne:



Oregon Health & Science University Hospital and Clinics Provider's Orders

OHSU ADULT AMBULATORY INFUSION ORDER Health Rezafungin (REZZAYO) Infusion

Page 3 of 3

ACCOUNT NO.
MED. REC. NO.
NAME
BIRTHDATE

Patient Identification

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.

Central Intake:

Phone: 971-262-9645 (providers only) Fax: 503-346-8058

Please check the appropriate box for the patient's preferred clinic location:

☐ Beaverton

OHSU Knight Cancer Institute 15700 SW Greystone Court Beaverton, OR 97006

Phone number: 971-262-9000 Fax number: 503-346-8058

☐ Gresham

Legacy Mount Hood campus Medical Office Building 3, Suite 140 24988 SE Stark Gresham, OR 97030

Phone number: 971-262-9500 Fax number: 503-346-8058

□ NW Portland

Legacy Good Samaritan campus Medical Office Building 3, Suite 150 1130 NW 22nd Ave. Portland, OR 97210

Phone number: 971-262-9600 Fax number: 503-346-8058

□ Tualatin

Legacy Meridian Park campus Medical Office Building 2, Suite 140 19260 SW 65th Ave. Tualatin, OR 97062

Phone number: 971-262-9700 Fax number: 503-346-8058

Infusion orders located at: www.ohsuknight.com/infusionorders