## Herceptin Hylecta® (trastuzumab and hyaluronidaseoysk)(Subcutaneous)

-E-

Document Number: OHSU HEALTHSERVICES-0544

Date Reviewed: 07/2025
Date of Origin: 06/02/2020

Dates Approved: 06/2020, 07/2020, 10/2020, 05/2021, 05/2022, 05/2023, 05/2024, 08/05/2025

### I. Length of Authorization <sup>1</sup>

Coverage is provided for 6 months and may be renewed (unless otherwise specified).

• Adjuvant therapy may be authorized for a total of fifty-two (52) weeks.

### II. Dosing Limits

#### Max Units (per dose and over time) [HCPCS Unit]:

• 60 billable units every 21 days

## III. Initial Approval Criteria 1-5

Coverage is provided in the following conditions:

Patient is at least 18 years of age; AND

#### **Universal Criteria**

- Left ventricular ejection fraction (LVEF) is within normal limits prior to initiating therapy and will be assessed at regular intervals (e.g., every 3 months) during treatment; AND
- Patient has cancer is human epidermal growth factor receptor 2 (HER2)-positive\* disease as
  determined by an FDA-approved or CLIA-compliant test\*; AND
- Females of reproductive potential have a negative pregnancy test prior to initiating treatment and will use effective contraception during treatment and for 7 months after the last dose; **AND**
- Therapy will not be substituted with or for ado-trastuzumab emtansine (Kadcyla) or famtrastuzumab deruxtecan-nxki (Enhertu); AND
- Therapy will not be used in combination with intravenous chemotherapy agents; AND

 Therapy will not be used in combination with trastuzumab (or any of its biosimilar products) or pertuzumab/trastuzumab and hyaluronidase-zzxf (Phesgo); AND

#### Breast Cancer\*\* † 1

- Used as adjuvant therapy; AND
  - Patient has ≥ T1 disease, node positive disease, or inflammatory disease (unless otherwise specified); AND
  - Used as a single agent following anthracycline-based therapy †; OR
- Used for metastatic disease; AND
  - Used as a single agent in patients who have received one or more prior chemotherapy regimens for metastatic disease †
  - \*\*NOTE: Coverage for Herceptin Hylecta will <u>NOT</u> encompass all FDA approved indications. Coverage will be <u>ONLY</u> provided when the above criteria is met.

#### FDA approved indications:

Adjuvant treatment of adults with HER2 overexpressing node positive or node negative (ER/PR negative or with one high risk feature) breast cancer:

- as part of a treatment regimen consisting of doxorubicin, cyclophosphamide, and either paclitaxel or docetaxel
- as part of a treatment regimen with docetaxel and carboplatin
- as a single agent following multi-modality anthracycline based therapy

#### Metastatic treatment of adults:

- In combination with paclitaxel for first-line treatment of HER2-overexpressing metastatic breast cancer
- As a single agent for treatment of HER2-overexpressing breast cancer in patients who have received one or more chemotherapy regimens for metastatic disease

#### \*HER2-positive overexpression criteria: 6,7

- Immunohistochemistry (IHC) assay 3+; OR
- Dual-probe in situ hybridization (ISH) assay HER2/CEP17 ratio ≥ 2.0 AND average HER2 copy number ≥ 4.0 signals/cell; OR
- Dual-probe in situ hybridization (ISH) assay AND concurrent IHC indicating one of the following:
  - HER2/CEP17 ratio ≥ 2.0 AND average HER2 copy number < 4.0 signals/cell AND concurrent IHC 3+; OR</li>
  - HER2/CEP17 ratio < 2.0 AND average HER2 copy number ≥ 6.0 signals/cell AND concurrent IHC 2+ or 3+; OR</li>
  - HER2/CEP17 ratio < 2.0 AND average HER2 copy number ≥ 4.0 and < 6.0 signals/cell AND concurrent IHC 3+</li>

Preferred therapies and recommendations are determined by review of clinical evidence. NCCN category of recommendation is taken into account as a component of this review. Regimens

deemed equally efficacious (i.e., those having the same NCCN categorization) are considered to be therapeutically equivalent.

❖ If confirmed using an immunotherapy assay - <a href="http://www.fda.gov/companiondiagnostics">http://www.fda.gov/companiondiagnostics</a>

† FDA Approved Indication(s); ‡ Compendia Recommended Indication(s); ♠ Orphan Drug

#### IV. Renewal Criteria <sup>1</sup>

Coverage may be renewed based upon the following criteria:

- Patient continues to meet the universal and other indication-specific relevant criteria such as concomitant therapy requirements (not including prerequisite therapy), performance status, etc. identified in section III; AND
- Duration of authorization has not been exceeded (refer to Section I); AND
- Disease response with treatment as defined by stabilization of disease or decrease in size of tumor or tumor spread; AND
- Absence of unacceptable toxicity from the drug. Examples of unacceptable toxicity include: cardiotoxicity (e.g., left ventricular dysfunction, cardiomyopathy, etc.), pulmonary toxicity (e.g., dyspnea, interstitial pneumonitis, etc.), neutropenia, severe administration-related reactions (e.g., hypersensitivity, anaphylaxis), etc.; AND
- Left ventricular ejection fraction (LVEF) within the previous 3 months as follows:
  - LVEF is within the institutional normal limits, and has not had an <u>absolute</u> decrease of ≥ 16% from pre-treatment baseline; **OR**
  - LVEF is below the institutional lower limits of normal, and has not had an <u>absolute</u> decrease of ≥ 10% from pre-treatment baseline

### V. Dosage/Administration <sup>1</sup>

Indication	Dose	
Breast Cancer	Administer 600 mg trastuzumab and 10,000 units hyaluronidase subcutaneously every three weeks.	
	<ul> <li>Adjuvant therapy: administer for 52 weeks or until disease recurrence</li> </ul>	
	Metastatic breast cancer: administer until disease progression	
Note:		

To be administered by a health care professional for subcutaneous use only in the thigh. Do not administer
intravenously.

## VI. Billing Code/Availability Information

**HCPCS Code:** 

OHSU Health Services ohsu.edu/healthshare Page | 3

- J9356 Injection, trastuzumab, 10 mg and hyaluronidase-oysk; 1 billable unit = 10 mg
   NDC:
- Herceptin Hylecta 600 mg/10,000 units (providing 600 mg trastuzumab and 10,000 units hyaluronidase per 5 mL) single-dose vials: 50242-0077-xx

### VII. References (STANDARD)

- 1. Herceptin Hylecta [package insert]. South San Francisco, CA; Genentech, Inc; June 2024. Accessed July 2025.
- 2. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) trastuzumab and hyaluronidase human. National Comprehensive Cancer Network, 2025. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed July 2025.
- 3. Jackisch C, Stroyakovskiy D, Pivot X, et al. Subcutaneous vs Intravenous Trastuzumab for Patients With ERBB2-Positive Early Breast Cancer: Final Analysis of the HannaH Phase 3 Randomized Clinical Trial. JAMA Oncol. 2019 May 1;5(5):e190339.
- 4. Gligorov J, Ataseven B, Verrill M, et al. Safety and tolerability of subcutaneous trastuzumab for the adjuvant treatment of human epidermal growth factor receptor 2-positive early breast cancer: SafeHer phase III study's primary analysis of 2573 patients. Eur J Cancer. 2017 Sep;82:237-246.
- 5. Pivot X, Verma S, Fallowfield L, et al. Efficacy and safety of subcutaneous trastuzumab and intravenous trastuzumab as part of adjuvant therapy for HER2-positive early breast cancer: Final analysis of the randomised, two-cohort PrefHer study. Eur J Cancer. 2017 Nov;86:82-90.
- 6. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines\*) for Breast Cancer 4.2025. National Comprehensive Cancer Network, 2025. NATIONAL COMPREHENSIVE CANCER NETWORK\*, NCCN\*, and NCCN GUIDELINES\* are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed July 2025.
- 7. Wolff AC, Hammond EH, Allison KH, et al. Human epidermal growth factor receptor 2 testing in breast cancer: American Society of Clinical Oncology/College of American Pathologists Clinical Practice Guideline Focused Update. J Clin Oncol 2018;36:2105-2122.

## VIII. References (ENHANCED)

**1e.** Woodward N, De Boer RH, Redfern A, et al. Results From the First Multicenter, Open-label, Phase IIIb Study Investigating the Combination of Pertuzumab With Subcutaneous Trastuzumab

OHSU Health Services ohsu.edu/healthshare Page | 4

- and a Taxane in Patients With HER2-positive Metastatic Breast Cancer (SAPPHIRE). Clin Breast Cancer. 2019;19(3):216–224. doi:10.1016/j.clbc.2019.02.008.
- 2e. Piccart-Gebhart MJ, Procter M, Leyland-Jones B, et al. Trastuzumab after adjuvant chemotherapy in HER2-positive breast cancer. N Engl J Med. 2005 Oct 20;353(16):1659-72.
- 3e. Smith I, Procter M, Gelber RD, et al. 2-year follow-up of trastuzumab after adjuvant chemotherapy in HER2-positive breast cancer: a randomised controlled trial. Lancet. 2007 Jan 6;369(9555):29-36.
- 4e. Prime Therapeutics Management. Trastuzumab SQ Clinical Literature Review Analysis. Last updated July 2025. Accessed July 2025.

## Appendix 1 – Covered Diagnosis Codes

ICD-10	ICD-10 Description	
C50.011	Malignant neoplasm of nipple and areola, right female breast	
C50.012	Malignant neoplasm of nipple and areola, left female breast	
C50.019	Malignant neoplasm of nipple and areola, unspecified female breast	
C50.021	Malignant neoplasm of nipple and areola, right female breast	
C50.022	Malignant neoplasm of nipple and areola, left female breast	
C50.029	Malignant neoplasm of nipple and areola, unspecified female breast	
C50.111	Malignant neoplasm of central portion of right female breast	
C50.112	Malignant neoplasm of central portion of left female breast	
C50.119	Malignant neoplasm of central portion of unspecified female breast	
C50.121	Malignant neoplasm of central portion of right male breast	
C50.122	Malignant neoplasm of central portion of left male breast	
C50.129	Malignant neoplasm of central portion of unspecified male breast	
C50.211	Malignant neoplasm of upper-inner quadrant of right female breast	
C50.212	Malignant neoplasm of upper-inner quadrant of left female breast	
C50.219	Malignant neoplasm of upper-inner quadrant of unspecified female breast	
C50.221	Malignant neoplasm of upper-inner quadrant of right male breast	
C50.222	Malignant neoplasm of upper-inner quadrant of left male breast	
C50.229	Malignant neoplasm of upper-inner quadrant of unspecified male breast	
C50.311	Malignant neoplasm of lower-inner quadrant of right female breast	
C50.312	Malignant neoplasm of lower-inner quadrant of left female breast	
C50.319	Malignant neoplasm of lower-inner quadrant of unspecified female breast	
C50.321	Malignant neoplasm of lower-inner quadrant of right male breast	

ICD-10	ICD-10 Description	
C50.322	Malignant neoplasm of lower-inner quadrant of left male breast	
C50.329	Malignant neoplasm of lower-inner quadrant of unspecified male breast	
C50.411	Malignant neoplasm of upper-outer quadrant of right female breast	
C50.412	Malignant neoplasm of upper-outer quadrant of left female breast	
C50.419	Malignant neoplasm of upper-outer quadrant of unspecified female breast	
C50.421	Malignant neoplasm of upper-outer quadrant of right male breast	
C50.422	Malignant neoplasm of upper-outer quadrant of left male breast	
C50.429	Malignant neoplasm of upper-outer quadrant of unspecified male breast	
C50.511	Malignant neoplasm of lower-outer quadrant of right female breast	
C50.512	Malignant neoplasm of lower-outer quadrant of left female breast	
C50.519	Malignant neoplasm of lower-outer quadrant of unspecified female breast	
C50.521	Malignant neoplasm of lower-outer quadrant of right male breast	
C50.522	Malignant neoplasm of lower-outer quadrant of left male breast	
C50.529	Malignant neoplasm of lower-outer quadrant of unspecified male breast	
C50.611	Malignant neoplasm of axillary tail of right female breast	
C50.612	Malignant neoplasm of axillary tail of left female breast	
C50.619	Malignant neoplasm of axillary tail of unspecified female breast	
C50.621	Malignant neoplasm of axillary tail of right male breast	
C50.622	Malignant neoplasm of axillary tail of left male breast	
C50.629	Malignant neoplasm of axillary tail of unspecified male breast	
C50.811	Malignant neoplasm of overlapping sites of right female breast	
C50.812	Malignant neoplasm of overlapping sites of left female breast	
C50.819	Malignant neoplasm of overlapping sites of unspecified female breast	
C50.821	Malignant neoplasm of overlapping sites of right male breast	
C50.822	Malignant neoplasm of overlapping sites of left male breast	
C50.829	Malignant neoplasm of overlapping sites of unspecified male breast	
C50.911	Malignant neoplasm of unspecified site of right female breast	
C50.912	Malignant neoplasm of unspecified site of left female breast	
C50.919	Malignant neoplasm of unspecified site of unspecified female breast	
C50.921	Malignant neoplasm of unspecified site of right male breast	
C50.922	Malignant neoplasm of unspecified site of left male breast	
C50.929	Malignant neoplasm of unspecified site of unspecified male breast	

ICD-10	ICD-10 Description	
Z85.3	Personal history of malignant neoplasm of breast	

### Appendix 2 – Centers for Medicare and Medicaid Services (CMS)

The preceding information is intended for non-Medicare coverage determinations. Medicare coverage for outpatient (Part B) drugs is outlined in the Medicare Benefit Policy Manual (Pub. 100-2), Chapter 15, §50 Drugs and Biologicals. In addition, National Coverage Determinations (NCDs) and/or Local Coverage Determinations (LCDs) may exist and compliance with these policies is required where applicable. Local Coverage Articles (LCAs) may also exist for claims payment purposes or to clarify benefit eligibility under Part B for drugs which may be self-administered. The following link may be used to search for NCD, LCD, or LCA documents: <a href="https://www.cms.gov/medicare-coverage-database/search.aspx">https://www.cms.gov/medicare-coverage-database/search.aspx</a>. Additional indications, including any preceding information, may be applied at the discretion of the health plan.

Medicare Part B Covered Diagnosis Codes (applicable to existing NCD/LCD/LCA): N/A

Medicare Part B Administrative Contractor (MAC) Jurisdictions				
Jurisdiction	Applicable State/US Territory	Contractor		
E (1)	CA,HI, NV, AS, GU, CNMI	Noridian Healthcare Solutions, LLC		
F (2 & 3)	AK, WA, OR, ID, ND, SD, MT, WY, UT, AZ	Noridian Healthcare Solutions, LLC		
5	KS, NE, IA, MO	Wisconsin Physicians Service Insurance Corp (WPS)		
6	MN, WI, IL	National Government Services, Inc. (NGS)		
H (4 & 7)	LA, AR, MS, TX, OK, CO, NM	Novitas Solutions, Inc.		
8	MI, IN	Wisconsin Physicians Service Insurance Corp (WPS)		
N (9)	FL, PR, VI	First Coast Service Options, Inc.		
J (10)	TN, GA, AL	Palmetto GBA		
M (11)	NC, SC, WV, VA (excluding below)	Palmetto GBA		
L (12)	DE, MD, PA, NJ, DC (includes Arlington & Fairfax counties and the city of Alexandria in VA)	Novitas Solutions, Inc.		
K (13 & 14)	NY, CT, MA, RI, VT, ME, NH	National Government Services, Inc. (NGS)		
15	ку, он	CGS Administrators, LLC		