



Oregon Health & Science University
Hospital and Clinics Provider's Orders

P07071



ADULT AMBULATORY INFUSION ORDER
Denosumab (PROLIA) Injection
Osteoporosis

Page 1 of 3

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.

ACCOUNT NO.
MED. REC. NO.
NAME
BIRTHDATE

Patient Identification

Weight: _____ kg Height: _____ cm

Allergies: _____

Diagnosis Code: _____

Treatment Start Date: _____ Patient to follow up with provider on date: _____

****This plan will expire after 365 days at which time a new order will need to be placed****

GUIDELINES FOR ORDERING

1. Send **FACE SHEET and H&P or most recent chart note**.
2. All patients should be prescribed daily calcium and vitamin D supplementation
3. In patients with severe renal impairment (creatinine clearance less than 30 mL/min), high risk of hypocalcemia, disturbances of mineral metabolism (e.g. hypoparathyroidism, thyroid surgery, parathyroid surgery, malabsorption syndromes, excision of small intestines) recommend clinical monitoring of calcium, magnesium and phosphorus levels within 14 days of Prolia injection.
4. Risk versus benefit regarding osteonecrosis of the jaw and hip fracture must be discussed prior to treatment.
5. A complete metabolic panel is recommended and a calcium level must be obtained within 60 days prior to starting treatment
6. The corrected calcium level should be greater than or equal to 8.4 mg/dL.
7. Pregnancy must be ruled out prior to administration. Perform pregnancy testing in all females of reproductive potential prior to administration of Prolia.
8. **Must complete and check the following box:**
 Provider confirms that the patient has had a recent oral or dental evaluation and/or has no contraindications to therapy related to dental issues prior to initiating therapy.

LABS:

- Complete metabolic panel, routine, ONCE, every visit

NURSING ORDERS:

1. Review previous creatinine clearance, serum calcium and albumin. If no results in past 60 days order CMP.
2. TREATMENT PARAMETER – Pharmacist to calculate corrected calcium. Hold and contact provider for corrected calcium less than 8.4 mg/dL.
3. Assess for new or unusual thigh, hip, groin, or jaw pain. Inform provider if positive findings or if patient is anticipating invasive dental work.
4. Do not hold treatment for CrCl less than 30 mL/min.
5. Remind patient to take at least 500 mg elemental calcium twice daily and 400 units vitamin D daily.

MEDICATIONS:

denosumab (PROLIA) injection, 60 mg, subcutaneous, every 6 months (26 weeks) x 2 doses,
Administer injection into upper arm, upper thigh, or abdomen



Oregon Health & Science University
Hospital and Clinics Provider's Orders

ADULT AMBULATORY INFUSION ORDER
Denosumab (PROLIA) Injection
Osteoporosis

Page 2 of 3

ACCOUNT NO.
MED. REC. NO.
NAME
BIRTHDATE

Patient Identification

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.

HYPERSensitivity MEDICATIONS:

1. **NURSING COMMUNICATION** – If hypersensitivity or infusion reactions develop, temporarily hold the infusion and notify provider immediately. Administer emergency medications per the Treatment Algorithm for Acute Infusion Reaction (OHSU HC-PAT-133-GUD, HMC C-132). Refer to algorithm for symptom monitoring and continuously assess as grade of severity may progress.
2. diphenhydrAMINE (BENADRYL) injection, 25-50 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
3. EPINEPHRine HCl (ADRENALIN) injection, 0.3 mg, intramuscular, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
4. hydrocortisone sodium succinate (SOLU-CORTEF) injection, 100 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
5. famotidine (PEPCID) injection, 20 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction

By signing below, I represent the following:

I am responsible for the care of the patient (*who is identified at the top of this form*);

I hold an active, unrestricted license to practice medicine in: Oregon _____ (check box that corresponds with state where you provide care to patient and where you are currently licensed. Specify state if not Oregon);

My physician license Number is # _____ **(MUST BE COMPLETED TO BE A VALID PRESCRIPTION);** and I am acting within my scope of practice and authorized by law to order Infusion of the medication described above for the patient identified on this form.

Provider signature: _____

Date/Time: _____

Printed Name: _____

Phone: _____

Fax: _____



Oregon Health & Science University

Hospital and Clinics Provider's Orders

ADULT AMBULATORY INFUSION ORDER

Denosumab (PROLIA) Injection
Osteoporosis

Page 3 of 3

ACCOUNT NO.

MED. REC. NO.

NAME

BIRTHDATE

Patient Identification

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.

Central Intake:

Phone: 971-262-9645 (providers only) Fax: 503-346-8058

Please check the appropriate box for the patient's preferred clinic location:

Beaverton

OHSU Knight Cancer Institute
15700 SW Greystone Court
Beaverton, OR 97006

Phone number: 971-262-9000

Fax number: 503-346-8058

NW Portland

Legacy Good Samaritan campus
Medical Office Building 3, Suite 150
1130 NW 22nd Ave
Portland, OR 97210

Phone number: 971-262-9600

Fax number: 503-346-8058

Gresham

Legacy Mount Hood campus
Medical Office Building 3, Suite 140
24988 SE Stark
Gresham, OR 97030

Phone number: 971-262-9500

Fax number: 503-346-8058

Tualatin

Legacy Meridian Park campus
Medical Office Building 2, Suite 140
19260 SW 65th Ave
Tualatin, OR 97062

Phone number: 971-262-9700

Fax number: 503-346-8058

Infusion orders located at: www.ohsuknight.com/infusionorders