



Oregon Health & Science University
Hospital and Clinics Provider's Orders

PO7071



ADULT AMBULATORY INFUSION ORDER
Ocrelizumab-hyaluronidase-ocsq
(OCREVUS ZUNOVO) Infusion

Page 1 of 3

Patient Identification

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.

Treatment Start Date: _____ Allergies: _____
Weight: _____ kg Height: _____ cm

REQUIRED ITEMS for all orders – necessary for insurance approval, scheduling, and patient safety

1. **FACE SHEET** with complete **INSURANCE** information and patient **CONTACT** information
2. Recent **VISIT NOTE** to support treatment (if not available in Epic)
3. **LAB RESULTS** for any required prescreening (if not available in Epic)
4. **DIAGNOSIS CODE**
5. **Patient NAME and DATE OF BIRTH** on **EVERY** page faxed

GUIDELINES FOR ORDERING

1. Send FACE SHEET and H&P or most recent chart note.
2. Hepatitis B (Hep B surface antigen and core antibody total) screening must be completed prior to initiation of treatment and the patient should not be infected. Please send results with order.
3. If patient is at high risk for TB exposure, a Tuberculin test must have been placed and read as negative prior to initiation of treatment (PPD or QuantiFERON Gold blood test). If result is indeterminate, a follow up chest X-ray must be performed to rule out TB. Please send results with order.
4. Please send results with order.

PRE-SCREENING

- Hepatitis B surface antigen and core antibody test results scanned with orders.
- Tuberculin skin test or QuantiFERON Gold blood test results scanned with orders if patient is at high risk for TB exposure.
- Chest X-Ray result scanned with orders if TB test result is indeterminate.
- TB screening is not necessary. Patient is not at high risk for TB exposure.

LABS:

- CBC with differential, Routine, Clinic Collect, ONCE, every visit
- Complete metabolic panel, Routine, Clinic Collect, ONCE, every visit

NURSING ORDERS:

1. **TREATMENT PARAMETER 1:** Hold treatment and contact provider if Hepatitis B surface antigen or core antibody total test result is positive, TB test result is positive, or if screening has not been performed.
2. RN to assess for active infection. If patient shows signs and symptoms of active infection or currently taking antibiotics. Hold treatment and notify provider.
3. For first injection, monitor patients for infusion-related reactions during injection and then for 1 hour after completion. For subsequent injections, monitor for 15 minutes after injection is complete. Lengthened monitoring recommended for previous infusion reactions, contact provider for guidance.
4. Do NOT substitute ocrelizumab (for IV administration) and ocrelizumab/hyaluronidase (for SUBQ administration); products have different dosing and are NOT interchangeable.
5. Allow refrigerated product to reach room temperature prior to administration.



**Oregon Health & Science University
Hospital and Clinics Provider's Orders**

**ADULT AMBULATORY INFUSION ORDER
Ocrelizumab-hyaluronidase-ocsq
(OCREVUS ZUNOVO) Infusion**

Page 2 of 3

ACCOUNT NO.
MED. REC. NO.
NAME
BIRTHDATE

Patient Identification

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.

6. Vial volume will account for dose volume of 23 mL plus volume needed to prime SUBQ infusion set. Withdraw entire contents of vial into a syringe using a 21-gauge transfer needle, attach an appropriate butterfly infusion set and prime the SUBQ infusion line with the drug product to eliminate air in the infusion line, stopping before the fluid reaches the needle. After priming, ensure the syringe contains exactly 23 mL of drug solution. Solution should be colorless to pale brown and clear to slightly opalescent.
7. Administer immediately to avoid needle clogging. Administer as subcutaneous injection over 10 minutes. Remaining priming volume in infusion tubing does NOT need to be administered.

PRE-MEDICATIONS: (Administer 30 minutes prior to infusion)

- acetaminophen (TYLENOL) tablet, 650 mg, oral, ONCE, every visit
- diphenhydRAMINE (BENADRYL) capsule, 50 mg, oral, ONCE, every visit. *Give either loratadine or diphenhydRAMINE, not both.*
- loratadine (CLARITIN) tablet, 10 mg, oral, ONCE AS NEEDED if diphenhydRAMINE is not given, every visit. *Give either loratadine or diphenhydRAMINE, not both.*
- dexAMETHasone tablet, 20 mg, oral, ONCE, every visit

MEDICATIONS: (must check one)

- Ocrelizumab-hyaluronidase-ocsq (OCREVUS ZUNOVO) injection, 920 mg, subcutaneously, ONCE, administer over 10 minutes, every 24 weeks.
Administer into the abdomen (avoid 2 inches around the navel) subcutaneously over approximately 10 minutes.

HYPERSensitivity MEDICATIONS:

1. NURSING COMMUNICATION – If hypersensitivity or infusion reactions develop, temporarily hold the infusion and notify provider immediately. Administer emergency medications per the Treatment Algorithm for Acute Infusion Reaction (Policy HC-CKT-133-GUD, Tuality C-132) . Refer to algorithm for symptom monitoring and continuously assess as grade of severity may progress.
2. DiphenhydRAMINE (BENADRYL) injection, 25–50 mg, intravenous, AS NEEDED X 1 dose for hypersensitivity reaction.
3. EPINEPPhrine HCl (ADRENALIN) injection, 0.3 mg, intramuscular, AS NEEDED x 1 dose for hypersensitivity reaction.
4. Hydrocortisone sodium succinate (SOLU-CORTEF) injection, 100 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity reaction. Dilute vial by either pressing chamber for Act-O-Vial or diluting powder vial with 2 mL SWFI or NS for injection.
5. Famotidine (PEPCID) injection, 20 mg, intravenous, AS NEEDED x 1 dose, for hypersensitivity reaction.
6. acetaminophen (TYLENOL) tablet, 650 mg, oral, EVERY 4 HOURS AS NEEDED for infusion related fever
7. meperidine (DEMEROL) injection, 25-50 mg, intravenous, EVERY 2 HOURS AS NEEDED for infusion-related severe rigors in the absence of hypotension, not to exceed 50 mg/hr.
8. sodium chloride 0.9% solution, 1000 mL, intravenous, AS NEEDED, Infuse at 200 mL/hr when infusion is stopped for emergency or PRN medications



Oregon Health & Science University
Hospital and Clinics Provider's Orders

ADULT AMBULATORY INFUSION ORDER
**Ocrelizumab-hyaluronidase-ocsq
(OCREVUS ZUNOVO) Infusion**

Page 3 of 3

ACCOUNT NO.
MED. REC. NO.
NAME
BIRTHDATE

Patient Identification

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.

By signing below, I represent the following:

- I am responsible for the care of the patient identified on this form
- I hold an active, unrestricted license to practice medicine
- I am acting within my scope of practice and authorized by law to order the medication described above for the patient identified on this form

ALL ITEMS BELOW MUST BE COMPLETED TO BE A VALID PRESCRIPTION

Signature: _____ **License #:** _____ **Date:** _____

Print Name: _____ **Phone:** _____ **Fax:** _____

Plan will expire 1 year after signature date at which time a new order will need to be placed

<p>Contact Referral Team directly for assistance at the centralized numbers below (do not fax/call individual clinics).</p> <p>INFUSION REFERRAL TEAM</p> <p>Fax completed orders to (503) 346-8058</p> <p>Phone (providers only) (971) 262-9645</p>	<input checked="" type="checkbox"/> Please indicate the patient's preferred clinic location below	
	<input type="checkbox"/> BEAVERTON OHSU Knight Cancer Institute	15700 SW Greystone Court Beaverton OR 97006
	<input type="checkbox"/> NW PORTLAND Legacy Good Samaritan campus	Medical Office Building 3 – Suite 150 1130 NW 22nd Ave, Portland OR 97210
	<input type="checkbox"/> GRESHAM Legacy Mount Hood campus	Medical Office Building 3 – Suite 140 24988 SE Stark, Gresham OR 97030
	<input type="checkbox"/> TUALATIN Legacy Meridian Park campus	Medical Office Building 2 – Suite 140 19260 SW 65th Ave, Tualatin OR 97062
	<input type="checkbox"/> Non-Legacy community providers only <input type="checkbox"/> EAST PORTLAND Adventist Health Portland campus	Pavilion – 10000 SE Main St – Suite 350 Portland, Oregon 97216
	Infusion orders located at: www.ohsuknight.com/infusionorders	