

Orencia® (abatacept) (Intravenous/Subcutaneous)

Document Number: OHSU HEALTHSERVICES-0091

Last Review Date: 03/04/2025

Date of Origin: 07/02/2010

Dates Reviewed: 07/2010, 09/2010, 12/2010, 02/2011, 03/2011, 06/2011, 09/2011, 12/2011, 03/2012, 06/2012, 09/2012, 12/2012, 03/2013, 06/2013, 09/2013, 12/2013, 03/2014, 06/2014, 09/2014, 12/2014, 03/2015, 06/2015, 09/2015, 12/2015, 03/2016, 06/2016, 09/2016, 12/2016, 03/2017, 04/2017, 06/2017, 08/2017, 09/2017, 12/2017, 03/2018, 06/2018, 08/2018, 08/2019, 01/2020, 08/2020, 08/2021, 01/2022, 08/2022, 08/2023, 12/2023, 08/2024, 03/2025

I. Length of Authorization

Initial coverage will be provided for 6 months and may be renewed annually thereafter, unless otherwise specified.

- Therapy for the Management of Immune Checkpoint Inhibitor-Related Toxicities and Prophylaxis for Acute Graft Versus Host Disease (aGVHD) may not be renewed.

II. Dosing Limits

Max Units (per dose and over time) [HCPCS Unit]:

- Management of Immune-Checkpoint Inhibitor-Related Toxicities
 - 675 billable units per 56 days
- Prophylaxis for aGVHD
 - 400 billable units per 29 days
- All other indications
 - Loading: 100 billable units at weeks 0, 2, & 4
 - Maintenance: 100 billable units per 4 weeks

III. Initial Approval Criteria ¹

Coverage is provided in the following conditions:

- Patient is at least 18 years of age (unless otherwise specified); **AND**
- Patient is up to date with all age-appropriate vaccinations, in accordance with current vaccination guidelines, prior to initiating therapy; **AND**

- Patient has been evaluated and screened for the presence of hepatitis B virus (HBV) prior to initiating treatment; **AND**

Universal Criteria¹

- Patient has been evaluated and screened for the presence of latent tuberculosis (TB) infection prior to initiating treatment and will receive ongoing monitoring for the presence of TB during treatment; **AND**
- Patient does not have an active infection, including clinically important localized infections; **AND**
- Patient will not receive live vaccines during therapy; **AND**
- Patient is not on concurrent treatment with another biologic therapy or targeted synthetic therapy unless otherwise specified; **AND**

Rheumatoid Arthritis (RA) †^{1,27,30}

- Physician has assessed baseline disease severity utilizing an objective measure/tool; **AND**
- Documented moderate to severe active disease; **AND**
 - Patient has had at least a 3-month trial and failed previous therapy with ONE conventional synthetic disease modifying anti-rheumatic drug (csDMARD) (e.g., methotrexate, azathioprine, auranofin, hydroxychloroquine, penicillamine, sulfasalazine, leflunomide, etc.); **OR**
 - Patient is already established on biologic or targeted synthetic therapy for the treatment of RA; **AND**
- May be used as a single agent or in combination with csDMARDs (e.g., methotrexate, hydroxychloroquine, leflunomide, sulfasalazine, etc.); **AND**

For Commercial Members Only

- Patient must try and have an inadequate response, contraindication, or intolerance to at least a three (3) month trial of ONE of the following: adalimumab*, Enbrel (etanercept), Xeljanz[‡] (tofacitinib), Rinvoq[‡] (upadacitinib), or tocilizumab SC**; **OR**
- Patient is continuing treatment

**Note: Preferred products are Hadlima (adalimumab-bwwd) and adalimumab-adaz*

‡ Note: Products are indicated in those who have had an inadequate response or intolerance to one or more TNF blockers

***Note: Preferred product is Tyenne SC (tocilizumab-aazg)*

For Medicaid Members Only

- Patient must try and have an inadequate response, contraindication, or intolerance to at least a three (3) month trial of ONE of the following: adalimumab*, Enbrel (etanercept), or tocilizumab SC**; **OR**

- Patient is continuing treatment

***Note: Preferred products are Hadlima (adalimumab-bwwd) and adalimumab-adaz**

****Note: Preferred product is Tyenne SC (tocilizumab-aazg)**

Polyarticular Juvenile Idiopathic Arthritis (pJIA) † 1,16

- Patient is at least 2 years of age (6 years of age for the IV formulation); **AND**
- Physician has assessed baseline disease severity utilizing an objective measure/tool; **AND**
- Documented moderate to severe active polyarticular disease; **AND**
 - Patient has had at least a 1-month trial and failure (unless contraindicated or intolerant) of previous therapy with either oral non-steroidal anti-inflammatory drugs (NSAIDs) OR a conventional synthetic disease-modifying anti-rheumatic drug (csDMARD) (e.g., methotrexate, leflunomide, sulfasalazine, etc.); **OR**
 - Patient is already established on biologic or targeted synthetic therapy for the treatment of pJIA; **AND**
- May be used as single agent or in combination with methotrexate; **AND**

For Commercial Members Only

- Patient must try and have an inadequate response, contraindication, or intolerance to at least a three (3) month trial of ONE of the following: adalimumab*, Enbrel (etanercept), Xeljanz[‡] (tofacitinib), Rinvoq[‡] (upadacitinib), or tocilizumab SC**; **OR**
- Patient is continuing treatment

***Note: Preferred products are Hadlima (adalimumab-bwwd) and adalimumab-adaz**

‡ Note: Products are indicated in those who have had an inadequate response or intolerance to one or more TNF blockers

****Note: Preferred product is Tyenne SC (tocilizumab-aazg)**

For Medicaid Members Only

- Patient must try and have an inadequate response, contraindication, or intolerance to at least a three (3) month trial of ONE of the following: adalimumab*, Enbrel (etanercept), or tocilizumab SC**; **OR**
- Patient is continuing treatment

***Note: Preferred products are Hadlima (adalimumab-bwwd) and adalimumab-adaz**

****Note: Preferred product is Tyenne SC (tocilizumab-aazg)**

Adult Psoriatic Arthritis (PsA) † 1,18,28,33,37,40

- Physician has assessed baseline disease severity utilizing an objective measure/tool; **AND**
- Documented moderate to severe active disease; **AND**

- For patients with predominantly axial disease OR enthesitis, a failure of at least a 4-week trial of ONE non-steroidal anti-inflammatory drug (NSAID), unless use is contraindicated; **OR**
- For patients with peripheral arthritis OR dactylitis, a failure of at least a 3-month trial of ONE conventional synthetic disease-modifying anti-rheumatic drug (csDMARD) (e.g., methotrexate, azathioprine, sulfasalazine, leflunomide, hydroxychloroquine, etc.); **OR**
- Patient is already established on biologic or targeted synthetic therapy for the treatment of PsA; **AND**
- May be used as a single agent or in combination with csDMARDs (e.g. methotrexate, hydroxychloroquine, leflunomide, sulfasalazine, etc.); **AND**

For Commercial Members Only

- Patient must try and have an inadequate response, contraindication, or intolerance to at least a three (3) month trial of ONE of the following: adalimumab*, Enbrel (etanercept), Cosentyx SC (secukinumab), ustekinumab SC^, Tremfya SC (guselkumab), Xeljanz[‡] (tofacitinib), Rinvoq[‡] (upadacitinib), Skyrizi SC (risankizumab), or Otezla (apremilast); **OR**
- Patient is continuing treatment

**Note: Preferred products are Hadlima (adalimumab-bwwd) and adalimumab-adaz*

^Note: Preferred products are Selarsdi SC (ustekinumab-aekn), Yesintek SC (ustekinumab-kfce), and Steqeyma SC (ustekinumab-stba)

‡ Note: Products are indicated in those who have had an inadequate response or intolerance to one or more TNF blockers

For Medicaid Members Only

- Patient must try and have an inadequate response, contraindication, or intolerance to at least a three (3) month trial of ONE of the following: adalimumab*, Enbrel (etanercept), Cosentyx SC (secukinumab), or ustekinumab SC^; **OR**
- Patient is continuing treatment

**Note: Preferred products are Hadlima (adalimumab-bwwd) and adalimumab-adaz*

^Note: Preferred products are Selarsdi SC (ustekinumab-aekn), Yesintek SC (ustekinumab-kfce), and Steqeyma SC (ustekinumab-stba)

Juvenile Psoriatic Arthritis (JPsA) † 1,4,16

- Patient is at least 2 years of age; **AND**
- Physician has assessed baseline disease severity utilizing an objective measure/tool; **AND**
- Documented moderate to severe active polyarticular disease; **AND**
- May be used as a single agent or in combination with methotrexate; **AND**

- Patient has had at least a 1-month trial and failure (unless contraindicated or intolerant) of previous therapy with either oral non-steroidal anti-inflammatory drugs (NSAIDs) OR a conventional synthetic disease-modifying anti-rheumatic drug (csDMARD) (e.g., methotrexate, leflunomide, sulfasalazine, etc.); **OR**
- Patient is already established on biologic or targeted synthetic therapy for the treatment of JPsA; **AND**

For Commercial Members Only

- For patients aged 6 years and older: patient must try and have an inadequate response, contraindication, or intolerance to at least a three (3) month trial of ONE of the following: Cosentyx SC (secukinumab), Enbrel (etanercept), Rinvoq[‡] (upadacitinib), or ustekinumab SC[^]; **OR**
- For patients aged 2 years and older: patient must try and have an inadequate response, contraindication, or intolerance to at least a three (3) month trial of ONE of the following: Cosentyx SC (secukinumab), Enbrel (etanercept), or Rinvoq[‡] (upadacitinib); **OR**
- Patient is continuing treatment

[^]Note: Preferred products are Selarsdi SC (ustekinumab-aekn), Yesintek SC (ustekinumab-kfce), and Steqeyma SC (ustekinumab-stba)

[‡]Note: Rinvoq (upadacitinib) is indicated in those who have had an inadequate response or intolerance to one or more TNF blockers

For Medicaid Members Only

- For patients aged 6 years and older: patient must try and have an inadequate response, contraindication, or intolerance to at least a three (3) month trial of ONE of the following: Cosentyx SC (secukinumab), Enbrel (etanercept), or ustekinumab SC[^]; **OR**
- For patients aged 2 years and older: patient must try and have an inadequate response, contraindication, or intolerance to at least a three (3) month trial of ONE of the following: Cosentyx SC (secukinumab) or Enbrel (etanercept); **OR**
- Patient is continuing treatment

[^]Note: Preferred products are Selarsdi SC (ustekinumab-aekn), Yesintek SC (ustekinumab-kfce), and Steqeyma SC (ustekinumab-stba)

Treatment of Chronic Graft Versus Host Disease (cGVHD) ‡^{1,19,20,22}

- Patient is post-allogeneic hematopoietic cell transplant (generally 3 or more months); **AND**
- Used as additional therapy in combination with systemic corticosteroids; **AND**
- Patient has no response (e.g., steroid-refractory disease) to first-line therapy options

Prophylaxis of Acute Graft Versus Host Disease (aGVHD) (IV formulation only) †^{1,35,36}

- Patient is undergoing a hematopoietic stem cell transplant (HSCT) from a matched or 1 allele-mismatched unrelated-donor; **AND**
- Patient is at least 2 years of age; **AND**
- Used in combination with a calcineurin inhibitor and methotrexate; **AND**
- Patient will receive antiviral prophylactic treatment for Epstein-Barr Virus (EBV) reactivation and prophylaxis will continue for 6 months post-transplantation; **AND**
- Patient will be monitored for both EBV reactivation and cytomegalovirus (CMV) infection/reactivation

Management of Immune Checkpoint Inhibitor-Related Toxicities (IV formulation only) †^{19,21,23}

- Patient has been receiving therapy with an immune checkpoint inhibitor; **AND**
 - Used as a single agent; **AND**
 - Used for the management of immunotherapy-related myocarditis; **AND**
 - Patient has had no improvement within 24-48 hours of starting high-dose methylprednisolone; **OR**
 - Used in combination with ruxolitinib; **AND**
 - Used for the management of immunotherapy-related concomitant myositis and myocarditis

† FDA Approved Indication(s); ‡ Compendia Recommended Indication(s); ◊ Orphan Drug

IV. Renewal Criteria ¹

Coverage can be renewed based upon the following criteria:

- Patient continues to meet the universal and other indication-specific relevant criteria identified in section III; **AND**
- Duration of authorization has not been exceeded (*refer to Section I*); **AND**
- Absence of unacceptable toxicity from the drug. Examples of unacceptable toxicity include: serious infections, severe hypersensitivity reactions, respiratory adverse events in those with predisposing conditions, reactivation of CMV or EBV, etc.; **AND**

Rheumatoid Arthritis ⁹⁻¹¹

- Disease response as indicated by improvement in signs and symptoms compared to baseline such as the number of tender and swollen joint counts, reduction of C reactive protein, improvement of patient global assessment, and/or an improvement on a disease activity scoring tool [e.g. an improvement on a composite scoring index such as Disease Activity Score-28 (DAS28) of 1.2 points or more or a ≥20% improvement on the American College of

Rheumatology-20 (ACR20) criteria, an improvement of disease severity on RAPID3 assessment, etc.].

Polyarticular Juvenile Idiopathic Arthritis ^{12,13}

- Disease response as indicated by improvement in signs and symptoms compared to baseline such as the number of tender and swollen joint counts, reduction of C-reactive protein, improvement of patient global assessment, and/or an improvement on a disease activity scoring tool [e.g., an improvement on a composite scoring index such as Juvenile Arthritis Disease Activity Score (JADAS) or the American College of Rheumatology (ACR) Pediatric (ACR-Pedi 30) of at least 30% improvement from baseline in three of six variables].

Adult Psoriatic Arthritis ^{14,34,41}

- Disease response as indicated by improvement in signs and symptoms compared to baseline such as the number of tender and swollen joint counts, reduction of C-reactive protein, improvement of patient global assessment, improvement on imaging (X-ray, ultrasound, or MRI), and/or an improvement on a disease activity scoring tool.

Juvenile Psoriatic Arthritis (JPsA) ^{1,12,13,41}

- Disease response as indicated by improvement in signs and symptoms compared to baseline such as the number of tender and swollen joint counts, reduction of C-reactive protein, improvement of patient global assessment, improvement on imaging (X-ray, ultrasound, or MRI), and/or an improvement on a disease activity scoring tool [e.g. an improvement on a composite scoring index such as Juvenile Arthritis Disease Activity Score (JADAS) or the American College of Rheumatology (ACR) Pediatric (ACR-Pedi 30) of at least 30% improvement from baseline in three of six variables].

Treatment of Chronic Graft Versus Host Disease (cGVHD) ^{20,22}

- Response to therapy with an improvement in one or more of the following:
 - Clinician assessments (e.g., NIH Skin Score, Upper GI Response Score, NIH Lung Symptom Score, etc.)
 - Patient-reported symptoms (e.g., Lee Symptom Scale, etc.)

V. Dosage/Administration ^{1,22,23,39}

Indication	Dose
Rheumatoid Arthritis	<p><u>Intravenous Dosing</u></p> <ul style="list-style-type: none"> • Weight < 60kg: <ul style="list-style-type: none"> ○ 500 mg at weeks 0, 2, & 4, then every 4 weeks thereafter • Weight 60 to 100 kg:

	<ul style="list-style-type: none"> ○ 750 mg at weeks 0, 2, & 4, then every 4 weeks thereafter ● Weight > 100 kg: <ul style="list-style-type: none"> ○ 1,000 mg at weeks 0, 2, & 4, then every 4 weeks thereafter <p><u>Subcutaneous Dosing (Orencia ClickJect or Orencia prefilled syringes)</u></p> <ul style="list-style-type: none"> ● Administer 125 mg by subcutaneous injection once weekly <ul style="list-style-type: none"> ○ May initiate with or without an IV loading dose ○ If initiated with an IV loading dose, a single weight-based IV infusion (see below) should be followed by the first 125 mg subcutaneous injection within a day <ul style="list-style-type: none"> ○ Weight < 60 kg: 500 mg ○ Weight 60 to 100 kg: 750 mg ○ Weight >100 kg: 1,000 mg <p>Patients transitioning from Orencia intravenous therapy to subcutaneous administration should administer the first subcutaneous dose instead of the next scheduled intravenous dose.</p>
<p>Adult Psoriatic Arthritis</p>	<p><u>Intravenous Dosing</u></p> <ul style="list-style-type: none"> ● Weight < 60kg: <ul style="list-style-type: none"> ○ 500 mg at weeks 0, 2, & 4, then every 4 weeks thereafter ● Weight 60 to 100 kg: <ul style="list-style-type: none"> ○ 750 mg at weeks 0, 2, & 4, then every 4 weeks thereafter ● Weight > 100 kg: <ul style="list-style-type: none"> ○ 1,000 mg at weeks 0, 2, & 4, then every 4 weeks thereafter <p><u>Subcutaneous Dosing (Orencia ClickJect or Orencia prefilled syringes)</u></p> <ul style="list-style-type: none"> ● Administer 125 mg by subcutaneous injection once weekly <ul style="list-style-type: none"> ○ No IV loading dose is needed <p>Patients transitioning from Orencia intravenous therapy to subcutaneous administration should administer the first subcutaneous dose instead of the next scheduled intravenous dose</p>
<p>Juvenile Psoriatic Arthritis</p>	<p><u>Subcutaneous Dosing (Orencia ClickJect or Orencia prefilled syringes)</u></p> <ul style="list-style-type: none"> ● Weight 10 to < 25 kg: 50 mg weekly ● Weight 25 to < 50 kg: 87.5 mg weekly ● Weight ≥ 50 kg: 125 mg weekly
<p>Polyarticular Juvenile Idiopathic Arthritis</p>	<p><u>Intravenous Dosing (patients aged 6 years or older ONLY)</u></p> <ul style="list-style-type: none"> ● Weight < 75 kg: <ul style="list-style-type: none"> ○ 10 mg/kg at weeks 0, 2, & 4, then every 4 weeks thereafter ● Weight 75 to 100 kg: <ul style="list-style-type: none"> ○ 750 mg at weeks 0, 2, & 4, then every 4 weeks thereafter ● Weight > 100 kg:

	<ul style="list-style-type: none"> ○ 1,000 mg at weeks 0, 2, & 4, then every 4 weeks thereafter <p><u>Subcutaneous Dosing (Orencia ClickJect or Orencia prefilled syringes and ONLY in patients aged 2 years or older)</u></p> <ul style="list-style-type: none"> ● Weight 10 to < 25 kg: 50 mg weekly ● Weight 25 to < 50 kg: 87.5 mg weekly ● Weight ≥ 50 kg: 125 mg weekly
Treatment of Chronic GVHD	<p><u>Intravenous/Subcutaneous Dosing</u></p> <ul style="list-style-type: none"> ● Up to 10 mg/kg (with a maximum dose of 1,000mg) at weeks 0, 2, & 4, then every 4 weeks thereafter until disease progression or unacceptable toxicity
Prophylaxis for Acute GVHD	<p><u>Intravenous Dosing</u></p> <ul style="list-style-type: none"> ● Patients ≥ 6 years old: 10 mg/kg (with a maximum dose of 1,000 mg) on the day before transplantation (Day -1), followed by administration on Days 5, 14, and 28 after transplantation ● Patients 2 to < 6 years old: 15 mg/kg on the day before transplantation (Day -1), followed by 12 mg/kg on Days 5, 14, and 28 after transplantation
Management of Immune Checkpoint Inhibitor-Related Toxicities	<p><u>Intravenous Dosing</u></p> <ul style="list-style-type: none"> ● Single agent: 500 mg every 2 weeks for a total of 5 doses ● In combination with ruxolitinib: Up to 20mg/kg on days 0, 5 and 14.

VI. Billing Code/Availability Information

HCPCS Code:

- J0129 – Injection, abatacept, 10 mg; 1 billable unit = 10 mg
(Code may be used for Medicare when drug is administered under the direct supervision of a physician; NOT for use when drug is self-administered)

NDC(s):

- Orencia 250 mg single-dose vial: 00003-2187-xx
- Orencia 125 mg/mL single-dose ClickJect™ Autoinjector: 00003-2188-xx
- Orencia 50 mg/0.4 mL single-dose prefilled syringe: 00003-2814-xx
- Orencia 87.5 mg/0.7 mL single-dose prefilled syringe: 00003-2818-xx
- Orencia 125 mg/mL single-dose prefilled syringe: 00003-2188-xx

VII. References

1. Orencia [package insert]. Princeton, NJ; Bristol-Myers Squibb; May 2024. Accessed January 2025.
2. Singh JA, Saag KG, Bridges SL Jr, et al. 2015 American College of Rheumatology Guideline for the Treatment of Rheumatoid Arthritis. Arthritis Care Res (Hoboken). 2015 Nov 6. doi: 10.1002/acr.22783.

3. Beukelman T, Patkar NM, Saag KG, et al. 2011 American College of Rheumatology recommendations for the treatment of juvenile idiopathic arthritis: initiation and safety monitoring of therapeutic agents for the treatment of arthritis and systemic features. *Arthritis Care Res (Hoboken)*. 2011 Apr;63(4):465-82.
4. Ringold S, Weiss PF, Beukelman T, et al. 2013 update of the 2011 American College of Rheumatology recommendations for the treatment of juvenile idiopathic arthritis: recommendations for the medical therapy of children with systemic juvenile idiopathic arthritis and tuberculosis screening among children receiving biologic medications. *Arthritis Rheum*. 2013 Oct;65(10):2499-512.
5. DeWitt EM, Kimura Y, Beukelman T, et al. Consensus treatment plans for new-onset systemic juvenile idiopathic arthritis. *Arthritis Care Res (Hoboken)*. 2012 Jul;64(7):1001-10.
6. Gottlieb A, Korman NJ, Gordon KB, et al. Guidelines of care for the management of psoriasis and psoriatic arthritis: Section 2. Psoriatic arthritis: overview and guidelines of care for treatment with an emphasis on the biologics. *J Am Acad Dermatol* 2008 May;58(5):851-64.
7. Gossec L, Smolen JS, Ramiro S, et al. European League Against Rheumatism (EULAR) recommendations for the management of psoriatic arthritis with pharmacological therapies: 2015 update. *Ann Rheum Dis*. 2015 Dec 7. pii: annrheumdis-2015-208337. doi: 10.1136/annrheumdis-2015-208337.
8. Smolen JS, Landewe R, Bijlsma J, et al. EULAR recommendations for the management of rheumatoid arthritis with synthetic and biological disease-modifying antirheumatic drugs: 2016 update. *Ann Rheum Dis*. 2017; 76: 935-938.
9. National Institute for Health and Care Excellence. NICE 2018. Rheumatoid Arthritis in Adults: Management. Published 11th July 2018. NICE Guideline [NG100]. <https://www.nice.org.uk/guidance/ng100/chapter/Recommendations>.
10. National Institute for Health and Care Excellence. NICE 2010. Adalimumab, etanercept, infliximab, rituximab and abatacept for the treatment of rheumatoid arthritis after failure of a TNF inhibitor. Published 10 October 2012. Clinical Guideline [TA195]. <https://www.nice.org.uk/guidance/ta195/resources/adalimumab-etanercept-infliximab-rituximab-and-abatacept-for-the-treatment-of-rheumatoid-arthritis-after-the-failure-of-a-tnf-inhibitor-pdf-82598558287813>.
11. Ward MM, Guthri LC, Alba MI. Rheumatoid Arthritis Response Criteria And Patient-Reported Improvement in Arthritis Activity: Is an ACR20 Response Meaningful to Patients?. *Arthritis Rheumatol*. 2014 Sep; 66(9): 2339–2343. doi: 10.1002/art.38705
12. Ringold S, Bittner R, Neggi T, et al. Performance of rheumatoid arthritis disease activity measures and juvenile arthritis disease activity scores in polyarticular-course juvenile idiopathic arthritis: Analysis of their ability to classify the American College of Rheumatology pediatric measures of response and the preliminary criteria for flare and inactive disease. *Arthritis Care Res (Hoboken)*. 2010 Aug;62(8):1095-102.
13. Consolaro A, Giancane G, Schiappapietra B, et al. Clinical outcome measures in juvenile idiopathic arthritis. *Pediatric Rheumatology* 18 April 2016 14:23.

14. National Institute for Health and Care Excellence. NICE 2017. Certolizumab pegol and secukinumab for treating active psoriatic arthritis after inadequate response to DMARDs. Published 24 May 2017. Technology Appraisal Guidance [TA445]. <https://www.nice.org.uk/guidance/ta445>. Accessed February 2025.
15. Ogdie A, Singh JA, Siegel E, Gladman DD, Husni ME. Treatment of psoriatic arthritis: a new ACR/NPF clinical guideline. Presented at: 2017 ACR/ARHP Annual Meeting; November 3-8, 2017; San Diego, CA. Scientific Session 5T064.
16. Ringold S, Angeles-Han ST, Beukelman T, et al. 2019 American College of Rheumatology/Arthritis Foundation Guideline for the Treatment of Juvenile Idiopathic Arthritis: Therapeutic Approaches for Non-Systemic Polyarthritis, Sacroiliitis, and Enthesitis. *Arthritis Care & Research*, Vol. 71, No. 6, June 2019, pp 717–734 DOI 10.1002/acr.23870.
17. Menter A, Strober BE, Kaplan DH, et al. Joint AAD-NPF guidelines of care for the management and treatment of psoriasis with biologics. *J Am Acad Dermatol* <https://doi.org/10.1016/j.jaad.2018.11.057>
18. Singh JA, Guyatt G, Ogdie A, et al. 2018 American College of Rheumatology/National Psoriasis Foundation Guideline for the Treatment of Psoriatic Arthritis. *Arthritis Rheumatol*. 2019 Jan;71(1):5-32. Doi: 10.1002/art.40726.
19. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) abatacept. National Comprehensive Cancer Network, 2025. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed February 2025.
20. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Hematopoietic Cell Transplantation (HCT). Version 2.2024. National Comprehensive Cancer Network, 2025. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed February 2025.
21. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Management of Immunotherapy-Related Toxicities. Version 1.2025. National Comprehensive Cancer Network, 2025. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed February 2025.
22. Nahas MR, Soiffer RJ, Kim HT, et al. Phase 1 clinical trial evaluating abatacept in patients with steroid-refractory chronic graft-versus-host disease. *Blood*. 2018;131(25):2836-2845. doi:10.1182/blood-2017-05-780239.

23. Salem JE, Allenbach Y, Vozy A, et al. Abatacept for Severe Immune Checkpoint Inhibitor-Associated Myocarditis. *N Engl J Med*. 2019;380(24):2377-2379. doi:10.1056/NEJMc1901677.
24. Vital EM, Emery P. Abatacept in the treatment of rheumatoid arthritis. *Ther Clin Risk Manag*. 2006;2(4):365-375. doi:10.2147/tcrm.2006.2.4.365.
25. Ruperto N, Lovell DJ, Quartier P, et al. Long-term safety and efficacy of abatacept in children with juvenile idiopathic arthritis. *Arthritis Rheum*. 2010;62(6):1792-1802. doi:10.1002/art.27431.
26. Mease P, Genovese MC, Gladstein G, et al. Abatacept in the treatment of patients with psoriatic arthritis: results of a six-month, multicenter, randomized, double-blind, placebo-controlled, phase II trial. *Arthritis Rheum*. 2011;63(4):939-948. doi:10.1002/art.30176.
27. Smolen JS, Landewé RBM, Bijlsma JWJ, et al. EULAR recommendations for the management of rheumatoid arthritis with synthetic and biological disease-modifying antirheumatic drugs: 2019 update. *Annals of the Rheumatic Diseases* 2020;79:685-699.
28. American Academy of Dermatology Work Group. Guidelines of care for the management of psoriasis and psoriatic arthritis: section 6. Guidelines of care for the treatment of psoriasis and psoriatic arthritis: case-based presentations and evidence-based conclusions. *J Am Acad Dermatol*. 2011 Jul;65(1):137-74.
29. Ramiro S, Smolen JS, Landewé R, et al. Pharmacological treatment of psoriatic arthritis: a systematic literature review for the 2015 update of the EULAR recommendations for the management of psoriatic arthritis. *Ann Rheum Dis* 2016;75:490-498 doi:10.1136/annrheumdis-2015-208466.
30. Fraenkel L, Bathon JM, England BR, et al. 2021 American College of Rheumatology Guideline for the Treatment of Rheumatoid Arthritis. *Arthritis Care Res (Hoboken)*. 2021 Jun 8. doi: 10.1002/acr.24596.
31. Genovese MC, Covarrubias A, Leon G, et al. Subcutaneous abatacept versus intravenous abatacept: a phase IIIb noninferiority study in patients with an inadequate response to methotrexate. *Arthritis Rheum*. 2011;63(10):2854-2864. doi:10.1002/art.30463.
32. Brunner HI, Tzaribachev N, Vega-Cornejo G, et al. Subcutaneous Abatacept in Patients With Polyarticular-Course Juvenile Idiopathic Arthritis: Results From a Phase III Open-Label Study. *Arthritis Rheumatol*. 2018;70(7):1144-1154. doi:10.1002/art.40466.
33. Gossec L, Baraliakos X, Kerschbaumer A, et al. EULAR recommendations for the management of psoriatic arthritis with pharmacological therapies: 2019 update. *Ann Rheum Dis*. 2020 Jun;79(6):700-712. doi: 10.1136/annrheumdis-2020-217159.
34. Mease PJ. Measures of psoriatic arthritis: Tender and Swollen Joint Assessment, Psoriasis Area and Severity Index (PASI), Nail Psoriasis Severity Index (NAPSI), Modified Nail Psoriasis Severity Index (mNAPSI), Mander/Newcastle Enthesitis Index (MEI), Leeds Enthesitis Index (LEI), Spondyloarthritis Research Consortium of Canada (SPARCC), Maastricht Ankylosing Spondylitis Enthesis Score (MASES), Leeds Dactylitis Index (LDI), Patient Global for Psoriatic Arthritis, Dermatology Life Quality Index (DLQI), Psoriatic Arthritis Quality of Life (PsAQOL), Functional Assessment of Chronic Illness Therapy-Fatigue (FACIT-F), Psoriatic Arthritis Response Criteria

(PsARC), Psoriatic Arthritis Joint Activity Index (PsAJAI), Disease Activity in Psoriatic Arthritis (DAPSA), and Composite Psoriatic Disease Activity Index (CPDAI). *Arthritis Care Res (Hoboken)*. 2011 Nov;63 Suppl 11:S64-85. doi: 10.1002/acr.20577.

35. Watkins B, Qayed M, McCracken C, et al. Phase II Trial of Costimulation Blockade With Abatacept for Prevention of Acute GVHD. *Journal of Clinical Oncology*. Volume 39, Issue 17

36. Kean L, Burns L, Kou T, et al. Improved Overall Survival of Patients Treated with Abatacept in Combination with a Calcineurin Inhibitor and Methotrexate Following 7/8 HLA-Matched Unrelated Allogeneic Hematopoietic Stem Cell Transplantation: Analysis of the Center for International Blood and Marrow Transplant Research Database. *Blood* (2021) 138 (Supplement 1): 3912. <https://doi.org/10.1182/blood-2021-150742>

37. Tucker L, Allen A, Chandler D, et al. The 2022 British Society for Rheumatology guideline for the treatment of psoriatic arthritis with biologic and targeted synthetic DMARDs. *Rheumatology (Oxford)*. 2022 Aug 30;61(9):e255-e266. doi: 10.1093/rheumatology/keac295.

38. Mease PJ, Gottlieb AB, van der Heijde D, et al. Efficacy and safety of abatacept, a T-cell modulator, in a randomised, double-blind, placebo-controlled, phase III study in psoriatic arthritis. *Ann Rheum Dis*. 2017 Sep;76(9):1550-1558. doi: 10.1136/annrheumdis-2016-210724

39. Salem JE, Bretagne M, Abbar B, et al. Abatacept/Ruxolitinib and Screening for Concomitant Respiratory Muscle Failure to Mitigate Fatality of Immune-Checkpoint Inhibitor Myocarditis. *Cancer Discov*. 2023 May 4;13(5):1100-1115. doi: 10.1158/2159-8290.CD-22-1180. PMID: 36815259.

40. Gossec L, Kerschbaumer A, Ferreira RJO, et al. EULAR recommendations for the management of psoriatic arthritis with pharmacological therapies: 2023 update. *Ann Rheum Dis*. 2024 May 15;83(6):706-719. doi: 10.1136/ard-2024-225531. PMID: 38499325

41. Tiwari V, Brent LH. Psoriatic Arthritis. [Updated 2024 Jan 7]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2024. <https://www.ncbi.nlm.nih.gov/books/NBK547710/>. Accessed February 04, 2025.

Appendix 1 – Covered Diagnosis Codes

ICD-10	ICD-10 Description
D89.810	Acute graft-versus-host disease
D89.811	Chronic graft-versus-host disease
D89.812	Acute on chronic graft-versus-host disease
D89.813	Graft-versus-host disease, unspecified
I30.8	Other forms of acute pericarditis
I30.9	Acute pericarditis, unspecified
I40.8	Other acute myocarditis
I40.9	Acute myocarditis, unspecified

ICD-10	ICD-10 Description
L40.50	Arthropathic psoriasis, unspecified
L40.51	Distal interphalangeal psoriatic arthropathy
L40.52	Psoriatic arthritis mutilans
L40.53	Psoriatic spondylitis
L40.54	Psoriatic juvenile arthropathy
L40.59	Other psoriatic arthropathy
M05.10	Rheumatoid lung disease with rheumatoid arthritis of unspecified site
M05.111	Rheumatoid lung disease with rheumatoid arthritis of right shoulder
M05.112	Rheumatoid lung disease with rheumatoid arthritis of left shoulder
M05.119	Rheumatoid lung disease with rheumatoid arthritis of unspecified shoulder
M05.121	Rheumatoid lung disease with rheumatoid arthritis of right elbow
M05.122	Rheumatoid lung disease with rheumatoid arthritis of left elbow
M05.129	Rheumatoid lung disease with rheumatoid arthritis of unspecified elbow
M05.131	Rheumatoid lung disease with rheumatoid arthritis of right wrist
M05.132	Rheumatoid lung disease with rheumatoid arthritis of left wrist
M05.139	Rheumatoid lung disease with rheumatoid arthritis of unspecified wrist
M05.141	Rheumatoid lung disease with rheumatoid arthritis of right hand
M05.142	Rheumatoid lung disease with rheumatoid arthritis of left hand
M05.149	Rheumatoid lung disease with rheumatoid arthritis of unspecified hand
M05.151	Rheumatoid lung disease with rheumatoid arthritis of right hip
M05.152	Rheumatoid lung disease with rheumatoid arthritis of left hip
M05.159	Rheumatoid lung disease with rheumatoid arthritis of unspecified hip
M05.161	Rheumatoid lung disease with rheumatoid arthritis of right knee
M05.162	Rheumatoid lung disease with rheumatoid arthritis of left knee
M05.169	Rheumatoid lung disease with rheumatoid arthritis of unspecified knee
M05.171	Rheumatoid lung disease with rheumatoid arthritis of right ankle and foot
M05.172	Rheumatoid lung disease with rheumatoid arthritis of left ankle and foot
M05.179	Rheumatoid lung disease with rheumatoid arthritis of unspecified ankle and foot
M05.19	Rheumatoid lung disease with rheumatoid arthritis of multiple sites
M05.20	Rheumatoid vasculitis with rheumatoid arthritis of unspecified site
M05.211	Rheumatoid vasculitis with rheumatoid arthritis of right shoulder
M05.212	Rheumatoid vasculitis with rheumatoid arthritis of left shoulder

ICD-10	ICD-10 Description
M05.219	Rheumatoid vasculitis with rheumatoid arthritis of unspecified shoulder
M05.221	Rheumatoid vasculitis with rheumatoid arthritis of right elbow
M05.222	Rheumatoid vasculitis with rheumatoid arthritis of left elbow
M05.229	Rheumatoid vasculitis with rheumatoid arthritis of unspecified elbow
M05.231	Rheumatoid vasculitis with rheumatoid arthritis of right wrist
M05.232	Rheumatoid vasculitis with rheumatoid arthritis of left wrist
M05.239	Rheumatoid vasculitis with rheumatoid arthritis of unspecified wrist
M05.241	Rheumatoid vasculitis with rheumatoid arthritis of right hand
M05.242	Rheumatoid vasculitis with rheumatoid arthritis of left hand
M05.249	Rheumatoid vasculitis with rheumatoid arthritis of unspecified hand
M05.251	Rheumatoid vasculitis with rheumatoid arthritis of right hip
M05.252	Rheumatoid vasculitis with rheumatoid arthritis of left hip
M05.259	Rheumatoid vasculitis with rheumatoid arthritis of unspecified hip
M05.261	Rheumatoid vasculitis with rheumatoid arthritis of right knee
M05.262	Rheumatoid vasculitis with rheumatoid arthritis of left knee
M05.269	Rheumatoid vasculitis with rheumatoid arthritis of unspecified knee
M05.271	Rheumatoid vasculitis with rheumatoid arthritis of right ankle and foot
M05.272	Rheumatoid vasculitis with rheumatoid arthritis of left ankle and foot
M05.279	Rheumatoid vasculitis with rheumatoid arthritis of unspecified ankle and foot
M05.29	Rheumatoid vasculitis with rheumatoid arthritis of multiple sites
M05.30	Rheumatoid heart disease with rheumatoid arthritis of unspecified site
M05.311	Rheumatoid heart disease with rheumatoid arthritis of right shoulder
M05.312	Rheumatoid heart disease with rheumatoid arthritis of left shoulder
M05.319	Rheumatoid heart disease with rheumatoid arthritis of unspecified shoulder
M05.321	Rheumatoid heart disease with rheumatoid arthritis of right elbow
M05.322	Rheumatoid heart disease with rheumatoid arthritis of left elbow
M05.329	Rheumatoid heart disease with rheumatoid arthritis of unspecified elbow
M05.331	Rheumatoid heart disease with rheumatoid arthritis of right wrist
M05.332	Rheumatoid heart disease with rheumatoid arthritis of left wrist
M05.339	Rheumatoid heart disease with rheumatoid arthritis of unspecified wrist
M05.341	Rheumatoid heart disease with rheumatoid arthritis of right hand
M05.342	Rheumatoid heart disease with rheumatoid arthritis of left hand

ICD-10	ICD-10 Description
M05.349	Rheumatoid heart disease with rheumatoid arthritis of unspecified hand
M05.351	Rheumatoid heart disease with rheumatoid arthritis of right hip
M05.352	Rheumatoid heart disease with rheumatoid arthritis of left hip
M05.359	Rheumatoid heart disease with rheumatoid arthritis of unspecified hip
M05.361	Rheumatoid heart disease with rheumatoid arthritis of right knee
M05.362	Rheumatoid heart disease with rheumatoid arthritis of left knee
M05.369	Rheumatoid heart disease with rheumatoid arthritis of unspecified knee
M05.371	Rheumatoid heart disease with rheumatoid arthritis of right ankle and foot
M05.372	Rheumatoid heart disease with rheumatoid arthritis of left ankle and foot
M05.379	Rheumatoid heart disease with rheumatoid arthritis of unspecified ankle and foot
M05.39	Rheumatoid heart disease with rheumatoid arthritis of multiple sites
M05.40	Rheumatoid myopathy with rheumatoid arthritis of unspecified site
M05.411	Rheumatoid myopathy with rheumatoid arthritis of right shoulder
M05.412	Rheumatoid myopathy with rheumatoid arthritis of left shoulder
M05.419	Rheumatoid myopathy with rheumatoid arthritis of unspecified shoulder
M05.421	Rheumatoid myopathy with rheumatoid arthritis of right elbow
M05.422	Rheumatoid myopathy with rheumatoid arthritis of left elbow
M05.429	Rheumatoid myopathy with rheumatoid arthritis of unspecified elbow
M05.431	Rheumatoid myopathy with rheumatoid arthritis of right wrist
M05.432	Rheumatoid myopathy with rheumatoid arthritis of left wrist
M05.439	Rheumatoid myopathy with rheumatoid arthritis of unspecified wrist
M05.441	Rheumatoid myopathy with rheumatoid arthritis of right hand
M05.442	Rheumatoid myopathy with rheumatoid arthritis of left hand
M05.449	Rheumatoid myopathy with rheumatoid arthritis of unspecified hand
M05.451	Rheumatoid myopathy with rheumatoid arthritis of right hip
M05.452	Rheumatoid myopathy with rheumatoid arthritis of left hip
M05.459	Rheumatoid myopathy with rheumatoid arthritis of unspecified hip
M05.461	Rheumatoid myopathy with rheumatoid arthritis of right knee
M05.462	Rheumatoid myopathy with rheumatoid arthritis of left knee
M05.469	Rheumatoid myopathy with rheumatoid arthritis of unspecified knee
M05.471	Rheumatoid myopathy with rheumatoid arthritis of right ankle and foot
M05.472	Rheumatoid myopathy with rheumatoid arthritis of left ankle and foot

ICD-10	ICD-10 Description
M05.479	Rheumatoid myopathy with rheumatoid arthritis of unspecified ankle and foot
M05.49	Rheumatoid myopathy with rheumatoid arthritis of multiple sites
M05.50	Rheumatoid polyneuropathy with rheumatoid arthritis of unspecified site
M05.511	Rheumatoid polyneuropathy with rheumatoid arthritis of right shoulder
M05.512	Rheumatoid polyneuropathy with rheumatoid arthritis of left shoulder
M05.519	Rheumatoid polyneuropathy with rheumatoid arthritis of unspecified shoulder
M05.521	Rheumatoid polyneuropathy with rheumatoid arthritis of right elbow
M05.522	Rheumatoid polyneuropathy with rheumatoid arthritis of left elbow
M05.529	Rheumatoid polyneuropathy with rheumatoid arthritis of unspecified elbow
M05.531	Rheumatoid polyneuropathy with rheumatoid arthritis of right wrist
M05.532	Rheumatoid polyneuropathy with rheumatoid arthritis of left wrist
M05.539	Rheumatoid polyneuropathy with rheumatoid arthritis of unspecified wrist
M05.541	Rheumatoid polyneuropathy with rheumatoid arthritis of right hand
M05.542	Rheumatoid polyneuropathy with rheumatoid arthritis of left hand
M05.549	Rheumatoid polyneuropathy with rheumatoid arthritis of unspecified hand
M05.551	Rheumatoid polyneuropathy with rheumatoid arthritis of right hip
M05.552	Rheumatoid polyneuropathy with rheumatoid arthritis of left hip
M05.559	Rheumatoid polyneuropathy with rheumatoid arthritis of unspecified hip
M05.561	Rheumatoid polyneuropathy with rheumatoid arthritis of right knee
M05.562	Rheumatoid polyneuropathy with rheumatoid arthritis of left knee
M05.569	Rheumatoid polyneuropathy with rheumatoid arthritis of unspecified knee
M05.571	Rheumatoid polyneuropathy with rheumatoid arthritis of right ankle and foot
M05.572	Rheumatoid polyneuropathy with rheumatoid arthritis of left ankle and foot
M05.579	Rheumatoid polyneuropathy with rheumatoid arthritis of unspecified ankle and foot
M05.59	Rheumatoid polyneuropathy with rheumatoid arthritis of multiple sites
M05.60	Rheumatoid arthritis of unspecified site with involvement of other organs and systems
M05.611	Rheumatoid arthritis of right shoulder with involvement of other organs and systems
M05.612	Rheumatoid arthritis of left shoulder with involvement of other organs and systems
M05.619	Rheumatoid arthritis of unspecified shoulder with involvement of other organs and systems
M05.621	Rheumatoid arthritis of right elbow with involvement of other organs and systems
M05.622	Rheumatoid arthritis of left elbow with involvement of other organs and systems
M05.629	Rheumatoid arthritis of unspecified elbow with involvement of other organs and systems

ICD-10	ICD-10 Description
M05.631	Rheumatoid arthritis of right wrist with involvement of other organs and systems
M05.632	Rheumatoid arthritis of left wrist with involvement of other organs and systems
M05.639	Rheumatoid arthritis of unspecified wrist with involvement of other organs and systems
M05.641	Rheumatoid arthritis of right hand with involvement of other organs and systems
M05.642	Rheumatoid arthritis of left hand with involvement of other organs and systems
M05.649	Rheumatoid arthritis of unspecified hand with involvement of other organs and systems
M05.651	Rheumatoid arthritis of right hip with involvement of other organs and systems
M05.652	Rheumatoid arthritis of left hip with involvement of other organs and systems
M05.659	Rheumatoid arthritis of unspecified hip with involvement of other organs and systems
M05.661	Rheumatoid arthritis of right knee with involvement of other organs and systems
M05.662	Rheumatoid arthritis of left knee with involvement of other organs and systems
M05.669	Rheumatoid arthritis of unspecified knee with involvement of other organs and systems
M05.671	Rheumatoid arthritis of right ankle and foot with involvement of other organs and systems
M05.672	Rheumatoid arthritis of left ankle and foot with involvement of other organs and systems
M05.679	Rheumatoid arthritis of unspecified ankle and foot with involvement of other organs and systems
M05.69	Rheumatoid arthritis of multiple sites with involvement of other organs and systems
M05.7A	Rheumatoid arthritis with rheumatoid factor of other specified site without organ or systems involvement
M05.711	Rheumatoid arthritis with rheumatoid factor of right shoulder without organ or systems involvement
M05.712	Rheumatoid arthritis with rheumatoid factor of left shoulder without organ or systems involvement
M05.719	Rheumatoid arthritis with rheumatoid factor of unspecified shoulder without organ or systems involvement
M05.721	Rheumatoid arthritis with rheumatoid factor of right elbow without organ or systems involvement
M05.722	Rheumatoid arthritis with rheumatoid factor of left elbow without organ or systems involvement
M05.729	Rheumatoid arthritis with rheumatoid factor of unspecified elbow without organ or systems involvement
M05.731	Rheumatoid arthritis with rheumatoid factor of right wrist without organ or systems involvement
M05.732	Rheumatoid arthritis with rheumatoid factor of left wrist without organ or systems involvement
M05.739	Rheumatoid arthritis with rheumatoid factor of unspecified wrist without organ or systems involvement
M05.741	Rheumatoid arthritis with rheumatoid factor of right hand without organ or systems involvement
M05.742	Rheumatoid arthritis with rheumatoid factor of left hand without organ or systems involvement
M05.749	Rheumatoid arthritis with rheumatoid factor of unspecified hand without organ or systems involvement
M05.751	Rheumatoid arthritis with rheumatoid factor of right hip without organ or systems involvement
M05.752	Rheumatoid arthritis with rheumatoid factor of left hip without organ or systems involvement
M05.759	Rheumatoid arthritis with rheumatoid factor of unspecified hip without organ or systems involvement

ICD-10	ICD-10 Description
M05.761	Rheumatoid arthritis with rheumatoid factor of right knee without organ or systems involvement
M05.762	Rheumatoid arthritis with rheumatoid factor of left knee without organ or systems involvement
M05.769	Rheumatoid arthritis with rheumatoid factor of unspecified knee without organ or systems involvement
M05.771	Rheumatoid arthritis with rheumatoid factor of right ankle and foot without organ or systems involvement
M05.772	Rheumatoid arthritis with rheumatoid factor of left ankle and foot without organ or systems involvement
M05.779	Rheumatoid arthritis with rheumatoid factor of unspecified ankle and foot without organ or systems involvement
M05.79	Rheumatoid arthritis with rheumatoid factor of multiple sites without organ or systems involvement
M05.8A	Other rheumatoid arthritis with rheumatoid factor of other specified site
M05.811	Other rheumatoid arthritis with rheumatoid factor of right shoulder
M05.812	Other rheumatoid arthritis with rheumatoid factor of left shoulder
M05.819	Other rheumatoid arthritis with rheumatoid factor of unspecified shoulder
M05.821	Other rheumatoid arthritis with rheumatoid factor of right elbow
M05.822	Other rheumatoid arthritis with rheumatoid factor of left elbow
M05.829	Other rheumatoid arthritis with rheumatoid factor of unspecified elbow
M05.831	Other rheumatoid arthritis with rheumatoid factor of right wrist
M05.832	Other rheumatoid arthritis with rheumatoid factor of left wrist
M05.839	Other rheumatoid arthritis with rheumatoid factor of unspecified wrist
M05.841	Other rheumatoid arthritis with rheumatoid factor of right hand
M05.842	Other rheumatoid arthritis with rheumatoid factor of left hand
M05.849	Other rheumatoid arthritis with rheumatoid factor of unspecified hand
M05.851	Other rheumatoid arthritis with rheumatoid factor of right hip
M05.852	Other rheumatoid arthritis with rheumatoid factor of left hip
M05.859	Other rheumatoid arthritis with rheumatoid factor of unspecified hip
M05.861	Other rheumatoid arthritis with rheumatoid factor of right knee
M05.862	Other rheumatoid arthritis with rheumatoid factor of left knee
M05.869	Other rheumatoid arthritis with rheumatoid factor of unspecified knee
M05.871	Other rheumatoid arthritis with rheumatoid factor of right ankle and foot
M05.872	Other rheumatoid arthritis with rheumatoid factor of left ankle and foot
M05.879	Other rheumatoid arthritis with rheumatoid factor of unspecified ankle and foot
M05.89	Other rheumatoid arthritis with rheumatoid factor of multiple sites
M05.9	Rheumatoid arthritis with rheumatoid factor, unspecified
M06.0A	Rheumatoid arthritis without rheumatoid factor, other specified site

ICD-10	ICD-10 Description
M06.011	Rheumatoid arthritis without rheumatoid factor, right shoulder
M06.012	Rheumatoid arthritis without rheumatoid factor, left shoulder
M06.019	Rheumatoid arthritis without rheumatoid factor, unspecified shoulder
M06.021	Rheumatoid arthritis without rheumatoid factor, right elbow
M06.022	Rheumatoid arthritis without rheumatoid factor, left elbow
M06.029	Rheumatoid arthritis without rheumatoid factor, unspecified elbow
M06.031	Rheumatoid arthritis without rheumatoid factor, right wrist
M06.032	Rheumatoid arthritis without rheumatoid factor, left wrist
M06.039	Rheumatoid arthritis without rheumatoid factor, unspecified wrist
M06.041	Rheumatoid arthritis without rheumatoid factor, right hand
M06.042	Rheumatoid arthritis without rheumatoid factor, left hand
M06.049	Rheumatoid arthritis without rheumatoid factor, unspecified hand
M06.051	Rheumatoid arthritis without rheumatoid factor, right hip
M06.052	Rheumatoid arthritis without rheumatoid factor, left hip
M06.059	Rheumatoid arthritis without rheumatoid factor, unspecified hip
M06.061	Rheumatoid arthritis without rheumatoid factor, right knee
M06.062	Rheumatoid arthritis without rheumatoid factor, left knee
M06.069	Rheumatoid arthritis without rheumatoid factor, unspecified knee
M06.071	Rheumatoid arthritis without rheumatoid factor, right ankle and foot
M06.072	Rheumatoid arthritis without rheumatoid factor, left ankle and foot
M06.079	Rheumatoid arthritis without rheumatoid factor, unspecified ankle and foot
M06.08	Rheumatoid arthritis without rheumatoid factor, vertebrae
M06.09	Rheumatoid arthritis without rheumatoid factor, multiple sites
M06.8A	Other specified rheumatoid arthritis, other specified site
M06.811	Other specified rheumatoid arthritis, right shoulder
M06.812	Other specified rheumatoid arthritis, left shoulder
M06.819	Other specified rheumatoid arthritis, unspecified shoulder
M06.821	Other specified rheumatoid arthritis, right elbow
M06.822	Other specified rheumatoid arthritis, left elbow
M06.829	Other specified rheumatoid arthritis, unspecified elbow
M06.831	Other specified rheumatoid arthritis, right wrist
M06.832	Other specified rheumatoid arthritis, left wrist

ICD-10	ICD-10 Description
M06.839	Other specified rheumatoid arthritis, unspecified wrist
M06.841	Other specified rheumatoid arthritis, right hand
M06.842	Other specified rheumatoid arthritis, left hand
M06.849	Other specified rheumatoid arthritis, unspecified hand
M06.851	Other specified rheumatoid arthritis, right hip
M06.852	Other specified rheumatoid arthritis, left hip
M06.859	Other specified rheumatoid arthritis, unspecified hip
M06.861	Other specified rheumatoid arthritis, right knee
M06.862	Other specified rheumatoid arthritis, left knee
M06.869	Other specified rheumatoid arthritis, unspecified knee
M06.871	Other specified rheumatoid arthritis, right ankle and foot
M06.872	Other specified rheumatoid arthritis, left ankle and foot
M06.879	Other specified rheumatoid arthritis, unspecified ankle and foot
M06.88	Other specified rheumatoid arthritis, vertebrae
M06.89	Other specified rheumatoid arthritis, multiple sites
M06.9	Rheumatoid arthritis, unspecified
M08.0A	Unspecified juvenile rheumatoid arthritis, other specified site
M08.011	Unspecified juvenile rheumatoid arthritis, right shoulder
M08.012	Unspecified juvenile rheumatoid arthritis, left shoulder
M08.019	Unspecified juvenile rheumatoid arthritis, unspecified shoulder
M08.021	Unspecified juvenile rheumatoid arthritis, right elbow
M08.022	Unspecified juvenile rheumatoid arthritis, left elbow
M08.029	Unspecified juvenile rheumatoid arthritis, unspecified elbow
M08.031	Unspecified juvenile rheumatoid arthritis, right wrist
M08.032	Unspecified juvenile rheumatoid arthritis, left wrist
M08.039	Unspecified juvenile rheumatoid arthritis, unspecified wrist
M08.041	Unspecified juvenile rheumatoid arthritis, right hand
M08.042	Unspecified juvenile rheumatoid arthritis, left hand
M08.049	Unspecified juvenile rheumatoid arthritis, unspecified hand
M08.051	Unspecified juvenile rheumatoid arthritis, right hip
M08.052	Unspecified juvenile rheumatoid arthritis, left hip
M08.059	Unspecified juvenile rheumatoid arthritis, unspecified hip

ICD-10	ICD-10 Description
M08.061	Unspecified juvenile rheumatoid arthritis, right knee
M08.062	Unspecified juvenile rheumatoid arthritis, left knee
M08.069	Unspecified juvenile rheumatoid arthritis, unspecified knee
M08.071	Unspecified juvenile rheumatoid arthritis, right ankle and foot
M08.072	Unspecified juvenile rheumatoid arthritis, left ankle and foot
M08.079	Unspecified juvenile rheumatoid arthritis, unspecified ankle and foot
M08.08	Unspecified juvenile rheumatoid arthritis, vertebrae
M08.09	Unspecified juvenile rheumatoid arthritis, multiple sites
M08.2A	Juvenile rheumatoid arthritis with systemic onset, other specified site
M08.211	Juvenile rheumatoid arthritis with systemic onset, right shoulder
M08.212	Juvenile rheumatoid arthritis with systemic onset, left shoulder
M08.219	Juvenile rheumatoid arthritis with systemic onset, unspecified shoulder
M08.221	Juvenile rheumatoid arthritis with systemic onset, right elbow
M08.222	Juvenile rheumatoid arthritis with systemic onset, left elbow
M08.229	Juvenile rheumatoid arthritis with systemic onset, unspecified elbow
M08.231	Juvenile rheumatoid arthritis with systemic onset, right wrist
M08.232	Juvenile rheumatoid arthritis with systemic onset, left wrist
M08.239	Juvenile rheumatoid arthritis with systemic onset, unspecified wrist
M08.241	Juvenile rheumatoid arthritis with systemic onset, right hand
M08.242	Juvenile rheumatoid arthritis with systemic onset, left hand
M08.249	Juvenile rheumatoid arthritis with systemic onset, unspecified hand
M08.251	Juvenile rheumatoid arthritis with systemic onset, right hip
M08.252	Juvenile rheumatoid arthritis with systemic onset, left hip
M08.259	Juvenile rheumatoid arthritis with systemic onset, unspecified hip
M08.261	Juvenile rheumatoid arthritis with systemic onset, right knee
M08.262	Juvenile rheumatoid arthritis with systemic onset, left knee
M08.269	Juvenile rheumatoid arthritis with systemic onset, unspecified knee
M08.271	Juvenile rheumatoid arthritis with systemic onset, right ankle and foot
M08.272	Juvenile rheumatoid arthritis with systemic onset, left ankle and foot
M08.279	Juvenile rheumatoid arthritis with systemic onset, unspecified ankle and foot
M08.28	Juvenile rheumatoid arthritis with systemic onset, vertebrae
M08.29	Juvenile rheumatoid arthritis with systemic onset, multiple sites

ICD-10	ICD-10 Description
M08.3	Juvenile rheumatoid polyarthritis (seronegative)
M08.3	Juvenile rheumatoid polyarthritis (seronegative)
M08.4A	Pauciarticular juvenile rheumatoid arthritis, other specified site
M08.411	Pauciarticular juvenile rheumatoid arthritis, right shoulder
M08.412	Pauciarticular juvenile rheumatoid arthritis, left shoulder
M08.419	Pauciarticular juvenile rheumatoid arthritis, unspecified shoulder
M08.421	Pauciarticular juvenile rheumatoid arthritis, right elbow
M08.422	Pauciarticular juvenile rheumatoid arthritis, left elbow
M08.429	Pauciarticular juvenile rheumatoid arthritis, unspecified elbow
M08.431	Pauciarticular juvenile rheumatoid arthritis, right wrist
M08.432	Pauciarticular juvenile rheumatoid arthritis, left wrist
M08.439	Pauciarticular juvenile rheumatoid arthritis, unspecified wrist
M08.441	Pauciarticular juvenile rheumatoid arthritis, right hand
M08.442	Pauciarticular juvenile rheumatoid arthritis, left hand
M08.449	Pauciarticular juvenile rheumatoid arthritis, unspecified hand
M08.451	Pauciarticular juvenile rheumatoid arthritis, right hip
M08.452	Pauciarticular juvenile rheumatoid arthritis, left hip
M08.459	Pauciarticular juvenile rheumatoid arthritis, unspecified hip
M08.461	Pauciarticular juvenile rheumatoid arthritis, right knee
M08.462	Pauciarticular juvenile rheumatoid arthritis, left knee
M08.469	Pauciarticular juvenile rheumatoid arthritis, unspecified knee
M08.471	Pauciarticular juvenile rheumatoid arthritis, right ankle and foot
M08.472	Pauciarticular juvenile rheumatoid arthritis, left ankle and foot
M08.479	Pauciarticular juvenile rheumatoid arthritis, unspecified ankle and foot
M08.48	Pauciarticular juvenile rheumatoid arthritis, vertebrae
M08.80	Other juvenile arthritis, unspecified site
M08.811	Other juvenile arthritis, right shoulder
M08.812	Other juvenile arthritis, left shoulder
M08.819	Other juvenile arthritis, unspecified shoulder
M08.821	Other juvenile arthritis, right elbow
M08.822	Other juvenile arthritis, left elbow
M08.829	Other juvenile arthritis, unspecified elbow

ICD-10	ICD-10 Description
M08.831	Other juvenile arthritis, right wrist
M08.832	Other juvenile arthritis, left wrist
M08.839	Other juvenile arthritis, unspecified wrist
M08.841	Other juvenile arthritis, right hand
M08.842	Other juvenile arthritis, left hand
M08.849	Other juvenile arthritis, unspecified hand
M08.851	Other juvenile arthritis, right hip
M08.852	Other juvenile arthritis, left hip
M08.859	Other juvenile arthritis, unspecified hip
M08.861	Other juvenile arthritis, right knee
M08.862	Other juvenile arthritis, left knee
M08.869	Other juvenile arthritis, unspecified knee
M08.871	Other juvenile arthritis, right ankle and foot
M08.872	Other juvenile arthritis, left ankle and foot
M08.879	Other juvenile arthritis, unspecified ankle and foot
M08.88	Other juvenile arthritis, other specified site
M08.89	Other juvenile arthritis, multiple sites
M08.9A	Juvenile arthritis, unspecified, other specified site
M08.911	Juvenile arthritis, unspecified, right shoulder
M08.912	Juvenile arthritis, unspecified, left shoulder
M08.919	Juvenile arthritis, unspecified, unspecified shoulder
M08.921	Juvenile arthritis, unspecified, right elbow
M08.922	Juvenile arthritis, unspecified, left elbow
M08.929	Juvenile arthritis, unspecified, unspecified elbow
M08.931	Juvenile arthritis, unspecified, right wrist
M08.932	Juvenile arthritis, unspecified, left wrist
M08.939	Juvenile arthritis, unspecified, unspecified wrist
M08.941	Juvenile arthritis, unspecified, right hand
M08.942	Juvenile arthritis, unspecified, left hand
M08.949	Juvenile arthritis, unspecified, unspecified hand
M08.951	Juvenile arthritis, unspecified, right hip
M08.952	Juvenile arthritis, unspecified, left hip

ICD-10	ICD-10 Description
M08.959	Juvenile arthritis, unspecified, unspecified hip
M08.961	Juvenile arthritis, unspecified, right knee
M08.962	Juvenile arthritis, unspecified, left knee
M08.969	Juvenile arthritis, unspecified, unspecified knee
M08.971	Juvenile arthritis, unspecified, right ankle and foot
M08.972	Juvenile arthritis, unspecified, left ankle and foot
M08.979	Juvenile arthritis, unspecified, unspecified ankle and foot
M08.98	Juvenile arthritis, unspecified, vertebrae
M08.99	Juvenile arthritis, unspecified, multiple sites
M60.811	Other myositis, right shoulder
M60.812	Other myositis, left shoulder
M60.819	Other myositis, unspecified shoulder
M60.821	Other myositis, right upper arm
M60.822	Other myositis, left upper arm
M60.829	Other myositis, unspecified upper arm
M60.831	Other myositis, right forearm
M60.832	Other myositis, left forearm
M60.839	Other myositis, unspecified forearm
M60.841	Other myositis, right hand
M60.842	Other myositis, left hand
M60.849	Other myositis, unspecified hand
M60.851	Other myositis, right thigh
M60.852	Other myositis, left thigh
M60.859	Other myositis, unspecified thigh
M60.861	Other myositis, right lower leg
M60.862	Other myositis, left lower leg
M60.869	Other myositis, unspecified lower leg
M60.871	Other myositis, right ankle and foot
M60.872	Other myositis, left ankle and foot
M60.879	Other myositis, unspecified ankle and foot
M60.88	Other myositis, other site
M60.89	Other myositis, multiple sites

ICD-10	ICD-10 Description
T86.09	Other complications of bone marrow transplant
Z94.81	Bone marrow transplant status

Appendix 2 – Centers for Medicare and Medicaid Services (CMS)

The preceding information is intended for non-Medicare coverage determinations. Medicare coverage for outpatient (Part B) drugs is outlined in the Medicare Benefit Policy Manual (Pub. 100-2), Chapter 15, §50 Drugs and Biologicals. In addition, National Coverage Determinations (NCDs) and/or Local Coverage Determinations (LCDs) may exist and compliance with these policies is required where applicable. Local Coverage Articles (LCAs) may also exist for claims payment purposes or to clarify benefit eligibility under Part B for drugs which may be self-administered. The following link may be used to search for NCD, LCD, or LCA documents: <https://www.cms.gov/medicare-coverage-database/search.aspx>. Additional indications, including any preceding information, may be applied at the discretion of the health plan.

Medicare Part B Covered Diagnosis Codes (applicable to existing NCD/LCD/LCA): N/A

Medicare Part B Administrative Contractor (MAC) Jurisdictions		
Jurisdiction	Applicable State/US Territory	Contractor
E (1)	CA, HI, NV, AS, GU, CNMI	Noridian Healthcare Solutions, LLC
F (2 & 3)	AK, WA, OR, ID, ND, SD, MT, WY, UT, AZ	Noridian Healthcare Solutions, LLC
5	KS, NE, IA, MO	Wisconsin Physicians Service Insurance Corp (WPS)
6	MN, WI, IL	National Government Services, Inc. (NGS)
H (4 & 7)	LA, AR, MS, TX, OK, CO, NM	Novitas Solutions, Inc.
8	MI, IN	Wisconsin Physicians Service Insurance Corp (WPS)
N (9)	FL, PR, VI	First Coast Service Options, Inc.
J (10)	TN, GA, AL	Palmetto GBA
M (11)	NC, SC, WV, VA (excluding below)	Palmetto GBA
L (12)	DE, MD, PA, NJ, DC (includes Arlington & Fairfax counties and the city of Alexandria in VA)	Novitas Solutions, Inc.
K (13 & 14)	NY, CT, MA, RI, VT, ME, NH	National Government Services, Inc. (NGS)
15	KY, OH	CGS Administrators, LLC