

## Yescarta® (axicabtagene ciloleucel) (Intravenous)

-E-

Document Number: OHSU HEALTHSERVICES-0454

Date Reviewed: 11/2025

Date of Origin: 05/01/2019

Dates Approved: 05/2019, 12/2019, 12/2020, 12/2021, 12/2022, 12/2023, 12/03/2024, 06/24/2025, 12/02/2025

### I. Length of Authorization

- Initial: Prior authorization validity will be provided initially for one treatment course (1 dose).
- Renewal: Prior authorization validity may NOT be renewed.

### II. Dosing Limits

**Max Units (per dose and over time) [HCPCS Unit]:**

- 1 billable unit (1 infusion up to 200 million autologous anti-CD19 CAR-positive viable T-cells)

### III. Initial Approval Criteria <sup>1</sup>

Submission of supporting clinical documentation (including but not limited to medical records, chart notes, lab results, and confirmatory diagnostics) related to the medical necessity criteria is REQUIRED on all requests for authorizations. Records will be reviewed at the time of submission as part of the evaluation of this request. Please provide documentation related to diagnosis, step therapy, and clinical markers (i.e., genetic, and mutational testing) supporting initiation when applicable. Please provide documentation via direct upload through the PA web portal or by fax. Failure to submit the medical records may result in the denial of the request due to inability to establish medical necessity in accordance with policy guidelines.

Prior authorization validity is provided in the following conditions:

- Patient is at least 18 years of age (unless otherwise specified); **AND**
- Patient does not have a clinically significant active systemic infection or inflammatory disorder; **AND**
- Patient has not received live vaccines within 6 weeks prior to the start of lymphodepleting chemotherapy, and will not receive live vaccines during axicabtagene ciloleucel treatment, and until immune recovery following treatment; **AND**

- Patient has been screened for hepatitis B virus (HBV), hepatitis C virus (HCV), and human immunodeficiency virus (HIV) in accordance with clinical guidelines prior to collection of cells (leukapheresis); **AND**
- Prophylaxis for infection will be followed according to local guidelines; **AND**
- Patient has not received prior chimeric antigen receptor (CAR)-T cell therapy; **AND**
- Patient has not received other anti-CD19 therapy (e.g., blinatumomab, tafasitamab, loncastuximab tesirine, etc.) OR patient previously received other anti-CD19 therapy and re-biopsy indicates CD-19 positive disease; **AND**
- Used as a single agent (not applicable to lymphodepleting or additional chemotherapy while awaiting manufacture); **AND**
- Patient has an Eastern Cooperative Oncology Group (ECOG) performance status of 0-1; **AND**
- Patient does not have primary central nervous system lymphoma; **AND**

**B-Cell Lymphomas † ‡ Φ<sup>1,7,12,13</sup>**

- Patient has histologic transformation of follicular lymphoma (indolent lymphoma) to diffuse large b-cell lymphoma (DLBCL); **AND**
  - Prior therapy included an anthracycline and an anti-CD20 monoclonal antibody, unless contraindicated; **AND**
    - Disease is refractory to first-line chemoimmunotherapy; **OR**
    - Used for treatment of disease that has relapsed within 12 months of first-line chemoimmunotherapy; **OR**
    - Patient has relapsed or refractory disease after two (2) or more prior lines of chemoimmunotherapy for indolent disease prior to histologic transformation; **OR**
- Patient has histologic transformation of marginal zone lymphoma (indolent lymphoma) to DLBCL OR Richter's transformation of Chronic Lymphocytic Leukemia (CLL) to DLBCL; **AND**
  - Patient has received two (2) or more prior lines of chemoimmunotherapy for indolent disease prior to histologic transformation which must have included an anthracycline and an anti-CD20 monoclonal antibody, unless contraindicated; **OR**
- Patient has HIV-related large B-cell lymphoma (i.e., HIV-related DLBCL or HHV8-positive DLBCL, not otherwise specified), DLBCL, primary mediastinal large B-cell lymphoma (PMBCL), high grade B-cell lymphoma, or monomorphic post-transplant lymphoproliferative disorder (B-cell type); **AND**
  - Prior therapy included an anthracycline and an anti-CD20 monoclonal antibody, unless contraindicated; **AND**
    - Used as additional therapy for relapsed disease >12 months after completion of first-line therapy and partial response following second-line therapy; **OR**

- Used as additional therapy for patients with primary refractory disease (partial response, no response, or progression) or relapsed disease <12 months after completion of first-line therapy; **OR**
  - Used for treatment of disease that is in second or greater relapse in patients with partial response, relapse, or progressive disease following therapy for relapsed or refractory disease; **OR**
- Patient has nodal marginal zone lymphoma (MZL) or follicular lymphoma (grade 1, 2 or 3A); **AND**
  - Used as third-line and subsequent therapy for partial response, no response, relapsed, or progressive disease after prior therapy including an anti-CD20 monoclonal antibody and an alkylating agent, unless contraindicated; **OR**
- Patient has extranodal marginal zone lymphoma (EMZL) of the stomach or of nongastric sites (noncutaneous); **AND**
  - Used as third-line and subsequent therapy for relapsed, refractory, or progressive disease after prior therapy including an anti-CD20 monoclonal antibody and an alkylating agent, unless contraindicated; **OR**
- Patient has splenic marginal zone lymphoma (MZL); **AND**
  - Used as third-line and subsequent therapy for disease recurrence after prior therapy including an anti-CD20 monoclonal antibody and an alkylating agent, unless contraindicated

**Pediatric Aggressive Mature B-Cell Lymphomas ‡<sup>2,7,11,15</sup>**

- Patient is ≤ 18 years of age\*; **AND**
- Patient has primary mediastinal large B-Cell lymphoma; **AND**
- Used as consolidation or additional therapy in patients with a partial response after therapy for relapsed or refractory disease; **AND**
- Patient previously received ≥ 2 prior chemoimmunotherapy regimens which included an anthracycline and an anti-CD20 monoclonal antibody, unless contraindicated

\* Pediatric Aggressive Mature B-Cell Lymphoma may be applicable to adolescent and young adult (AYA) patients older than 18 years of age and less than 39 years of age, who are treated in the pediatric oncology setting.

**Preferred therapies and recommendations are determined by review of clinical evidence. NCCN category of recommendation is taken into account as a component of this review. Regimens deemed equally efficacious (i.e., those having the same NCCN categorization) are considered to be therapeutically equivalent.**

**Enhanced Oncology Value (EOV) Program – Redacted indications**

Uses not listed above have inadequate data to support efficacy and are excluded from prior authorization validity.

Other treatment options including, but are not limited to, the following may be appropriate: radiation therapy, surgery, traditional chemotherapy (e.g., platinum, taxane), compassionate use/expanded access programs, clinical trials, supportive care, integrative and complementary therapies.

† FDA Approved Indication(s); ‡ Compendia Recommended Indication(s); ◊ Orphan Drug

#### IV. Renewal Criteria <sup>1</sup>

Duration of authorization has not been exceeded (*refer to Section I*)

#### V. Dosage/Administration <sup>1</sup>

Indication	Dose
B-Cell Lymphomas	<p><b><u>Lymphodepleting chemotherapy:</u></b></p> <ul style="list-style-type: none"> <li>Administer cyclophosphamide 500 mg/m<sup>2</sup> and fludarabine 30 mg/m<sup>2</sup> intravenously on the fifth, fourth, and third day before infusion of Yescarta</li> </ul> <p><b><u>Yescarta infusion:</u></b></p> <ul style="list-style-type: none"> <li>Each single infusion bag of Yescarta contains a suspension of chimeric antigen receptor (CAR)-positive T cells in approximately 68 mL. The target dose is 2 × 10<sup>6</sup> CAR-positive viable T cells per kg body weight, with a maximum of 2 × 10<sup>8</sup> CAR-positive viable T cells (for patients 100 kg and above).</li> </ul>
<p><b>For autologous use only. For intravenous use only.</b></p> <ul style="list-style-type: none"> <li>Yescarta is prepared from the patient’s peripheral blood mononuclear cells, which are obtained via a standard leukapheresis procedure</li> <li>One treatment course consists of lymphodepleting chemotherapy followed by a single infusion of Yescarta.</li> <li>Confirm Yescarta availability prior to starting the lymphodepleting regimen.</li> <li>Confirm the patient’s identity matches the patient identifiers on the Yescarta cassette.</li> </ul>	
<p><b><u>Premedication:</u></b></p> <ul style="list-style-type: none"> <li>Premedicate with 650 mg acetaminophen and 12.5 mg diphenhydramine 1 hour prior to infusion. Only consider the use of prophylactic systemic corticosteroids in patients after weighing the potential benefits and risks.</li> </ul>	
<p><b><u>Monitoring after infusion:</u></b></p> <ul style="list-style-type: none"> <li>Monitor patients at least daily for 7 days following infusion for signs and symptoms of cytokine release syndrome (CRS) and neurologic toxicities.</li> <li>Instruct patients to remain within proximity of a healthcare facility for at least 2 weeks following infusion.</li> <li>Instruct patients to refrain from driving for at least 2 weeks following infusion.</li> </ul>	
<ul style="list-style-type: none"> <li>Store infusion bag in the vapor phase of liquid nitrogen (less than or equal to minus 150°C). Thaw prior to infusion.</li> <li>In case of manufacturing failure, a second manufacturing may be attempted.</li> <li>Additional chemotherapy (not the lymphodepletion) may be necessary while the patient awaits the product.</li> <li>Ensure that 2 doses of tocilizumab and emergency equipment are available prior to infusion and during the recovery period.</li> </ul>	

• Yescarta contains human blood cells that are genetically modified with replication incompetent retroviral vector. Follow universal precautions and local biosafety guidelines for handling and disposal.

## VI. Billing Code/Availability Information

### HCPCS Code:

- Q2041 – Axicabtagene Ciloleucel, up to 200 million autologous anti-cd19 car-positive viable t cells, including leukapheresis and dose preparation procedures, per therapeutic dose

### NDC:

- Yescarta suspension for intravenous infusion; 1 infusion bag (~68 mL): 71287-0119-xx

## VII. References (STANDARD)

1. Yescarta [package insert]. Santa Monica, CA; Kite Pharma, Inc., October 2025. Accessed November 2025.
2. Locke FL, Neelapu SS, Bartlett NL, et al. Phase 1 Results of ZUMA-1: A Multicenter Study of KTE-C19 Anti-CD19 CAR T Cell Therapy in Refractory Aggressive Lymphoma. *Mol Ther*. 2017 Jan 4;25(1):285-295.
3. Mejstrikova E, Hrusak O, Borowitz MJ, et al. CD19-negative relapse of pediatric B-cell precursor acute lymphoblastic leukemia following blinatumomab treatment. *Blood Cancer J*. 2017; 659. DOI 10.1038/s41408-017-0023-x
4. Ruella M, Maus MV. Catch me if you can: Leukemia Escape after CD19-Directed T Cell Immunotherapies. *Computational and Structural Biotechnology Journal* 14 (2016) 357–362.
5. Braig F, Brandt A, Goebeler M, et al. Resistance to anti-CD19/CD3 BiTE in acute lymphoblastic leukemia may be mediated by disrupted CD19 membrane trafficking. *Blood*; 129:1, 2017 Jan.
6. Majzner RG, Mackall CL. Tumor Antigen Escape from CAR T-cell Therapy. *Cancer Discov* 2018;8:1219-1226.
7. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) axicabtagene ciloleucel. National Comprehensive Cancer Network, 2025. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed November 2025.
8. Locke FL, Ghobadi A, Jacobson CA, et al. Long-term safety and activity of axicabtagene ciloleucel in refractory large B-cell lymphoma (ZUMA-1): a single-arm, multicentre, phase 1-2 trial. *Lancet Oncol*. 2019;20(1):31-42. doi:10.1016/S1470-2045(18)30864-7.
9. Neelapu SS, Jacobson CA, Oluwole OO, et al. Outcomes of older patients in ZUMA-1, a pivotal study of axicabtagene ciloleucel in refractory large B-cell lymphoma. *Blood*. 2020;135(23):2106-2109. doi:10.1182/blood.2019004162.

10. Neelapu SS, Locke FL, Bartlett NL, et al. Axicabtagene ciloleucel CAR T-cell therapy in refractory large B-cell lymphoma. *N Engl J Med*. 2017;377(26):2531-2544. doi: 10.1056/NEJMoa1707447.
11. Jacobson CA, Chavez JC, Sehgal AR, et al. Interim analysis of ZUMA-5: A phase II study of axicabtagene ciloleucel (axi-cel) in patients (pts) with relapsed/refractory indolent non-Hodgkin lymphoma (R/R iNHL). *Journal of Clinical Oncology* 2020 38:15\_suppl, 8008-8008.
12. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Chronic Lymphocytic Leukemia/Small Lymphocytic Lymphoma Version 3.2025. National Comprehensive Cancer Network, 2025. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed November 2025.
13. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) B-Cell Lymphomas, Version 3.2025. National Comprehensive Cancer Network, 2025. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the NCCN Guidelines, go online to NCCN.org. Accessed November 2025.
14. Locke FL, Miklos DB, Jacobson CA, et al; All ZUMA-7 Investigators and Contributing Kite Members. Axicabtagene Ciloleucel as Second-Line Therapy for Large B-Cell Lymphoma. *N Engl J Med*. 2022 Feb 17;386(7):640-654. doi: 10.1056/NEJMoa2116133.
15. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Pediatric Aggressive Mature B-Cell Lymphomas Version 2.2025. National Comprehensive Cancer Network, 2025. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed November 2025.

## VIII. References (ENHANCED)

- 1e. Schuster S, Bishop M, Tam C, et al. Tisagenlecleucel in Adult Relapsed or Refractory Diffuse Large B-Cell Lymphoma. *N Engl J Med* 2019; 380:45-56.
- 2e. Abramson JS, Palomba ML, Gordon LI, et al. Lisocabtagene maraleucel for patients with relapsed or refractory large B-cell lymphomas (TRANSCEND NHL 001): a multicentre seamless design study. *Lancet*. 2020 Sep 19;396(10254):839-852. doi: 10.1016/S0140-6736(20)31366-0. Epub 2020 Sep 1.
- 3e. Caimi PF, Ai WZ, Alderuccio JP, et al. Loncastuximab tesirine in relapsed or refractory diffuse large B-cell lymphoma (LOTIS-2): a multicentre, open-label, single-arm, phase 2 trial. *Lancet Oncol* 2021;22:790-800.

- 4e. Kalakonda N, Maerevoet M, Cavallo F, et al. Selinexor in patients with relapsed or refractory diffuse large B-cell lymphoma (SADAL): a single-arm, multinational, multicentre, open-label, phase 2 trial. *Lancet Haematol*. 2020 Jul;7(7):e511-e522. doi: 10.1016/S2352-3026(20)30120-4.
- 5e. Salles G, Duell J, González Barca E, et al. Tafasitamab plus lenalidomide in relapsed or refractory diffuse large B-cell lymphoma (L-MIND): a multicentre, prospective, single-arm, phase 2 study. *Lancet Oncol*. 2020 Jul;21(7):978-988. doi: 10.1016/S1470-2045(20)30225-4.
- 6e. Dreyling, M., Santoro, A., Mollica, L., et al. (2017) COPANLISIB IN PATIENTS WITH RELAPSED OR REFRACTORY INDOLENT B-CELL LYMPHOMA (CHRONOS-1). *Hematological Oncology*, 35(S2): 119–120. doi: 10.1002/hon.2437\_107.
- 7e. Jacobson CA, Chavez JC, Sehgal AR, et al. Axicabtagene ciloleucel in relapsed or refractory indolent non-Hodgkin lymphoma (ZUMA-5): a single-arm, multicentre, phase 2 trial. *Lancet Oncol*. 2022 Jan;23(1):91-103. doi: 10.1016/S1470-2045(21)00591-X. Epub 2021 Dec 8.
- 8e. Mounier N, El Gnaoui T, Tilly H, et al. Rituximab plus gemcitabine and oxaliplatin in patients with refractory/relapsed diffuse large B-cell lymphoma who are not candidates for high-dose therapy. A phase II Lymphoma Study Association trial. *Haematologica*. 2013;98(11):1726-1731. doi:10.3324/haematol.2013.090597.
- 9e. Morschhauser F, Tilly H, Chaidos A, et al. Tazemetostat for patients with relapsed or refractory follicular lymphoma: an open-label, single-arm, multicentre, phase 2 trial. *Lancet Oncol* 2020;21:1433-1442.
- 10e. Fowler NH, Dickinson M, Dreyling M, et al. Tisagenlecleucel in adult relapsed or refractory follicular lymphoma: the phase 2 ELARA trial. *Nat Med*. 2022 Feb;28(2):325-332. doi: 10.1038/s41591-021-01622-0.
- 11e. Thieblemont C, Phillips T, Ghesquieres H, et al. Epcoritamab, a Novel, Subcutaneous CD3xCD20 Bispecific T-Cell-Engaging Antibody, in Relapsed or Refractory Large B-Cell Lymphoma: Dose Expansion in a Phase I/II Trial. *J Clin Oncol*. 2023 Apr 20;41(12):2238-2247.
- 12e. Dickinson MJ, Carlo-Stella C, Morschhauser F, et al. Glofitamab for Relapsed or Refractory Diffuse Large B-Cell Lymphoma. *N Engl J Med* 2022;387:22220-2231.
- 13e. Bartlett NL, Sehn LH, Matasar MJ, et al. Mosunetuzumab Monotherapy Demonstrates Durable Efficacy with a Manageable Safety Profile in Patients with Relapsed/Refractory Follicular Lymphoma Who Received  $\geq 2$  Prior Therapies: Updated Results from a Pivotal Phase II Study [abstract]. *Blood* 2022;140:1467-1470.
- 14e. Barta SK, Noy A, Pasquini MC, et al. Observational Cohort Study of People Living with HIV (PWH) Treated with CD19-Directed CAR T Cell Therapy for B-Cell Lymphoid Malignancies - Interim Results of AIDS Malignancy Consortium (AMC) Study AMC-113. *Blood* 2022;140:1847-1848.
- 15e. Epstein-Peterson ZD, Lee JY, Barta SK, et al. Trial-in-Progress: AMC 112: Axicabtagene Ciloleucel in Relapsed or Refractory HIV-Associated Aggressive B-Cell Non-Hodgkin Lymphoma (NCT05077527). *Blood* 2022;140:12788-12789.

16e. Kittai AS, Bond D, Huang Y, et al. Anti-CD19 chimeric antigen receptor T-cell therapy for Richter transformation: An international, multicenter, retrospective study. J Clin Oncol 2024;42:2071-2079.

17e. Prime Therapeutics Management. Yescarta Clinical Literature Review Analysis. Last updated November 2025. Accessed November 2025.

## Appendix A – Non-Quantitative Treatment Limitations (NQTL) Factor Checklist

Non-quantitative treatment limitations (NQTLs) refer to the methods, guidelines, standards of evidence, or other conditions that can restrict how long or to what extent benefits are provided under a health plan. These may include things like utilization review or prior authorization. The utilization management NQTL applies comparably, and not more stringently, to mental health/substance use disorder (MH/SUD) Medical Benefit Prescription Drugs and medical/surgical (M/S) Medical Benefit Prescription Drugs. The table below lists the factors that were considered in designing and applying prior authorization to this drug/drug group, and a summary of the conclusions that Prime’s assessment led to for each.

Factor	Conclusion
Indication	Yes: Consider for PA
Safety and efficacy	Yes: Consider for PA
Potential for misuse/abuse	No: PA not a priority
Cost of drug	Yes: Consider for PA

## Appendix 1 – Covered Diagnosis Codes

ICD-10	ICD-10 Description
C82.00	Follicular lymphoma grade I, unspecified site
C82.01	Follicular lymphoma grade I, lymph nodes of head, face and neck
C82.02	Follicular lymphoma, grade I, intrathoracic lymph nodes
C82.03	Follicular lymphoma grade I, intra-abdominal lymph nodes
C82.04	Follicular lymphoma grade I, lymph nodes of axilla and upper limb
C82.05	Follicular lymphoma grade I, lymph nodes of inguinal regional and lower limb
C82.06	Follicular lymphoma grade I, intrapelvic lymph nodes
C82.07	Follicular lymphoma grade I, spleen
C82.08	Follicular lymphoma grade I, lymph nodes of multiple sites
C82.09	Follicular lymphoma grade I, extranodal and solid organ sites
C82.10	Follicular lymphoma grade II, unspecified site
C82.11	Follicular lymphoma grade II, lymph nodes of head, face and neck
C82.12	Follicular lymphoma, grade II, intrathoracic lymph nodes
C82.13	Follicular lymphoma grade II, intra-abdominal lymph nodes

C82.14	Follicular lymphoma grade II, lymph nodes of axilla and upper limb
C82.15	Follicular lymphoma grade II, lymph nodes of inguinal region and lower limb
C82.16	Follicular lymphoma grade II, intrapelvic lymph nodes
C82.17	Follicular lymphoma grade II, spleen
C82.18	Follicular lymphoma grade II, lymph nodes of multiple sites
C82.19	Follicular lymphoma grade II, extranodal and solid organ sites
C82.20	Follicular lymphoma grade III, unspecified, unspecified site
C82.21	Follicular lymphoma grade III, unspecified, lymph nodes of head, face and neck
C82.22	Follicular lymphoma, grade III, unspecified, intrathoracic lymph nodes
C82.23	Follicular lymphoma grade III, unspecified, intra-abdominal lymph nodes
C82.24	Follicular lymphoma grade III, unspecified, lymph nodes of axilla and upper limb
C82.25	Follicular lymphoma grade III, unspecified, lymph nodes of inguinal region and lower limb
C82.26	Follicular lymphoma grade III, unspecified, intrapelvic lymph nodes
C82.27	Follicular lymphoma grade III, unspecified, spleen
C82.28	Follicular lymphoma grade III, unspecified, lymph nodes of multiple sites
C82.29	Follicular lymphoma grade III, unspecified, extranodal and solid organ sites
C82.30	Follicular lymphoma grade IIIa, unspecified site
C82.31	Follicular lymphoma grade IIIa, lymph nodes of head, face and neck
C82.32	Follicular lymphoma, grade IIIa, intrathoracic lymph nodes
C82.33	Follicular lymphoma grade IIIa, intra-abdominal lymph nodes
C82.34	Follicular lymphoma grade IIIa, lymph nodes of axilla and upper limb
C82.35	Follicular lymphoma grade IIIa, lymph nodes of inguinal region and lower limb
C82.36	Follicular lymphoma grade IIIa, intrapelvic lymph nodes
C82.37	Follicular lymphoma grade IIIa, spleen
C82.38	Follicular lymphoma grade IIIa, lymph nodes of multiple sites
C82.39	Follicular lymphoma grade IIIa, extranodal and solid organ sites
C82.40	Follicular lymphoma grade IIIb, unspecified site
C82.41	Follicular lymphoma grade IIIb, lymph nodes of head, face, and neck
C82.42	Follicular lymphoma grade IIIb, intrathoracic lymph nodes
C82.43	Follicular lymphoma grade IIIb, intra-abdominal lymph nodes
C82.44	Follicular lymphoma grade IIIb, lymph nodes of axilla and upper limb
C82.45	Follicular lymphoma grade IIIb, lymph nodes of inguinal region and lower limb
C82.46	Follicular lymphoma grade IIIb, intrapelvic lymph nodes

C82.47	Follicular lymphoma grade IIIb, spleen
C82.48	Follicular lymphoma grade IIIb, lymph nodes of multiple sites
C82.49	Follicular lymphoma grade IIIb, extranodal and solid organ sites
C82.50	Diffuse follicle center lymphoma, unspecified site
C82.51	Diffuse follicle center lymphoma, lymph nodes of head, face and neck
C82.52	Diffuse follicle center lymphoma, intrathoracic lymph nodes
C82.53	Diffuse follicle center lymphoma, intra-abdominal lymph nodes
C82.54	Diffuse follicle center lymphoma, lymph nodes of axilla and upper limb
C82.55	Diffuse follicle center lymphoma, lymph nodes of inguinal region and lower limb
C82.56	Diffuse follicle center lymphoma, intrapelvic lymph nodes
C82.57	Diffuse follicle center lymphoma, spleen
C82.58	Diffuse follicle center lymphoma, lymph nodes of multiple sites
C82.59	Diffuse follicle center lymphoma, extranodal and solid organ sites
C82.60	Cutaneous follicle center lymphoma, unspecified site
C82.61	Cutaneous follicle center lymphoma, lymph nodes of head, face and neck
C82.62	Cutaneous follicle center lymphoma, intrathoracic lymph nodes
C82.63	Cutaneous follicle center lymphoma, intra-abdominal lymph nodes
C82.64	Cutaneous follicle center lymphoma, lymph nodes of axilla and upper limb
C82.65	Cutaneous follicle center lymphoma, lymph nodes of inguinal region and lower limb
C82.66	Cutaneous follicle center lymphoma, intrapelvic lymph nodes
C82.67	Cutaneous follicle center lymphoma, spleen
C82.68	Cutaneous follicle center lymphoma, lymph nodes of multiple sites
C82.69	Cutaneous follicle center lymphoma, extranodal and solid organ sites
C82.80	Other types of follicular lymphoma, unspecified site
C82.81	Other types of follicular lymphoma, lymph nodes of head, face and neck
C82.82	Other types of follicular lymphoma, intrathoracic lymph nodes
C82.83	Other types of follicular lymphoma, intra-abdominal lymph nodes
C82.84	Other types of follicular lymphoma, lymph nodes of axilla and upper limb
C82.85	Other types of follicular lymphoma, lymph nodes of inguinal region and lower limb
C82.86	Other types of follicular lymphoma, intrapelvic lymph nodes
C82.87	Other types of follicular lymphoma, spleen
C82.88	Other types of follicular lymphoma, lymph nodes of multiple sites
C82.89	Other types of follicular lymphoma, extranodal and solid organ sites

C82.90	Follicular lymphoma, unspecified, unspecified site
C82.91	Follicular lymphoma, unspecified, lymph nodes of head, face and neck
C82.92	Follicular lymphoma, unspecified, intrathoracic lymph nodes
C82.93	Follicular lymphoma, unspecified, intra-abdominal lymph nodes
C82.94	Follicular lymphoma, unspecified, lymph nodes of axilla and upper limb
C82.95	Follicular lymphoma, unspecified lymph nodes of inguinal region and lower limb
C82.96	Follicular lymphoma, unspecified, intrapelvic lymph nodes
C82.97	Follicular lymphoma, unspecified, spleen
C82.98	Follicular lymphoma, unspecified, lymph nodes of multiple sites
C82.99	Follicular lymphoma, unspecified, extranodal and solid organ sites
C83.00	Small cell B-cell lymphoma, unspecified site
C83.01	Small cell B-cell lymphoma, lymph nodes of head, face and neck
C83.02	Small cell B-cell lymphoma, intrathoracic lymph nodes
C83.03	small cell B-cell lymphoma, intra-abdominal lymph nodes
C83.04	Small cell B-cell lymphoma, lymph nodes of axilla and upper limb
C83.05	Small cell B-cell lymphoma, lymph nodes of inguinal region and lower limb
C83.06	Small cell B-cell lymphoma, intrapelvic lymph nodes
C83.07	Small cell B-cell lymphoma, spleen
C83.08	Small cell B-cell lymphoma, lymph nodes of multiple sites
C83.09	Small cell B-cell lymphoma, extranodal and solid organ sites
C83.30	Diffuse large B-cell lymphoma unspecified site
C83.31	Diffuse large B-cell lymphoma, lymph nodes of head, face, and neck
C83.32	Diffuse large B-cell lymphoma intrathoracic lymph nodes
C83.33	Diffuse large B-cell lymphoma intra-abdominal lymph nodes
C83.34	Diffuse large B-cell lymphoma lymph nodes of axilla and upper limb
C83.35	Diffuse large B-cell lymphoma, lymph nodes of inguinal region and lower limb
C83.36	Diffuse large B-cell lymphoma intrapelvic lymph nodes
C83.37	Diffuse large B-cell lymphoma, spleen
C83.38	Diffuse large B-cell lymphoma lymph nodes of multiple sites
C83.39	Diffuse large B-cell lymphoma extranodal and solid organ sites
C83.90	Non-follicular (diffuse) lymphoma, unspecified site
C83.91	Non-follicular (diffuse) lymphoma, unspecified lymph nodes of head, face, and neck
C83.92	Non-follicular (diffuse) lymphoma, unspecified intrathoracic lymph nodes

C83.93	Non-follicular (diffuse) lymphoma, unspecified intra-abdominal lymph nodes
C83.94	Non-follicular (diffuse) lymphoma, unspecified lymph nodes of axilla and upper limb
C83.95	Non-follicular (diffuse) lymphoma, unspecified lymph nodes of inguinal region and lower limb
C83.96	Non-follicular (diffuse) lymphoma, unspecified intrapelvic lymph nodes
C83.97	Non-follicular (diffuse) lymphoma, unspecified spleen
C83.98	Non-follicular (diffuse) lymphoma, unspecified lymph nodes of multiple sites
C83.99	Non-follicular (diffuse) lymphoma, unspecified extranodal and solid organ sites
C85.10	Unspecified B-cell lymphoma, unspecified site
C85.11	Unspecified B-cell lymphoma, lymph nodes of head, face, and neck
C85.12	Unspecified B-cell lymphoma, intrathoracic lymph nodes
C85.13	Unspecified B-cell lymphoma, intra-abdominal lymph nodes
C85.14	Unspecified B-cell lymphoma, lymph nodes of axilla and upper limb
C85.15	Unspecified B-cell lymphoma, lymph nodes of inguinal region and lower limb
C85.16	Unspecified B-cell lymphoma, intrapelvic lymph nodes
C85.17	Unspecified B-cell lymphoma, spleen
C85.18	Unspecified B-cell lymphoma, lymph nodes of multiple sites
C85.19	Unspecified B-cell lymphoma, extranodal and solid organ sites
C85.20	Mediastinal (thymic) large B-cell lymphoma, unspecified site
C85.21	Mediastinal (thymic) large B-cell lymphoma, lymph nodes of head, face and neck
C85.22	Mediastinal (thymic) large B-cell lymphoma, intrathoracic lymph nodes
C85.23	Mediastinal (thymic) large B-cell lymphoma, intra-abdominal lymph nodes
C85.24	Mediastinal (thymic) large B-cell lymphoma, lymph nodes of axilla and upper limb
C85.25	Mediastinal (thymic) large B-cell lymphoma, lymph nodes of inguinal region and lower limb
C85.26	Mediastinal (thymic) large B-cell lymphoma, intrapelvic lymph nodes
C85.27	Mediastinal (thymic) large B-cell lymphoma, spleen
C85.28	Mediastinal (thymic) large B-cell lymphoma, lymph nodes of multiple sites
C85.29	Mediastinal (thymic) large B-cell lymphoma, extranodal and solid organ sites
C85.80	Other specified types of non-Hodgkin lymphoma, unspecified site
C85.81	Other specified types of non-Hodgkin lymphoma, lymph nodes of head, face and neck
C85.82	Other specified types of non-Hodgkin lymphoma, intrathoracic lymph nodes
C85.83	Other specified types of non-Hodgkin lymphoma, intra-abdominal lymph nodes
C85.84	Other specified types of non-Hodgkin lymphoma, lymph nodes of axilla and upper limb
C85.85	Other specified types of non-Hodgkin lymphoma, lymph nodes of inguinal region of lower limb

C85.86	Other specified types of non-Hodgkin lymphoma, intrapelvic lymph nodes
C85.87	Other specified types of non-Hodgkin lymphoma, spleen
C85.88	Other specified types of non-Hodgkin lymphoma, lymph nodes of multiple sites
C85.89	Other specified types of non-Hodgkin lymphoma, extranodal and solid organ sites
C85.87	Other specified types of non-Hodgkin lymphoma, spleen
C91.10	Chronic lymphocytic leukemia of B-cell type not having achieved remission
C91.12	Chronic lymphocytic leukemia of B-cell type in relapse
D47.21	Post-transplant lymphoproliferative disorder (PTLD)

## Appendix 2 – Centers for Medicare and Medicaid Services (CMS)

The preceding information is intended for non-Medicare coverage determinations. Medicare coverage for outpatient (Part B) drugs is outlined in the Medicare Benefit Policy Manual (Pub. 100-2), Chapter 15, §50 Drugs and Biologicals. In addition, National Coverage Determinations (NCDs) and/or Local Coverage Determinations (LCDs) may exist and compliance with these policies is required where applicable. Local Coverage Articles (LCAs) may also exist for claims payment purposes or to clarify benefit eligibility under Part B for drugs which may be self-administered. The following link may be used to search for NCD, LCD, or LCA documents: <https://www.cms.gov/medicare-coverage-database/search.aspx>. Additional indications, including any preceding information, may be applied at the discretion of the health plan.

Medicare Part B Covered Diagnosis Codes (applicable to existing NCD/LCA/LCD): N/A

Medicare Part B Administrative Contractor (MAC) Jurisdictions		
Jurisdiction	Applicable State/US Territory	Contractor
E (1)	CA, HI, NV, AS, GU, CNMI	Noridian Healthcare Solutions, LLC
F (2 & 3)	AK, WA, OR, ID, ND, SD, MT, WY, UT, AZ	Noridian Healthcare Solutions, LLC
5	KS, NE, IA, MO	Wisconsin Physicians Service Insurance Corp (WPS)
6	MN, WI, IL	National Government Services, Inc. (NGS)
H (4 & 7)	LA, AR, MS, TX, OK, CO, NM	Novitas Solutions, Inc.
8	MI, IN	Wisconsin Physicians Service Insurance Corp (WPS)
N (9)	FL, PR, VI	First Coast Service Options, Inc.
J (10)	TN, GA, AL	Palmetto GBA
M (11)	NC, SC, WV, VA (excluding below)	Palmetto GBA
L (12)	DE, MD, PA, NJ, DC (includes Arlington & Fairfax counties and the city of Alexandria in VA)	Novitas Solutions, Inc.
K (13 & 14)	NY, CT, MA, RI, VT, ME, NH	National Government Services, Inc. (NGS)
15	KY, OH	CGS Administrators, LLC