



Oregon Health & Science University
Hospital and Clinics Provider's Orders

PO7071



ADULT AMBULATORY INFUSION ORDER
**Agalsidase Beta (FABRAZYME)
Infusion**

Page 1 of 2

ACCOUNT NO.
MED. REC. NO.
NAME
BIRTHDATE

Patient Identification

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.

Weight: _____ kg Height: _____ cm

Allergies: _____

Diagnosis Code: _____

Treatment Start Date: _____ Patient to follow up with provider on date: _____

****This plan will expire after 365 days at which time a new order will need to be placed****

GUIDELINES FOR ORDERING

1. **Send FACE SHEET and H&P or most recent chart note.**
2. Indicated for use in patients with Fabry disease. Reduces globotriaosylceramide (GL-3) deposition in capillary endothelium of the kidney and certain other cell types
3. Please encourage patients to enroll in the Fabry registry by visiting www.fabryregistry.com or calling 1-800-745-4447.
4. Patients with advanced Fabry disease may have compromised cardiac function which may predispose them to a higher risk of severe complications from infusion reactions.

NURSING ORDERS:

1. Follow facility policies and/or protocols for vascular access maintenance with appropriate flush solution, dec clotting (alteplase), and/or dressing changes.
2. VITAL SIGNS – Monitor and record vital signs, tolerance, and presence of infusion-related reactions prior to infusion, with every rate increase, then hourly until infusion is complete. Observe patient for 60 minutes following infusion (unless prescriber indicates this is not necessary).
3. Initial rate should not exceed 30 mL/hour. If no previous infusion reactions, rate me by increased in increments of 10 mL/hour with each subsequent infusion. Minimum total infusion time 1.5 hours (MAX RATE = total volume/1.5). If previous infusion reaction contact provider for guidance.
4. Reschedule patient for next weekly infusion.

PRE-MEDICATIONS: (Administer 30 minutes prior to infusion)

Note to provider: Please select which medications below, if any, you would like the patient to receive prior to treatment by checking the appropriate box(s)

- acetaminophen (TYLENOL) tablet, 650 mg, oral, ONCE, every visit
 - diphenhydrAMINE (BENADRYL) capsule, 50 mg, oral, ONCE, every visit
 - loratadine (CLARITIN) tablet, 10 mg, oral, ONCE, every visit
- (Choose as alternative to diphenhydrAMINE if needed)**

MEDICATIONS:

Agalsidase beta (FABRAZYME) 1 mg/kg = _____ mg in sodium chloride 0.9% IV infusion, ONCE, every 2 weeks x _____ doses (Pharmacist will round dose to nearest 5 mg vial and modify during order verification)

Administer using an in-line low protein binding 0.2 micron filter.



Oregon Health & Science University
Hospital and Clinics Provider's Orders

ADULT AMBULATORY INFUSION ORDER
**Agalsidase Beta (FABRAZYME)
Infusion**

Page 2 of 2

ACCOUNT NO.
MED. REC. NO.
NAME
BIRTHDATE

Patient Identification

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.

HYPERSENSITIVITY MEDICATIONS:

1. NURSING COMMUNICATION – If hypersensitivity or infusion reactions develop, temporarily hold the infusion and notify provider immediately. Administer emergency medications per the Treatment Algorithm for Acute Infusion Reaction (OHSU HC-PAT-133-GUD, HMC C-132). Refer to algorithm for symptom monitoring and continuously assess as grade of severity may progress.
2. diphenhydrAMINE (BENADRYL) injection, 25-50 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
3. EPINEPHrine HCl (ADRENALIN) injection, 0.3 mg, intramuscular, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
4. hydrocortisone sodium succinate (SOLU-CORTEF) injection, 100 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
5. famotidine (PEPCID) injection, 20 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction

By signing below, I represent the following:

I am responsible for the care of the patient (*who is identified at the top of this form*);

I hold an active, unrestricted license to practice medicine in: Oregon _____ (*check box that corresponds with state where you provide care to patient and where you are currently licensed. Specify state if not Oregon*);

My physician license Number is # _____ (MUST BE COMPLETED TO BE A VALID PRESCRIPTION); and I am acting within my scope of practice and authorized by law to order Infusion of the medication described above for the patient identified on this form.

Provider signature: _____	Date/Time: _____
Printed Name: _____	Phone: _____ Fax: _____

INFUSION REFERRAL TEAM Phone (providers only) (971) 262-9645 Fax completed orders to (503) 346-8058 <i>Infusion orders located at:</i> www.ohsuknight.com/infusionorders	<input checked="" type="checkbox"/> Please indicate the patient's preferred clinic location below	
	<input type="checkbox"/> BEAVERTON OHSU Knight Cancer Institute	15700 SW Greystone Court Beaverton OR 97006
	<input type="checkbox"/> NW PORTLAND Legacy Good Samaritan campus	Medical Office Building 3 – Suite 150 1130 NW 22nd Ave, Portland OR 97210
	<input type="checkbox"/> GRESHAM Legacy Mount Hood campus	Medical Office Building 3 – Suite 140 24988 SE Stark, Gresham OR 97030
	<input type="checkbox"/> TUALATIN Legacy Meridian Park campus	Medical Office Building 2 – Suite 140 19260 SW 65th Ave, Tualatin OR 97062