

## Zynyz® (retifanlimab-dlwr) (Intravenous)

-E-

Document Number: OHSU HEALTHSERVICES-0746

Date Reviewed: 12/2025

Date of Origin: 06/04/2024

Dates Approved: 06/2024, 08/2024, 11/05/2024, 02/04/2025, 05/05/2025, 06/24/2025, 08/05/2025, 11/04/2025, 02/03/2026

### I. Length of Authorization <sup>Δ 1,13-15</sup>

- Initial: Prior authorization validity will be provided initially for 6 months (180 days).
- Renewal: Prior authorization validity may be renewed every 6 months (180 days) thereafter for up to a maximum of 24 months (26 total doses) of therapy, unless otherwise specified.
  - Squamous Cell Carcinoma of the Anal Canal in combination with chemotherapy: Prior authorization validity may be renewed for up to a maximum of 12 months (13 total doses).

### II. Dosing Limits

**Max Units (per dose and over time) [HCPCS Unit]:**

- 500 billable units every 4 weeks

### III. Initial Approval Criteria <sup>1</sup>

Prior authorization validity is provided in the following conditions:

- Patient is at least 18 years of age; **AND**

**Universal Criteria** <sup>1,2</sup>

- Patient has not received previous therapy with a programmed death (PD-1/PD-L1)-directed therapy, unless otherwise specified <sup>A</sup>; **AND**
- Used as single agent therapy, unless otherwise specified; **AND**

**Squamous Cell Carcinoma of the Anal Canal (SCAC) † ‡ Φ<sup>1-2,9-10,16</sup>**

- Used in combination with carboplatin and paclitaxel for first-line treatment of inoperable locally recurrent or metastatic disease †; **OR**
- Used for locally recurrent or metastatic disease with progression on or intolerance to platinum-based chemotherapy

**Merkel Cell Carcinoma (MCC) † Φ<sup>1-4,1e</sup>**

- Used as first-line therapy; **AND**
  - Patient has M1 disseminated disease †; **OR**
  - Patient has recurrent locally advanced disease †; **AND**
    - Both curative surgery and curative radiation therapy are not feasible; **OR**
  - Patient has recurrent regional disease ‡; **AND**
    - Both curative surgery and curative radiation therapy are not feasible **OR**
  - Patient has recurrent in-transit N+ regional disease ‡; **AND**
    - Both curative surgery and curative radiation therapy are not feasible

**Preferred therapies and recommendations are determined by review of clinical evidence. NCCN category of recommendation is taken into account as a component of this review. Regimens deemed equally efficacious (i.e., those having the same NCCN categorization) are considered to be therapeutically equivalent.**

**Enhanced Oncology Value (EOV) Program – Redacted indications**

Uses not listed above have inadequate data to support efficacy and are excluded from prior authorization validity.

Other treatment options including, but are not limited to, the following may be appropriate: radiation therapy, surgery, traditional chemotherapy (e.g., platinum, taxane), compassionate use/expanded access programs, clinical trials, supportive care, integrative and complementary therapies.

† FDA Approved Indication(s); ‡ Compendia Recommended Indication(s); Φ Orphan Drug

#### IV. Renewal Criteria<sup>Δ 1,2</sup>

Prior authorization validity may be renewed based upon the following criteria:

- Patient continues to meet the universal and other indication-specific relevant criteria identified in section III; **AND**
- Duration of authorization has not been exceeded (*refer to Section I*); **AND**

- Disease response with treatment as defined by stabilization of disease or decrease in size of tumor or tumor spread; **AND**
- Absence of unacceptable toxicity from the drug. Examples of unacceptable toxicity include: severe infusion-related reactions, severe immune-mediated adverse reactions (e.g., pneumonitis, hepatitis, colitis, endocrinopathies, nephritis with renal dysfunction, dermatologic adverse reactions/rash, etc.), complications of allogeneic hematopoietic stem cell transplantation (HSCT), solid organ transplant rejection, etc.

<sup>Δ</sup> <u>Notes:</u>	
•	Patients responding to therapy who relapse ≥ 6 months after discontinuation due to duration (i.e., receipt of 24 months of therapy) are eligible to re-initiate PD-directed therapy.
•	Patients previously presenting with aggressive disease who are exhibiting stable disease on treatment as their best response (or if therapy improved performance status) may be eligible for continued therapy beyond the 24-month limit without interruption or discontinuation.

## V. Dosage/Administration <sup>Δ</sup> 1,9,15,17

Indication	Dose
Squamous Cell Carcinoma of the Anal Canal	<p><u>Combination Therapy:</u> Administer 500 mg intravenously every four weeks until disease progression or unacceptable toxicity, or up to 12 months.</p> <p><u>Monotherapy:</u> Administer 500 mg intravenously every four weeks until disease progression or unacceptable toxicity, or up to 24 months.</p>
All other indications	Administer 500 mg intravenously every four weeks until disease progression or unacceptable toxicity, or up to 24 months.

## VI. Billing Code/Availability Information

HCPCS Code:

- J9345 – Injection, retifanlimab-dlwr, 1 mg; 1 billable unit = 1 mg

NDC:

- Zynyz 500 mg/20 mL solution in a single-dose vial: 50881-0006-xx

## VII. References (STANDARD)

1. Zynyz [package insert]. Wilmington, DE; Incyte Corporation; May 2025. Accessed December 2025.
2. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) retifanlimab-dlwr. National Comprehensive Cancer Network, 2025. The NCCN

Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed December 2025.

3. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Merkel Cell Carcinoma. Version 2.2026. National Comprehensive Cancer Network, 2025. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed December 2025.
4. Grignani G, Rutkowski P, Lebbé C. A Phase 2 Study of Retifanlimab in Patients With Advanced or Metastatic Merkel Cell Carcinoma (POD1UM-201) Presented at the Society for Immunotherapy of Cancer's 36th Annual Meeting Washington, DC •November 10–14, 2021 [Epub ahead of print]
5. Gupta S, Sonpavde G, Grivas P, et al. Defining “platinum-ineligible” patients with metastatic urothelial cancer (mUC). *J Clin Oncol*. 2019 Mar 1;37(7\_suppl):451.
6. Fahrenbruch R, Kintzel P, Bott AM, et al. Dose Rounding of Biologic and Cytotoxic Anticancer Agents: A Position Statement of the Hematology/Oncology Pharmacy Association. *J Oncol Pract*. 2018 Mar;14(3):e130-e136.
7. Hematology/Oncology Pharmacy Association (2019). Intravenous Cancer Drug Waste Issue Brief. Retrieved from [https://www.hoparx.org/documents/287/HOPA\\_Drug\\_Waste\\_Issue\\_Brief\\_-\\_Updated\\_02.22.24.pdf](https://www.hoparx.org/documents/287/HOPA_Drug_Waste_Issue_Brief_-_Updated_02.22.24.pdf)
8. Bach PB, Conti RM, Muller RJ, et al. Overspending driven by oversized single dose vials of cancer drugs. *BMJ*. 2016 Feb 29;352:i788.
9. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Anal Carcinoma. Version 5.2025. National Comprehensive Cancer Network, 2025. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed December 2025.
10. Rao S, Anandappa G, Capdevila J, et al. A phase II study of retifanlimab (INCMGA00012) in patients with squamous carcinoma of the anal canal who have progressed following platinum-based chemotherapy (POD1UM-202). *ESMO Open*. 2022 Aug;7(4):100529. doi: 10.1016/j.esmoop.2022.100529. Epub 2022 Jul 8. PMID: 35816951; PMCID: PMC9463376.
11. Berton D, Pautier P, Lorusso D, et al. Antitumor activity and safety of the PD-1 inhibitor retifanlimab in patients with recurrent microsatellite instability-high or deficient mismatch repair endometrial cancer: Final safety and efficacy results from cohort H of the POD1UM-101

- phase I study. *Gynecol Oncol* 2024 Jul;186:191-198. doi: 10.1016/j.ygyno.2024.05.025. Epub 2024 Jun 1.
12. Lakhani N, Cosman R, Banerji U, et al. A first-in-human phase I study of the PD-1 inhibitor, retifanlimab (INCMGA00012), in patients with advanced solid tumors (POD1UM-101). *ESMO Open*. 2024 Apr;9(4):102254. doi: 10.1016/j.esmoop.2024.102254. Epub 2024 Feb 21.
  13. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Small Bowel Adenocarcinoma. Version 4.2025. National Comprehensive Cancer Network, 2025. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed December 2025.
  14. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Colon Cancer. Version 5.2025. National Comprehensive Cancer Network, 2025. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed December 2025.
  15. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Rectal Cancer. Version 4.2025. National Comprehensive Cancer Network, 2025. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed December 2025.
  16. Rao S, Jones M, Bowman J, et al. POD1UM-303/InterAACT 2: A phase III, global, randomized, double-blind study of retifanlimab or placebo plus carboplatin-paclitaxel in patients with locally advanced or metastatic squamous cell anal carcinoma. *Front Oncol*. 2022 Aug 24;12:935383. doi: 10.3389/fonc.2022.935383. PMID: 36091159; PMCID: PMC9449327.
  17. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Appendiceal Neoplasms and Cancers. Version 1.2026. National Comprehensive Cancer Network, 2025. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed December 2025.

## VIII. References (ENHANCED)

- 1e. Grignani G, Rutkowski P, Lebbé C. 1146P Updated results from POD1UM-201: A phase II study of retifanlimab in patients with advanced or metastatic Merkel cell carcinoma (MCC). Presented at the European Society for Medical Oncology Congress 2023. Madrid, Spain; October 20-24, 2023.
- 2e. Kaufman HL, Russell JS, Hamid O, et al. Updated efficacy of avelumab in patients with previously treated metastatic Merkel cell carcinoma after  $\geq 1$  year of follow-up: JAVELIN Merkel 200, a phase 2 clinical trial. *Journal for ImmunoTherapy of Cancer*. 2018;6(1). doi:<https://doi.org/10.1186/s40425-017-0310-x>.
- 3e. D'Angelo SP, Russell J, Lebbé C, et al. Efficacy and Safety of First-line Avelumab Treatment in Patients With Stage IV Metastatic Merkel Cell Carcinoma. *JAMA Oncology*. 2018;4(9):e180077. doi:<https://doi.org/10.1001/jamaoncol.2018.0077>.
- 4e. Nghiem PT, Bhatia S, Lipson EJ, et al. PD-1 Blockade with Pembrolizumab in Advanced Merkel-Cell Carcinoma. *The New England journal of medicine*. 2016;374(26):2542-2552. doi:<https://doi.org/10.1056/NEJMoa160370>.
- 5e. Ott PA, Piha-Paul SA, Munster P, et al. Safety and antitumor activity of the anti-PD-1 antibody pembrolizumab in patients with recurrent carcinoma of the anal canal. *Annals of Oncology: Official Journal of the European Society for Medical Oncology*. 2017;28(5):1036-1041. doi:<https://doi.org/10.1093/annonc/mdx029>.
- 6e. Aurelien Marabelle, Cassier PA, Fakih M, et al. Pembrolizumab for previously treated advanced anal squamous cell carcinoma: results from the non-randomised, multicohort, multicentre, phase 2 KEYNOTE-158 study. *The Lancet Gastroenterology & hepatology*. 2022;7(5):446-454. doi:[https://doi.org/10.1016/s2468-1253\(21\)00382-4](https://doi.org/10.1016/s2468-1253(21)00382-4).
- 7e. Morris VK, Salem ME, Nimeiri H, et al. Nivolumab for previously treated unresectable metastatic anal cancer (NCI9673): a multicentre, single-arm, phase 2 study. *The Lancet Oncology*. 2017;18(4):446-453. doi:[https://doi.org/10.1016/S1470-2045\(17\)30104-3](https://doi.org/10.1016/S1470-2045(17)30104-3).
- 8e. Overman MJ, McDermott R, Leach JL, et al. Nivolumab in patients with metastatic DNA mismatch repair-deficient or microsatellite instability-high colorectal cancer (CheckMate 142): an open-label, multicentre, phase 2 study. *The Lancet Oncology*. 2017;18(9):1182-1191. doi:[https://doi.org/10.1016/S1470-2045\(17\)30422-9](https://doi.org/10.1016/S1470-2045(17)30422-9).
- 9e. Andre T, Shiu KK, Kim TW, et al. Pembrolizumab versus chemotherapy for microsatellite instability-high/mismatch repair deficient metastatic colorectal cancer: The phase 3 KEYNOTE-177 Study. *Journal of Clinical Oncology*. 2020;38(18\_suppl):LBA4-LBA4. doi:[https://doi.org/10.1200/jco.2020.38.18\\_suppl.lba4](https://doi.org/10.1200/jco.2020.38.18_suppl.lba4).
- 10e. Chung HC, Ros W, Delord JP, et al. Efficacy and Safety of Pembrolizumab in Previously Treated Advanced Cervical Cancer: Results From the Phase II KEYNOTE-158 Study. *Journal of Clinical Oncology*. 2019;37(17):1470-1478. doi:<https://doi.org/10.1200/jco.18.01265>.

- 11e. Oaknin A, Tinker AV, Gilbert L, et al. Clinical Activity and Safety of the Anti–Programmed Death 1 Monoclonal Antibody Dostarlimab for Patients With Recurrent or Advanced Mismatch Repair–Deficient Endometrial Cancer. *JAMA Oncology*. 2020;6(11):1766. doi:<https://doi.org/10.1001/jamaoncol.2020.4515>.
- 12e. Lakhani N, Cosman R, Banerji U, et al. A first-in-human phase I study of the PD-1 inhibitor, retifanlimab (INCMGA00012), in patients with advanced solid tumors (POD1UM-101). *ESMO Open*. 2024;9(4):102254-102254. doi:<https://doi.org/10.1016/j.esmooop.2024.102254>.
- 13e. Migden MR, Khushalani NI, Chang ALS, et al. Cemiplimab in locally advanced cutaneous squamous cell carcinoma: results from an open-label, phase 2, single-arm trial. *The Lancet Oncology*. 2020;21(2):294-305. doi:[https://doi.org/10.1016/s1470-2045\(19\)30728-4](https://doi.org/10.1016/s1470-2045(19)30728-4).
- 14e. Xia F, Wang Y, Wang H, et al. Randomized Phase II Trial of Immunotherapy-Based Total Neoadjuvant Therapy for Proficient Mismatch Repair or Microsatellite Stable Locally Advanced Rectal Cancer (TORCH). *Journal of Clinical Oncology*. 2024;42(28):3308-3318. doi:<https://doi.org/10.1200/jco.23.02261>.
- 15e. Migden MR, Rischin D, Schmults CD, et al. PD-1 Blockade with Cemiplimab in Advanced Cutaneous Squamous-Cell Carcinoma. *New England Journal of Medicine*. 2018;379(4):341-351. doi:<https://doi.org/10.1056/nejmoa1805131>.
- 16e. André T, Berton D, Curigliano G, et al. Antitumor Activity and Safety of Dostarlimab Monotherapy in Patients With Mismatch Repair Deficient Solid Tumors: A Nonrandomized Controlled Trial. *JAMA network open*. 2023;6(11):e2341165. doi:<https://doi.org/10.1001/jamanetworkopen.2023.41165>.
- 17e. Li J, Xu Y, Zang A, et al. Tislelizumab in previously treated, locally advanced unresectable/metastatic microsatellite instability-high/mismatch repair-deficient solid tumors. *PubMed*. 2024;36(3):257-269. doi:<https://doi.org/10.21147/j.issn.1000-9604.2024.03.03>.
- 18e. Chen Y, Wang Y, Zhang H, et al. Short-course radiotherapy combined with chemotherapy and PD-1 inhibitor in low-lying early rectal cancer: study protocol for a single-arm, multicentre, prospective, phase II trial (TORCH-E). *BMJ Open*. 2023;13(10):e076048-e076048. doi:<https://doi.org/10.1136/bmjopen-2023-076048>.
- 19e. Topalian SL, Bhatia S, Amin A, et al. Neoadjuvant nivolumab for patients with resectable Merkel cell carcinoma in the CheckMate 358 Trial. *J Clin Oncol* 2020;38:2476-2487.
- 20e. Le DT, Kim TW, Van Cutsem E, et al. Phase II Open-Label Study of Pembrolizumab in Treatment-Refractory, Microsatellite Instability-High/Mismatch Repair-Deficient Metastatic Colorectal Cancer: KEYNOTE-164. *J Clin Oncol*. 2020 Jan 1;38(1):11-19. Doi: 10.1200/JCO.19.02107. Epub 2019 Nov 14.
- 21e. Le DT, Uram JN, Wang H, et al. PD-1 Blockade in Tumors with Mismatch-Repair Deficiency. *N Engl J Med*. 2015 Jun 25;372(26):2509-20. doi: 10.1056/NEJMoa1500596. Epub 2015 May 30. PMID: 26028255; PMCID: PMC4481136.

- 22e. Rao S, Samalin-Scalzi E, Evesque L, et al. LBA2 POD1UM303/InterAACT 2: Phase III study of retifanlimab with carboplatinpaclitaxel (c-p) in patients (Pts) with inoperable locally recurrent or metastatic squamous cell carcinoma of the anal canal (SCAC) not previously treated with systemic chemotherapy (Chemo) [abstract]. *Ann Oncol* 2024;35:S1217.
- 23e. Zeitouni NC, Giordano CN, Kane JM 3rd. In-transit Merkel cell carcinoma treated with isolated limb perfusion or isolated limb infusion: a case series of 12 patients. *Dermatol Surg.* 2011 Mar;37(3):357-64. doi: 10.1111/j.1524-4725.2011.01883.x. Epub 2011 Feb 16.
- 24e. Hindmarch JJ, Coker DJ, Waugh R, et al. Treatment of in-transit Merkel cell carcinoma by isolated limb infusion with cytotoxic drugs. *J Surg Case Rep.* 2022 Apr 12;2022(4):rjac172. doi: 10.1093/jscr/rjac172. PMID: 35422991; PMCID: PMC9005209.
- 25e. Kiyohara T, Shijimaya T, Miyamoto M, et al. In-transit recurrence of Merkel cell carcinoma associated with Bowen's disease: The first reported case successfully treated by avelumab. *J Dermatol.* 2019 May;46(5):440-443. doi: 10.1111/1346-8138.14829. Epub 2019 Feb 27.
- 26e. Rusheen J, Clune J, et al. Case report: Metastatic Merkel cell carcinoma presenting seven years after loco-regional disease resection of primary tumor with interval in-transit and nodal metastases. *Front Oncol.* 2023 Jul 5;13:1217816. doi: 10.3389/fonc.2023.1217816. PMID: 37476373; PMCID: PMC10354444.
- 27e. Prime Therapeutics Management. Zynyz Clinical Literature Review Analysis. Last updated December 2025. Accessed December 2025.

## Appendix A – Non-Quantitative Treatment Limitations (NQTL) Factor Checklist

Non-quantitative treatment limitations (NQTLs) refer to the methods, guidelines, standards of evidence, or other conditions that can restrict how long or to what extent benefits are provided under a health plan. These may include things like utilization review or prior authorization. The utilization management NQTL applies comparably, and not more stringently, to mental health/substance use disorder (MH/SUD) Medical Benefit Prescription Drugs and medical/surgical (M/S) Medical Benefit Prescription Drugs. The table below lists the factors that were considered in designing and applying prior authorization to this drug/drug group, and a summary of the conclusions that Prime’s assessment led to for each.

Factor	Conclusion
Indication	Yes: Consider for PA
Safety and efficacy	No: PA not a priority
Potential for misuse/abuse	No: PA not a priority
Cost of drug	Yes: Consider for PA

## Appendix 1 – Covered Diagnosis Codes

ICD-10	ICD-10 Description
C21.0	Malignant neoplasm of anus, unspecified

ICD-10	ICD-10 Description
C21.1	Malignant neoplasm of anal canal
C21.2	Malignant neoplasm of cloacogenic zone
C21.8	Malignant neoplasm of overlapping sites of rectum, anus and anal canal
C4A.0	Merkel cell carcinoma of lip
C4A.10	Merkel cell carcinoma of unspecified eyelid, including canthus
C4A.111	Merkel cell carcinoma of right upper eyelid, including canthus
C4A.112	Merkel cell carcinoma of right lower eyelid, including canthus
C4A.121	Merkel cell carcinoma of left upper eyelid, including canthus
C4A.122	Merkel cell carcinoma of left lower eyelid, including canthus
C4A.20	Merkel cell carcinoma of unspecified ear and external auricular canal
C4A.21	Merkel cell carcinoma of right ear and external auricular canal
C4A.22	Merkel cell carcinoma of left ear and external auricular canal
C4A.30	Merkel cell carcinoma of unspecified part of face
C4A.31	Merkel cell carcinoma of nose
C4A.39	Merkel cell carcinoma of other parts of face
C4A.4	Merkel cell carcinoma of scalp and neck
C4A.51	Merkel cell carcinoma of anal skin
C4A.52	Merkel cell carcinoma of skin of breast
C4A.59	Merkel cell carcinoma of other part of trunk
C4A.60	Merkel cell carcinoma of unspecified upper limb, including shoulder
C4A.61	Merkel cell carcinoma of right upper limb, including shoulder
C4A.62	Merkel cell carcinoma of left upper limb, including shoulder
C4A.70	Merkel cell carcinoma of unspecified lower limb, including hip
C4A.71	Merkel cell carcinoma of right lower limb, including hip
C4A.72	Merkel cell carcinoma of left lower limb, including hip
C4A.8	Merkel cell carcinoma of overlapping sites
C4A.9	Merkel cell carcinoma, unspecified
C7B.1	Secondary Merkel cell carcinoma
Z85.821	Personal history of Merkel cell carcinoma

## Appendix 2 – Centers for Medicare and Medicaid Services (CMS)

The preceding information is intended for non-Medicare coverage determinations. Medicare coverage for outpatient (Part B) drugs is outlined in the Medicare Benefit Policy Manual (Pub. 100-2), Chapter 15, §50 Drugs and Biologicals. In addition, National Coverage Determinations (NCDs) and/or Local Coverage Determinations (LCDs) may exist and compliance with these policies is required where applicable. Local Coverage Articles (LCAs) may also exist for claims payment purposes or to clarify benefit eligibility under

Part B for drugs which may be self-administered. The following link may be used to search for NCD, LCD, or LCA documents: <https://www.cms.gov/medicare-coverage-database/search.aspx>. Additional indications, including any preceding information, may be applied at the discretion of the health plan.

Medicare Part B Covered Diagnosis Codes (applicable to existing NCD/LCD/LCA): N/A

Medicare Part B Administrative Contractor (MAC) Jurisdictions		
Jurisdiction	Applicable State/US Territory	Contractor
E (1)	CA, HI, NV, AS, GU, CNMI	Noridian Healthcare Solutions, LLC
F (2 & 3)	AK, WA, OR, ID, ND, SD, MT, WY, UT, AZ	Noridian Healthcare Solutions, LLC
5	KS, NE, IA, MO	Wisconsin Physicians Service Insurance Corp (WPS)
6	MN, WI, IL	National Government Services, Inc. (NGS)
H (4 & 7)	LA, AR, MS, TX, OK, CO, NM	Novitas Solutions, Inc.
8	MI, IN	Wisconsin Physicians Service Insurance Corp (WPS)
N (9)	FL, PR, VI	First Coast Service Options, Inc.
J (10)	TN, GA, AL	Palmetto GBA
M (11)	NC, SC, WV, VA (excluding below)	Palmetto GBA
L (12)	DE, MD, PA, NJ, DC (includes Arlington & Fairfax counties and the city of Alexandria in VA)	Novitas Solutions, Inc.
K (13 & 14)	NY, CT, MA, RI, VT, ME, NH	National Government Services, Inc. (NGS)
15	KY, OH	CGS Administrators, LLC