

Mylotarg™ (gemtuzumab ozogamicin) (Intravenous)

-E-

Document Number: IC-0491

Date Reviewed: 05/2026**Date of Origin: 09/03/2019****Dates Approved: 09/2019, 12/2019, 11/2020, 12/2021, 12/2022, 12/2023, 11/2024, 05/05/2025, 06/24/2025, 05/05/2026, 06/02/2026**

I. Length of Authorization ^{1,5-8,11}

Newly-Diagnosed AML

- Initial: Prior authorization validity will be provided initially for 6 months (180 days), unless otherwise specified.
 - In combination with daunorubicin and cytarabine (adult): Prior authorization validity will be provided for 6 months consisting of 3 cycles (1 induction and 2 consolidation).
 - In combination with daunorubicin and cytarabine (pediatric): Prior authorization validity will be provided for 6 months consisting of 2 cycles (1 induction and 1 consolidation).
 - In combination with idarubicin and cytarabine (adult): Prior authorization validity will be provided for 6 months consisting of 3 cycles (1 induction and 2 consolidation).
 - In combination with high-dose cytarabine (adult) as consolidation therapy: Prior authorization validity will be provided for 6 months consisting of 2 cycles (2 doses).
- Renewal: Prior authorization validity may NOT be renewed, unless otherwise specified.
 - Single-agent therapy: Prior authorization validity may be renewed for up to a maximum of 8 cycles of continuation.

Relapsed or Refractory AML

- Initial: Prior authorization validity will be provided initially for 6 months (180 days) consisting of 1 cycle (3 doses).
- Renewal: Prior authorization validity may NOT be renewed.

Acute Promyelocytic Leukemia (APL)

- Induction/Consolidation Therapy:
 - Initial: Prior authorization validity will be provided initially for 6 months (180 days) [*1 cycle of induction therapy followed by consolidation therapy*].
 - Renewal: Prior authorization validity may be renewed every 6 months (180 days) thereafter, based on response to therapy (*refer to Section V*).

II. Dosing Limits

Max Units (per dose and over time) [HCPCS Unit]:

- AML:
 - Induction: 135 billable units on Day 1, 90 billable units on Day 4, 90 billable units on Day 7 of a 28-day cycle (1 cycle only)
 - Consolidation/Continuation: 225 billable units every 28 days
- APL:
 - Induction: 180 billable units on Day 1
 - Consolidation: 270 billable units every 28 days

III. Initial Approval Criteria ¹

Prior authorization validity is provided in the following conditions:

- Member is at least 18 years of age (unless otherwise specified); **AND**
- Member has not previously received gemtuzumab ozogamicin (unless otherwise specified); **AND**
- Baseline electrocardiogram (ECG) has been obtained in members with a history of or predisposition for QTc prolongation; **AND**

Universal Criteria ¹

- Member has CD33-positive disease; **AND**

Acute Myeloid Leukemia (AML) † ‡ Φ ^{1,6,10}

- Member has newly-diagnosed disease; **AND**
 - Used in combination with daunorubicin and cytarabine †; **AND**
 - Member is at least 1 month of age; **OR**
 - Used as a single agent †; **OR**
 - Used in combination with cytarabine and idarubicin; **AND**
 - Used as induction or consolidation therapy; **AND**
 - Member has favorable or intermediate-risk AML; **OR**
 - Used in combination with high-dose cytarabine; **AND**
 - Used as consolidation therapy; **AND**
 - Member has favorable-risk AML; **OR**
- Member has relapsed or refractory disease; **AND**

- Used as a single agent †; **AND**
- Member is at least 2 years of age; **OR**
- Member has acute promyelocytic leukemia (APL); **AND**
 - Used for high-risk disease (white blood cell count >10 x 10⁹/L); **AND**
 - Used as induction therapy; **AND**
 - Used in combination with tretinoin (ATRA) and arsenic trioxide (ATO); **OR**
 - Used as consolidation therapy if previously received as induction therapy; **AND**
 - Used in combination with tretinoin (ATRA) or arsenic trioxide (ATO)

Preferred therapies and recommendations are determined by review of clinical evidence. NCCN category of recommendation is taken into account as a component of this review. Regimens deemed equally efficacious (i.e., those having the same NCCN categorization) are considered to be therapeutically equivalent.

Enhanced Oncology Value (EOV) Program – Redacted indications

Uses not listed above have inadequate data to support efficacy and are excluded from coverage.

Other treatment options including, but are not limited to, the following may be appropriate: radiation therapy, surgery, traditional chemotherapy (e.g., platinum, taxane), compassionate use/expanded access programs, clinical trials, supportive care, integrative and complementary therapies.

† FDA Approved Indication(s); ‡ Compendium Recommended Indication(s); Ⓢ Orphan Drug

IV. **Renewal Criteria** ^{1,6}

Prior authorization validity may be renewed based upon the following criteria:

- Member continues to meet the universal and indication-specific relevant criteria such as concomitant therapy requirements (not including prerequisite therapy), performance status, etc. identified in section III; **AND**
- Duration of authorization has not been exceeded (*refer to Section I*); **AND**
- Disease stabilization or improvement as evidenced by a complete response [CR] (i.e. morphologic, cytogenetic or molecular complete response CR), complete hematologic response or a partial response by CBC, bone marrow cytogenetic analysis, QPCR, or FISH; **AND**
- Absence of unacceptable toxicity from the drug. Examples of unacceptable toxicity include: severe infusion-related reactions (including anaphylaxis), hemorrhage, hepatotoxicity (e.g.,

veno-occlusive liver disease [VOD], sinusoidal obstruction syndrome [SOS], etc.), QTc interval prolongation, etc.

V. Dosage/Administration ^{1,5-8,11}

Indication	Dose
Acute Myeloid Leukemia	Newly Diagnosed AML
	<u>Adult (≥ 18 years old) – Combination regimen:</u>
	- Induction Therapy (1 cycle only): - Administer 3 mg/m² (up to one 4.5 mg vial) on Days 1, 4, and 7 in combination with daunorubicin and cytarabine - For members requiring a second induction cycle, do not administer gemtuzumab ozogamicin during the second induction cycle - Consolidation Therapy (maximum of 2 cycles): - Administer 3 mg/m² (up to one 4.5 mg vial) on Day 1 in combination with daunorubicin or idarubicin and cytarabine - Administer 3 mg/m² (up to one 4.5 mg vial) on Day 1 in combination with high-dose cytarabine
	<u>Pediatric (1 month to < 18 years old) – Combination regimen:</u>
	- Induction Therapy (1 cycle only): - Administer 3 mg/m² (BSA ≥ 0.6 m²) or 0.1 mg/kg (BSA < 0.6 m²) on Day 6 in combination with daunorubicin and cytarabine - For members requiring a second induction cycle, do not administer gemtuzumab ozogamicin during the second induction cycle - Consolidation/Intensification Therapy (1 cycle only): - Administer 3 mg/m² (BSA ≥ 0.6 m²) or 0.1 mg/kg (BSA < 0.6 m²) on Day 7 in intensification 2
	<u>Adult (≥ 18 years old) – Single-agent regimen:</u>
	- Induction Therapy (1 cycle only): - Administer 6 mg/m² as a single agent on Day 1 and 3 mg/m² on Day 8 - Continuation Therapy (maximum of 8 cycles): - Administer 2 mg/m² as a single agent on Day 1 every 4 weeks; OR - Administer 6 mg/m² as a single agent on Day 1 and 3 mg/m² on Day 8
	Relapsed or Refractory AML
	Administer 3 mg/m² (up to one 4.5 mg vial) on Days 1, 4, and 7 (1 cycle only)
	Acute Promyelocytic Leukemia (APL)

	<u>High-Risk Disease:</u> • Induction Therapy (1 cycle only): ◦ Administer 6-9 mg/m² on Day 1 (or Day 2, Day 3, or Day 4) in combination with ATRA + ATO • Consolidation Therapy: ◦ ATRA and ATO are used for consolidation. If ATRA or ATO are discontinued due to toxicity, a single dose of gemtuzumab ozogamicin 6 or 9 mg/m² may be given once every 4-5 weeks provided platelets and ANC recover to $\geq 100 \times 10^9/L$ and $\geq 1.0 \times 10^9/L$, respectively, until molecular complete remission.
--	--

VI. Billing Code/Availability Information

HCPCS Code:

- J9203 – Injection, gemtuzumab ozogamicin, 0.1 mg: 1 billable unit = 0.1 mg

NDC:

- Mylotarg 4.5 mg single-dose vial: 00008-4510-xx

VII. References (STANDARD)

1. Mylotarg [package insert]. Philadelphia, PA; Pfizer Inc., August 2021. Accessed March 2026.
2. Castaigne S, Pautas C, Terré C, et al. Effect of gemtuzumab ozogamicin on survival of adult patients with de-novo acute myeloid leukaemia (ALFA-0701): a randomised, open-label, phase 3 study. *Lancet*. 2012 Apr 21;379(9825):1508-16
3. Amadori S, Suci S, Selleslag D, et al. Gemtuzumab Ozogamicin Versus Best Supportive Care in Older Patients With Newly Diagnosed Acute Myeloid Leukemia Unsuitable for Intensive Chemotherapy: Results of the Randomized Phase III EORTC-GIMEMA AML-19 Trial. *J Clin Oncol*. 2016 Mar 20;34(9):972-9.
4. Taksin AL, Legrand O, Raffoux E, et al. High efficacy and safety profile of fractionated doses of Mylotarg as induction therapy in patients with relapsed acute myeloblastic leukemia: A prospective study of the ALFA group. *Leukemia* 2007;21:66–71.
5. Abaza Y, Kantarjian H, Garcia-Mannero G, et al. Long-term outcome of acute promyelocytic leukemia treated with all-transretinoic acid, arsenic trioxide, and gemtuzumab. *Blood* 2017;129:1275-1283.
6. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Acute Myeloid Leukemia. Version 3.2026. National Comprehensive Cancer Network, 2026. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most

recent and complete version of the Compendium, go online to NCCN.org. Accessed March 2026.

7. Burnett AK, Hills RK, Milligan D, et al. Identification of patients with acute myeloblastic leukemia who benefit from the addition of gemtuzumab ozogamicin: results of the MRC AML15 trial. *J Clin Oncol.* 2011 Feb 1;29(4):369-77. doi: 10.1200/JCO.2010.31.4310.
8. Hills RK, Castaigne S, Appelbaum FR, et al. Addition of gemtuzumab ozogamicin to induction chemotherapy in adult patients with acute myeloid leukaemia: a meta-analysis of individual patient data from randomised controlled trials. *Lancet Oncol.* 2014;15(9):986–996. doi:10.1016/S1470-2045(14)70281-5.
9. Gamis AS, Alonzo TA, Meshinchi S, et al. Gemtuzumab ozogamicin in children and adolescents with de novo acute myeloid leukemia improves event-free survival by reducing relapse risk: results from the randomized phase III Children’s Oncology Group trial AAML0531. *J Clin Oncol.* 2014;32(27):3021-3032. doi:10.1200/JCO.2014.55.3628.
10. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) gemtuzumab ozogamicin. National Comprehensive Cancer Network, 2026. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed March 2026.
11. Estey EH, Giles FJ, Beran M, et al. Experience with gemtuzumab ozogamycin ("mylotarg") and all-trans retinoic acid in untreated acute promyelocytic leukemia. *Blood.* 2002 Jun 1;99(11):4222-4. doi: 10.1182/blood-2001-12-0174. PMID: 12010830.
12. Lo-Coco F, Cimino G, Breccia M, et al. Gemtuzumab ozogamicin (Mylotarg) as a single agent for molecularly relapsed acute promyelocytic leukemia. *Blood.* 2004 Oct 1;104(7):1995-9.
13. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Gemtuzumab ozogamicin: Acute Myeloid Leukemia Chemotherapy Order Template, AML88. National Comprehensive Cancer Network, 2026. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed March 2026.

VIII. References (ENHANCED)

- 1e. Burnett AK, Russell NH, Hills RK, et al. Addition of gemtuzumab ozogamicin to induction chemotherapy improves survival in older patients with acute myeloid leukemia. *J Clin Oncol.* 2012 Nov 10;30(32):3924-31. doi: 10.1200/JCO.2012.42.2964. *J Clin Oncol.* 2013 Dec 10;31(35):4424-30. doi: 10.1200/JCO.2013.49.0771.

- 2e. Amadori S, Suci S, Stasi R, et al. Sequential combination of gemtuzumab ozogamicin and standard chemotherapy in older patients with newly diagnosed acute myeloid leukemia: results of a randomized phase III trial by the EORTC and GIMEMA consortium (AML-17).
- 3e. Ravandi F, Estey E, Jones D, et al. Effective treatment of acute promyelocytic leukemia with all-trans-retinoic acid, arsenic trioxide, and gemtuzumab ozogamicin. *J Clin Oncol*. 2009;27(4):504–510. doi:10.1200/JCO.2008.18.6130.
- 4e. Burnett AK, Russell NH, Hills RK, et al. Arsenic trioxide and all-trans retinoic acid treatment for acute promyelocytic leukaemia in all risk groups (AML17): results of a randomised, controlled, phase 3 trial. *Lancet Oncol*. 2015 Oct;16(13):1295-305. doi: 10.1016/S1470-2045(15)00193-X.
- 5e. Lancet JE, Moseley A, Komrokji RS, et al. ATRA, Arsenic Trioxide (ATO), and Gemtuzumab Ozogamicin (GO) Is Safe and Highly Effective in Patients with Previously Untreated High-Risk Acute Promyelocytic Leukemia (APL): Final Results of the SWOG/Alliance/ECOG S0535 Trial. *Blood*. 2016;128:896.
- 6e. Aribi A, Kantarjian HM, Estey EH, et al. Combination therapy with arsenic trioxide, all-trans retinoic acid, and gemtuzumab ozogamicin in recurrent acute promyelocytic leukemia. *Cancer*. 2007 Apr 1;109(7):1355-9.
- 7e. Burnett AK, Hills RK, Milligan D, et al. Identification of patients with acute myeloblastic leukemia who benefit from the addition of gemtuzumab ozogamicin: results of the MRC AML15 trial. *J Clin Oncol*. 2011 Feb 1;29(4):369-77.
- 8e. Jaramillo S, Benner A, Krauter J, et al. Condensed versus standard schedule of high-dose cytarabine consolidation therapy with pegfilgrastim growth factor support in acute myeloid leukemia. *Blood Cancer J* 2017;7:e564.
- 9e. Sanz MA, Montesinos P, Rayon C, et al. Risk-adapted treatment of acute promyelocytic leukemia based on all-trans retinoic acid and anthracycline with addition of cytarabine in consolidation therapy for high-risk patients: further improvements in treatment outcome. *Blood* 2010;115:5137-5146.
- 10e. Pautas C, Merabet F, Thomas X, et al. Randomized study of intensified anthracycline doses for induction and recombinant interleukin-2 for maintenance in patients with acute myeloid leukemia age 50 to 70 years: results of the ALFA-9801 study. *J Clin Oncol*. 2010 Feb 10;28(5):808-14. doi: 10.1200/JCO.2009.23.2652. Epub 2010 Jan 4. PMID: 20048183.
- 11e. Teuffel O, Leibundgut K, Lehrnbecher T, et al. Anthracyclines during induction therapy in acute myeloid leukaemia: a systematic review and meta-analysis. *Br J Haematol*. 2013 Apr;161(2):192-203. doi: 10.1111/bjh.12233. Epub 2013 Feb 8. PMID: 23398482.
- 12e. Prime Therapeutics Management. Mylotarg Clinical Literature Review Analysis. Last updated March 2026. Accessed March 2026.

Appendix A – Non-Quantitative Treatment Limitations (NQTL) Factor Checklist

Page 7

Medical Necessity Criteria

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2026 Prime Therapeutics Management, LLC



Non-quantitative treatment limitations (NQTLs) refer to the methods, guidelines, standards of evidence, or other conditions that can restrict how long or to what extent benefits are provided under a health plan. These may include things like utilization review or prior authorization. The utilization management NQTL applies comparably, and not more stringently, to mental health/substance use disorder (MH/SUD) Medical Benefit Prescription Drugs and medical/surgical (M/S) Medical Benefit Prescription Drugs. The table below lists the factors that were considered in designing and applying prior authorization to this drug/drug group, and a summary of the conclusions that Prime's assessment led to for each.

Factor	Conclusion
Indication	Yes: Consider for PA
Safety and efficacy	Yes: Consider for PA
Potential for misuse/abuse	No: PA not a priority
Cost of drug	Yes: Consider for PA

Appendix 1 – Covered Diagnosis Codes

ICD-10	ICD-10 Description
C92.00	Acute myeloblastic leukemia not having achieved remission
C92.01	Acute myeloblastic leukemia in remission
C92.02	Acute myeloblastic leukemia in relapse
C92.40	Acute promyelocytic leukemia not having achieved remission
C92.41	Acute promyelocytic leukemia in remission
C92.42	Acute promyelocytic leukemia in relapse
C92.50	Acute myelomonocytic leukemia not having achieved remission
C92.51	Acute myelomonocytic leukemia in remission
C92.52	Acute myelomonocytic leukemia in relapse
C92.60	Acute myeloid leukemia with 11q23-abnormality not having achieved remission
C92.61	Acute myeloid leukemia with 11q23-abnormality in remission
C92.62	Acute myeloid leukemia with 11q23-abnormality in relapse
C92.A0	Acute myeloid leukemia with multilineage dysplasia not having achieved remission
C92.A1	Acute myeloid leukemia with multilineage dysplasia in remission
C92.A2	Acute myeloid leukemia with multilineage dysplasia in relapse
C93.00	Acute monoblastic/monocytic leukemia not having achieved remission
C93.01	Acute monoblastic/monocytic leukemia in remission

C93.02	Acute monoclastic/monocytic leukemia in relapse
--------	---

Appendix 2 – Centers for Medicare and Medicaid Services (CMS)

The preceding information is intended for non-Medicare coverage determinations. Medicare coverage for outpatient (Part B) drugs is outlined in the Medicare Benefit Policy Manual (Pub. 100-2), Chapter 15, §50 Drugs and Biologicals. In addition, National Coverage Determinations (NCDs) and/or Local Coverage Determinations (LCDs) may exist and compliance with these policies is required where applicable. Local Coverage Articles (LCAs) may also exist for claims payment purposes or to clarify benefit eligibility under Part B for drugs which may be self-administered. The following link may be used to search for NCD, LCD, or LCA documents: <https://www.cms.gov/medicare-coverage-database/search.aspx>. Additional indications, including any preceding information, may be applied at the discretion of the health plan.

Medicare Part B Covered Diagnosis Codes (applicable to existing NCD/LCD/LCA): N/A

Medicare Part B Administrative Contractor (MAC) Jurisdictions		
Jurisdiction	Applicable State/US Territory	Contractor
E (1)	CA, HI, NV, AS, GU, CNMI	Noridian Healthcare Solutions, LLC
F (2 & 3)	AK, WA, OR, ID, ND, SD, MT, WY, UT, AZ	Noridian Healthcare Solutions, LLC
5	KS, NE, IA, MO	Wisconsin Physicians Service Insurance Corp (WPS)
6	MN, WI, IL	National Government Services, Inc. (NGS)
H (4 & 7)	LA, AR, MS, TX, OK, CO, NM	Novitas Solutions, Inc.
8	MI, IN	Wisconsin Physicians Service Insurance Corp (WPS)
N (9)	FL, PR, VI	First Coast Service Options, Inc.
J (10)	TN, GA, AL	Palmetto GBA
M (11)	NC, SC, WV, VA (excluding below)	Palmetto GBA
L (12)	DE, MD, PA, NJ, DC (includes Arlington & Fairfax counties and the city of Alexandria in VA)	Novitas Solutions, Inc.
K (13 & 14)	NY, CT, MA, RI, VT, ME, NH	National Government Services, Inc. (NGS)
15	KY, OH	CGS Administrators, LLC