

Opdualag™ (nivolumab/relatlimab-rmbw) (Intravenous)

-E-

Document Number: IC-0685

Date Reviewed: 06/2026

Date of Origin: 12/01/2022

Dates Approved: 12/2022, 06/2023, 10/2023, 01/04/2024, 05/02/2024, 09/04/2025, 08/04/2026

I. Length of Authorization ^{1,5}

- Initial: Prior authorization validity will be provided initially for 6 months (180 days), unless otherwise specified.
 - Neoadjuvant treatment of Cutaneous Melanoma: Prior authorization validity will be provided initially for 2 doses only.
- Renewal: Prior authorization validity may be renewed every 6 months (180 days) thereafter, unless otherwise specified.
 - Neoadjuvant treatment of Cutaneous Melanoma: Prior authorization validity may NOT be renewed.

II. Dosing Limits

Max Units (per dose and over time) [HCPCS Unit]:

- 160 billable units (480 mg nivolumab/160 mg relatlimab) every 28 days

III. Initial Approval Criteria ¹

Prior authorization validity is provided for the following conditions:

- Member is at least 12 years of age; **AND**

Universal Criteria ¹⁻³

- Member weighs at least 40 kg; **AND**
- Member has not received previous therapy with a programmed death (PD-1/PD-L1)-directed therapy, unless otherwise specified ^Δ; **AND**

Cutaneous Melanoma † ‡ Φ ^{1-4,9e}

- Used as first-line therapy for unresectable or metastatic* disease; **OR**
- Used as subsequent therapy unresectable or metastatic* disease; **AND**

- Used for disease progression, intolerance, and/or projected risk of progression with BRAF-targeted therapy (e.g., dabrafenib/trametinib, vemurafenib/cobimetinib, encorafenib/binimetinib, etc.); **OR**
 - Used as re-induction therapy in members with prior nivolumab/relatlimab that resulted in disease control (*i.e., complete response, partial response, or stable disease*) with no residual toxicity, but with disease progression or relapse >3 months after treatment discontinuation; **OR**
 - Used as neoadjuvant therapy; **AND**
 - Used for one of the following:
 - Member has stage III disease; **AND**
 - Used as primary treatment for clinically positive, resectable nodal disease; **OR**
 - Used for limited resectable disease with clinical satellite/in-transit metastases; **OR**
 - Member has limited resectable local satellite/in-transit recurrence; **OR**
 - Member has resectable disease limited to nodal recurrence; **AND**
- Use of nivolumab/relatlimab will be restricted to members with a contraindication or intolerance to pembrolizumab

**Metastatic disease includes stage III unresectable/borderline resectable disease with clinically positive node(s) or clinical satellite/in-transit metastases, as well as unresectable/borderline resectable local satellite/in-transit recurrence, unresectable nodal recurrence, oligometastatic disease, and widely disseminated distant metastatic disease and/or brain metastases.*

Preferred therapies and recommendations are determined by review of clinical evidence. NCCN category of recommendation is taken into account as a component of this review. Regimens deemed equally efficacious (i.e., those having the same NCCN categorization) are considered to be therapeutically equivalent.

† FDA Approved Indication(s); ‡ Compendia Recommended Indication(s); Ⓢ Orphan Drug

IV. Renewal Criteria ¹⁻⁴

Prior authorization validity may be renewed based upon the following criteria:

- Member continues to meet the universal and other indication-specific relevant criteria such as concomitant therapy requirements (not including prerequisite therapy), performance status, etc. identified in section III; **AND**
- Duration of authorization has not been exceeded (*refer to Section I*); **AND**
- Absence of unacceptable toxicity from the drug. Examples of unacceptable toxicity include: severe infusion-related reactions, complications of allogeneic hematopoietic stem cell

transplantation (HSCT), severe immune-mediated adverse reactions (e.g., pneumonitis, colitis, hepatitis, endocrinopathies, nephritis/renal dysfunction, dermatologic adverse reactions/rash, myocarditis, etc.), etc.; **AND**

- Disease response with treatment as defined by stabilization of disease or decrease in size of tumor or tumor spread

Δ Notes:

- Members who complete adjuvant therapy and progress ≥ 6 months after discontinuation are eligible to re-initiate PD-directed therapy for metastatic disease.
- Members whose tumors, upon re-biopsy, demonstrate a change in actionable mutation (e.g., MSS initial biopsy; MSI-H subsequent biopsy) may be eligible to re-initiate PD-directed therapy and will be evaluated on a case-by-case basis.

V. Dosage/Administration ^{Δ 1,5}

Indication	Dose
Cutaneous Melanoma	Adult and pediatric members ≥ 12 years of age who weigh at least 40 kg: • <u>Neoadjuvant treatment:</u> Administer 480 mg nivolumab and 160 mg relatlimab (contents of 2 vials) intravenously every 4 weeks for 2 doses • <u>All other treatment settings:</u> Administer 480 mg nivolumab and 160 mg relatlimab (contents of 2 vials) intravenously every 4 weeks until disease progression or unacceptable toxicity

VI. Billing Code/Availability Information

HCPSC Code:

- J9298 – Injection, nivolumab and relatlimab-rmbw, 3 mg/1 mg; 1 billable unit = 3 mg nivolumab/1 mg relatlimab-rmbw

NDC:

- Opdualag 240 mg of nivolumab and 80 mg of relatlimab per 20 mL single-dose vial: 00003-7125-xx

VII. References (STANDARD)

1. Opdualag [package insert]. Princeton, NJ; Bristol-Myers Squibb Company; June 2026. Accessed June 2026.
2. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) nivolumab-relatlimab. National Comprehensive Cancer Network, 2026. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL

COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed June 2026.

3. Tawbi HA, Schadendorf D, Lipson EJ; RELATIVITY-047 Investigators, et al. Relatlimab and nivolumab versus nivolumab in untreated advanced melanoma. *N Engl J Med*. 2022;386:24-34.
4. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Melanoma: Cutaneous. Version 2.2026. National Comprehensive Cancer Network, 2026. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed June 2026.
5. Amaria RN, Postow M, Burton EM, et al. Neoadjuvant relatlimab and nivolumab in resectable melanoma. *Nature* 2022;611:155-160. Erratum in: *Nature* 2023;615:E23.

VIII. References (ENHANCED)

- 1e. Robert C, Long GV, Brady B, et al. Nivolumab in previously untreated melanoma without BRAF mutation. *N Engl J Med*. 2015 Jan 22;372(4):320-30. doi: 10.1056/NEJMoa1412082.
- 2e. Larkin J, Chiarion-Sileni V, Gonzalez R, et al. Combined Nivolumab and Ipilimumab or Monotherapy in Untreated Melanoma [published correction appears in *N Engl J Med*. 2018 Nov 29;379(22):2185]. *N Engl J Med*. 2015;373(1):23–34. doi:10.1056/NEJMoa1504030.
- 3e. Robert C, Schachter J, Long GV, et al. Pembrolizumab versus Ipilimumab in Advanced Melanoma. *N Engl J Med*. 2015 Jun 25;372(26):2521-32. doi: 10.1056/NEJMoa1503093.
- 4e. Ascierto PA, Lipson EJ, Dummer R, et al. Nivolumab and Relatlimab in Patients With Advanced Melanoma That Had Progressed on Anti-Programmed Death-1/Programmed Death Ligand 1 Therapy: Results From the Phase I/IIa RELATIVITY-020 Trial. *J Clin Oncol* 2023; :JCO2202072.
- 5e. Burton EM, Milton DR, Tetzlaff MT, et al. Long-Term Survival and Biomarker Analysis Evaluating Neoadjuvant Plus Adjuvant Relatlimab (anti-LAG3) and Nivolumab (anti-PD1) in Patients With Resectable Melanoma. *J Clin Oncol*. 2025 Jul 10;JCO2500494.
- 6e. Versluis JM, Menzies AM, Sikorska K, et al. Survival update of neoadjuvant ipilimumab plus nivolumab in macroscopic stage III melanoma in the OpACIN and OpACIN-neo trials. *Ann Oncol*. 2023 Apr;34(4):420-430.
- 7e. Tawbi H. Nivolumab + relatlimab (NIVO + RELA) in advanced melanoma: 5-year update of RELATIVITY-047. - ASCO. Asco.org. Published 2026. Accessed June 26, 2026. <https://www.asco.org/abstracts-presentations/260898>
- 8e. Regan MM, Ascierto PA MD, Lipson EJ, et al. Analysis of treatment-free survival of patients with advanced melanoma receiving nivolumab as monotherapy or in combination with relatlimab in

RELATIVITY-047. *J Immunother Cancer*. 2025;13(9):e012747. Published 2025 Sep 12. doi:10.1136/jitc-2025-012747.

9e. Patel SP, Othus M, Chen Y, et al. Neoadjuvant-Adjuvant or Adjuvant-Only Pembrolizumab in Advanced Melanoma. *N Engl J Med*. 2023;388(9):813-823. doi:10.1056/NEJMoa2211437

10e. Prime Therapeutics Management. Opdualag Clinical Literature Review Analysis. Last updated June 2026. Accessed June 2026.

Appendix A – Non-Quantitative Treatment Limitations (NQTL) Factor Checklist

Non-quantitative treatment limitations (NQTLs) refer to the methods, guidelines, standards of evidence, or other conditions that can restrict how long or to what extent benefits are provided under a health plan. These may include things like utilization review or prior authorization. The utilization management NQTL applies comparably, and not more stringently, to mental health/substance use disorder (MH/SUD) Medical Benefit Prescription Drugs and medical/surgical (M/S) Medical Benefit Prescription Drugs. The table below lists the factors that were considered in designing and applying prior authorization to this drug/drug group, and a summary of the conclusions that Prime's assessment led to for each.

Factor	Conclusion
Indication	Yes: Consider for PA
Safety and efficacy	No: PA not a priority
Potential for misuse/abuse	No: PA not a priority
Cost of drug	Yes: Consider for PA

Appendix 1 – Covered Diagnosis Codes

ICD-10	ICD-10 Description
C43.0	Malignant melanoma of lip
C43.10	Malignant melanoma of unspecified eyelid, including canthus
C43.111	Malignant melanoma of right upper eyelid, including canthus
C43.112	Malignant melanoma of right lower eyelid, including canthus
C43.121	Malignant melanoma of left upper eyelid, including canthus
C43.122	Malignant melanoma of left lower eyelid, including canthus
C43.20	Malignant melanoma of unspecified ear and external auricular canal
C43.21	Malignant melanoma of right ear and external auricular canal
C43.22	Malignant melanoma of left ear and external auricular canal
C43.30	Malignant melanoma of unspecified part of face

C43.31	Malignant melanoma of nose
C43.39	Malignant melanoma of other parts of face
C43.4	Malignant melanoma of scalp and neck
C43.51	Malignant melanoma of anal skin
C43.52	Malignant melanoma of skin of breast
C43.59	Malignant melanoma of other part of trunk
C43.60	Malignant melanoma of unspecified upper limb, including shoulder
C43.61	Malignant melanoma of right upper limb, including shoulder
C43.62	Malignant melanoma of left upper limb, including shoulder
C43.70	Malignant melanoma of unspecified lower limb, including hip
C43.71	Malignant melanoma of right lower limb, including hip
C43.72	Malignant melanoma of left lower limb, including hip
C43.8	Malignant melanoma of overlapping sites of skin
C43.9	Malignant melanoma of skin, unspecified

Appendix 2 – Centers for Medicare and Medicaid Services (CMS)

The preceding information is intended for non-Medicare coverage determinations. Medicare coverage for outpatient (Part B) drugs is outlined in the Medicare Benefit Policy Manual (Pub. 100-2), Chapter 15, §50 Drugs and Biologicals. In addition, National Coverage Determinations (NCDs) and/or Local Coverage Determinations (LCDs) may exist and compliance with these policies is required where applicable. Local Coverage Articles (LCAs) may also exist for claims payment purposes or to clarify benefit eligibility under Part B for drugs which may be self-administered. The following link may be used to search for NCD, LCD, or LCA documents: <https://www.cms.gov/medicare-coverage-database/search.aspx>. Additional indications, including any preceding information, may be applied at the discretion of the health plan.

Medicare Part B Covered Diagnosis Codes (applicable to existing NCD/LCD/LCA): N/A

Medicare Part B Administrative Contractor (MAC) Jurisdictions		
Jurisdiction	Applicable State/US Territory	Contractor
E (1)	CA, HI, NV, AS, GU, CNMI	Noridian Healthcare Solutions, LLC
F (2 & 3)	AK, WA, OR, ID, ND, SD, MT, WY, UT, AZ	Noridian Healthcare Solutions, LLC
5	KS, NE, IA, MO	Wisconsin Physicians Service Insurance Corp (WPS)
6	MN, WI, IL	National Government Services, Inc. (NGS)
H (4 & 7)	LA, AR, MS, TX, OK, CO, NM	Novitas Solutions, Inc.

Medicare Part B Administrative Contractor (MAC) Jurisdictions		
Jurisdiction	Applicable State/US Territory	Contractor
8	MI, IN	Wisconsin Physicians Service Insurance Corp (WPS)
N (9)	FL, PR, VI	First Coast Service Options, Inc.
J (10)	TN, GA, AL	Palmetto GBA
M (11)	NC, SC, WV, VA (excluding below)	Palmetto GBA
L (12)	DE, MD, PA, NJ, DC (includes Arlington & Fairfax counties and the city of Alexandria in VA)	Novitas Solutions, Inc.
K (13 & 14)	NY, CT, MA, RI, VT, ME, NH	National Government Services, Inc. (NGS)
15	KY, OH	CGS Administrators, LLC