

Tecentriq® (atezolizumab) (Intravenous)

-E-

Document Number: IC-0388

Date Reviewed: 06/2026

Date of Origin: 08/05/2019

Dates Approved: 08/2019, 10/2019, 01/2020, 04/2020, 07/2020, 09/2020, 01/2021, 04/2021, 06/2021, 10/2021, 12/2021, 02/2022, 04/2022, 07/2022, 10/2022, 01/2023, 04/2023, 07/2023, 10/2023, 01/2024, 04/2024, 08/2024, 11/2024, 02/04/2025, 05/05/2025, 06/24/2025, 08/05/2025, 11/04/2025, 12/02/2025, 02/03/2026, 05/05/2026, 07/01/2026, 08/04/2026

I. Length of Authorization ^{Δ 1,23}

- Initial: Prior authorization validity will be provided initially for 6 months (180 days).
- Renewal: Prior authorization validity may be renewed every 6 months (180 days) thereafter, unless otherwise specified.
 - Colon Cancer adjuvant therapy: Prior authorization validity may be renewed up to a maximum of 12 months of therapy.*
 - Small Bowel Adenocarcinoma adjuvant therapy: Prior authorization validity may be renewed up to a maximum of 12 months of therapy.*
 - Non-Small Cell Lung Cancer (NSCLC) and Urothelial Carcinoma (Bladder Cancer) when used as adjuvant therapy: Prior authorization validity may be renewed up to a maximum of 12 months of therapy.*
 - Thymic carcinoma: Prior authorization validity may be renewed up to a maximum of 24 months of therapy.*
 - Chronic Lymphocytic Leukemia/Small Lymphocytic Lymphoma (CLL/SLL): Prior authorization validity may be renewed up to a maximum of 18 cycles.

***Note: The maximum number of doses is dependent on the dosing frequency and duration of therapy. Refer to Section V for exact dosage.**

Dosing Frequency	Maximum length of therapy	Maximum number of doses
2 weeks	1 year	26 doses
3 weeks	1 year	18 doses
	2 years	35 doses
4 weeks	1 year	13 doses

II. Dosing Limits

Max Units (per dose and over time) [HCPCS Unit]:

- Mesotheliomas (*peritoneal, pericardial, & tunica vaginalis testis*), Thymic Carcinoma, and CLL/SLL: 120 billable units every 21 days
- All other indications: 504 billable units every 84 days

III. Initial Approval Criteria ¹

Prior authorization validity is provided in the following conditions:

- Member is at least 18 years of age (unless otherwise specified); **AND**

Universal Criteria

- Member has not received previous therapy with a programmed death (PD-1/PD-L1)-directed therapy unless otherwise specified ^Δ (*Note: Not applicable when used as switch-therapy with subcutaneous atezolizumab*); **AND**
- Therapy will not be used concomitantly with subcutaneous atezolizumab; **AND**

Non-Small Cell Lung Cancer (NSCLC) † ‡ ^{1,5,6,8,11,12,17,23,9e-11e,14e,49e}

- Used for recurrent, advanced, or metastatic disease; **AND**
 - Used as first-line therapy; **AND**
 - Used as a single agent; **AND**
 - Members with performance status (PS) 0-2 who have tumors that are negative for actionable biomarkers* (may be KRAS G12C mutation positive) and PD-L1 ≥50% (*PD-L1 stained ≥ 50% of tumor cells [TC ≥ 50%] or PD-L1 stained tumor-infiltrating immune cells [IC] covering ≥ 10% of the tumor area [IC ≥ 10%]*), as determined by an FDA-approved test or Clinical Laboratory Improvement Amendments (CLIA)-compliant test❖; **AND**

– Use of atezolizumab will be restricted to members with a contraindication or intolerance to single-agent cemiplimab; **OR**
 - Members with PS 3 who have tumors that are negative for actionable biomarkers* (may be KRAS G12C mutation positive) regardless of PD-L1 status; **AND**
 - Member is platinum-ineligible; **OR**
 - Members with PS 3 who have tumors positive for one of the following biomarkers: EGFR exon 20, BRAF V600E, NTRK1/2/3 gene fusion, MET exon-14 skipping, ERBB2 (HER2), NRG1 gene fusion; **AND**
 - Member is platinum-ineligible; **OR**
 - Used in combination with one of the following:

- Carboplatin, paclitaxel, and bevacizumab
- Carboplatin and albumin-bound paclitaxel; **AND**
- Used for non-squamous disease; **AND**
 - Tumor is negative for actionable biomarkers* (may be KRAS G12C mutation positive); **AND**

◆ Use of atezolizumab will be restricted to members with a contraindication or intolerance to cemiplimab/(paclitaxel or pemetrexed)/(carboplatin or cisplatin); **OR**
 - Tumor is positive for one of the following biomarkers: BRAF V600E, NTRK1/2/3 gene fusion, MET exon-14 skipping, ERBB2 (HER2), or NRG1 gene; **AND**

◆ Use of atezolizumab will be restricted to members with a contraindication or intolerance to cemiplimab/(paclitaxel or pemetrexed)/(carboplatin or cisplatin); **OR**
 - Tumor is positive for EGFR exon 20; **AND**

In combination with carboplatin and albumin-bound paclitaxel ONLY:

◆ Use of atezolizumab will be restricted to members with a contraindication or intolerance to cemiplimab/(paclitaxel or pemetrexed)/(carboplatin or cisplatin); **OR**
- Used as subsequent therapy; **AND**
 - Used as a single agent; **AND**
 - Members with PS 0-2; **OR**
 - Members with PS 3 who are positive for one of the following biomarkers: BRAF V600E, NTRK1/2/3 gene fusion, MET exon-14 skipping, or ERBB2 (HER2); **AND**
 - Member is platinum-ineligible; **OR**
 - Members with PS 3 who are positive for one of the following biomarkers and received prior targeted therapy§: EGFR exon 19 deletion or L858R, EGFR S768I, L861Q and/or G719X, RET gene fusion, ALK gene fusion, or ROS1 gene fusion; **AND**
 - Member is platinum-ineligible; **OR**
 - Used in combination with one of the following:
 - Carboplatin, paclitaxel, and bevacizumab
 - Carboplatin and albumin-bound paclitaxel; **AND**
 - Used for non-squamous disease; **AND**
 - Used for one of the following:

- ◆ Member is positive for one of the following biomarkers: BRAF V600E, NTRK1/2/3 gene fusion, MET exon-14 skipping, or ERBB2 (HER2); **OR**
- ◆ Member is positive for one of the following biomarkers and received prior targeted therapy§: EGFR S768I, L861Q, and/or G719X mutation; **AND**

In combination with carboplatin and albumin-bound paclitaxel ONLY:

- Use of atezolizumab will be restricted to members with a contraindication or intolerance to cemiplimab/(paclitaxel or pemetrexed)/(carboplatin or cisplatin); **OR**
- Used as continuation maintenance therapy in members who have achieved a tumor response or stable disease following initial therapy; **AND**
 - Used in combination with bevacizumab following a first-line regimen with atezolizumab, carboplatin, paclitaxel, and bevacizumab for non-squamous histology; **OR**
 - Used as a single agent following a first-line regimen with atezolizumab, carboplatin, and albumin-bound paclitaxel for non-squamous histology; **OR**
 - Used as a single agent following a first-line regimen with single agent atezolizumab; **OR**
- Used as adjuvant therapy as a single agent; **AND**
 - Tumor expresses PD-L1 ≥1% as determined by an FDA-approved test or CLIA-compliant test❖; **AND**
 - Used following resection and previous adjuvant platinum-based chemotherapy; **AND**
 - Member has stage II to IIIA disease †; **OR**
 - Member has stage IB or IIIB (T2-T3, N2b; T4, N2) disease ‡; **AND**
 - Member has no known EGFR mutations, or ALK gene fusion

**Note: Actionable biomarkers include EGFR, KRAS, ALK, ROS1, BRAF, NTRK1/2/3, MET, RET, NRG1, and ERBB2 (HER2). Complete biomarker testing including molecular assessment of EGFR, KRAS, ALK, ROS1, BRAF, NTRK1/2/3, MET, RET, NRG1, and ERBB2 (HER2), via biopsy and/or plasma testing. If a clinically actionable marker is found, it is reasonable to start therapy based on the identified marker. Treatment is guided by available results and, if unknown, these members are treated as though they do not have driver oncogenes.*

§ Note: Genomic Aberration/Mutational Driver Targeted Therapies: Refer to guidelines for appropriate use.

Small Cell Lung Cancer (SCLC) † ‡ Φ^{1,6,14,18,39}

- Member has extensive stage disease (ES-SCLC); **AND**
 - Used as first-line therapy in combination with carboplatin and etoposide †; **OR**
 - Used as maintenance therapy if disease has not progressed following 4 cycles of first-line therapy with atezolizumab (intravenous or subcutaneous), carboplatin and etoposide; **AND**

- Used in combination with lurbinectedin †; **OR**
- Used as a single agent ‡; **OR**
- Member has disease progression or relapse after a prolonged disease-free interval ‡; **AND**
 - Used as subsequent therapy in combination with carboplatin and etoposide; **OR**
 - Used as maintenance therapy; **AND**
 - Used as a single agent following 4 cycles of subsequent therapy with atezolizumab (intravenous or subcutaneous), carboplatin, and etoposide; **OR**
 - Used in combination with lurbinectedin (if lurbinectedin has not been used previously); **AND**
 - Member has at least stable disease following 4 cycles of subsequent therapy with atezolizumab (intravenous or subcutaneous), carboplatin, and etoposide; **AND**
 - Member has an Eastern Cooperative Oncology Group (ECOG) performance status of 0-1; **AND**
 - Member has no history of brain metastases

Hepatocellular Carcinoma (HCC) † ‡ Φ 1,6,15,16,21,30e

- Used in combination with bevacizumab; **AND**
 - Member does not have Child-Turcotte-Pugh (CTP) Class C liver disease; **AND**
 - Used as first-line therapy; **AND**
 - Used for one of the following:
 - Member has unresectable disease †; **OR**
 - Member has extrahepatic/metastatic disease and is deemed ineligible for resection, transplant, or locoregional therapy; **AND**
- Use of atezolizumab will be restricted to members with a contraindication or intolerance to durvalumab/tremelimumab; **OR**
- Used as subsequent-line therapy for disease progression on or after systemic therapy

Peritoneal Mesothelioma (PeM) ‡ 6,24,27,22e**

- Used as subsequent therapy in combination with bevacizumab; **AND**
 - Member received previous treatment with platinum and pemetrexed, unless contraindicated; **AND**
- Member must demonstrate an inadequate response to nivolumab, unless there is a contraindication or intolerance, prior to approval of atezolizumab

**** Note:** May also be used for pericardial mesothelioma and tunica vaginalis testis mesothelioma.

Cutaneous Melanoma † ‡ Φ^{1,6,19,20,29}

- Member has BRAF V600 mutation positive disease as detected by an FDA approved or CLIA compliant test❖; **AND**
- Used in combination with cobimetinib and vemurafenib; **AND**
- Member has unresectable or metastatic disease; **AND**
 - Used as first-line therapy; **OR**
 - Used as re-induction therapy in members who experienced disease control (i.e., complete response, partial response, or stable disease with no residual toxicity) from prior combination cobimetinib/vemurafenib + atezolizumab therapy, but subsequently have disease progression/relapse > 3 months after treatment discontinuation

Alveolar Soft Part Sarcoma (ASPS) † ‡ Φ^{1,6,26}

- Member is at least 2 years of age; **AND**
- Used as a single agent; **AND**
- Member has unresectable or metastatic disease that is not curable by surgery

Cervical Cancer ‡^{6,14,37}

- Member has small cell neuroendocrine carcinoma of the cervix (NECC); **AND**
 - Used as first-line therapy for persistent, recurrent, or metastatic disease; **AND**
 - Used in combination with etoposide AND either cisplatin or carboplatin; **OR**
 - Used as single-agent maintenance therapy after initial therapy with atezolizumab, etoposide, AND either carboplatin or cisplatin; **OR**
- Member has adenocarcinoma, adenosquamous, or squamous cell carcinoma; **AND**
 - Used as first-line therapy; **AND**
 - Member has recurrent or metastatic disease; **AND**
 - Used in combination with bevacizumab, paclitaxel, AND either cisplatin or carboplatin; **OR**
 - Used in combination with bevacizumab as maintenance therapy after initial therapy with atezolizumab, bevacizumab, paclitaxel, AND cisplatin or carboplatin

Colon Cancer ‡^{6,37,38}

- Member has microsatellite instability-high (MSI-H)/deficient mismatch repair (dMMR) disease as determined by an FDA-approved or CLIA-compliant test❖; **AND**

- Used as adjuvant treatment in combination with FOLFOX (fluorouracil, leucovorin, and oxaliplatin) or CAPEOX (capecitabine and oxaliplatin) followed by single agent maintenance therapy; **AND**
 - Member has stage IIC, stage III, or metastatic disease; **OR**
- Used as a single agent for locally unresectable, medically inoperable, advanced or metastatic disease; **AND**
 - Used as initial treatment; **OR**
 - Used in subsequent therapy; **AND**
 - Use of atezolizumab will be restricted to members with a contraindication or intolerance to durvalumab

Rectal Cancer ‡^{6,45}

- Member has MSI-H/dMMR disease as determined by an FDA-approved or CLIA-compliant test❖; **AND**
- Used as a single agent for locally unresectable, medically inoperable, advanced or metastatic disease; **AND**
 - Used as initial treatment; **OR**
 - Used as subsequent therapy; **AND**
 - Use of atezolizumab will be restricted to members with a contraindication or intolerance to durvalumab

Small Bowel Adenocarcinoma ‡^{6,37,44}

- Member has MSI-H/dMMR disease as determined by an FDA-approved or CLIA-compliant test❖; **AND**
 - Used as adjuvant therapy in combination with FOLFOX or CAPEOX, followed by single agent maintenance therapy; **AND**
 - Member has T4, N0, M0 or T Any, N1-2 disease; **OR**
 - Used as a single agent; **AND**
 - Member has advanced or metastatic disease; **AND**
 - Used as initial treatment; **OR**
 - Used as subsequent therapy; **AND**
 - Use of atezolizumab will be restricted to members with a contraindication or intolerance to durvalumab; **OR**
 - Member has locally unresectable or medically inoperable disease; **AND**

- Used as primary treatment

Thymic Carcinoma ‡^{6,41}

- Used in combination with carboplatin and paclitaxel; **AND**
- Used as first-line therapy for recurrent, advanced, or metastatic disease

Chronic Lymphocytic Leukemia/Small Lymphocytic Lymphoma (CLL/SLL) ‡^{6,42}

- Used in combination with venetoclax and obinutuzumab for histologic transformation (Richter); **AND**
 - Used as additional therapy for partial response, refractory disease, or progression while on prior treatment [^]; **OR**
 - Used as first-line treatment for Richter transformation if previously treated for CLL; **OR**
 - Used as continuation therapy for complete response until progression

[^]Prior treatment could have included immune checkpoint inhibitor therapy (e.g., PD-1/PD-L1-directed therapy)

Urothelial Carcinoma (Bladder Cancer) †^{1,46}

- Member has muscle invasive bladder cancer (MIBC); **AND**
- Used as a single agent; **AND**
- Used as adjuvant therapy following cystectomy; **AND**
- Member has circulating tumor DNA molecular residual disease (ctDNA MRD) as determined by an FDA-approved or CLIA-compliant test[❖], detected within 1 year post-cystectomy

Preferred therapies and recommendations are determined by review of clinical evidence. NCCN category of recommendation is taken into account as a component of this review. Regimens deemed equally efficacious (i.e., those having the same NCCN categorization) are considered to be therapeutically equivalent.

Enhanced Oncology Value (EOV) Program – Redacted indications

Uses not listed above have inadequate data to support efficacy and are excluded from prior authorization validity.

Other treatment options including, but are not limited to, the following may be appropriate: radiation therapy, surgery, traditional chemotherapy (e.g., platinum, taxane), compassionate use/expanded access programs, clinical trials, supportive care, integrative and complementary therapies.

❖ If confirmed using an FDA approved assay – <http://www.fda.gov/companiondiagnostics>

† FDA Approved Indication(s); ‡ Compendia Recommended Indication(s); Ⓢ Orphan Drug

IV. Renewal Criteria ^{Δ 1,6}

Prior authorization validity can be renewed based upon the following criteria:

- Member continues to meet the universal and other indication-specific relevant criteria such as concomitant therapy requirements (not including prerequisite therapy), performance status, etc. identified in section III; **AND**
- Duration of authorization has not been exceeded (*refer to Section I*); **AND**
- Disease response with treatment as defined by stabilization of disease or decrease in size of tumor or tumor spread, unless otherwise specified in section III; **AND**
- Absence of unacceptable toxicity from the drug. Examples of unacceptable toxicity include: immune-mediated adverse reactions (e.g., pneumonitis, hepatitis, colitis, endocrinopathies, nephritis/renal dysfunction, rash/dermatitis [including Stevens-Johnson syndrome (SJS), drug rash with eosinophilia and systemic symptoms (DRESS), and toxic epidermal necrolysis (TEN)], myocarditis, pericarditis, vasculitis, solid organ transplant rejection, etc.), severe infusion-related reactions, complications of allogeneic hematopoietic stem cell transplantation (HSCT), etc.

^Δ Notes:

- Members responding to therapy who relapse ≥ 6 months after discontinuation due to duration) are eligible to re-initiate PD-directed therapy.
- Members previously presenting with aggressive disease who are exhibiting stable disease on treatment as their best response (or if therapy improved performance status) may be eligible for continued therapy without interruption or discontinuation.
- Members who complete adjuvant therapy and progress ≥ 6 months after discontinuation are eligible to re-initiate PD-directed therapy for metastatic disease.
- Members whose tumors, upon re-biopsy, demonstrate a change in actionable mutation (e.g., MSS initial biopsy; MSI-H subsequent biopsy) may be eligible to re-initiate PD-directed therapy and will be evaluated on a case-by-case basis.

V. Dosage/Administration ^{Δ 1,14,27,28,36-45}

Indication	Dose
------------	------

NSCLC, SCLC, Cervical Cancer, Rectal Cancer, Urothelial Carcinoma (Bladder Cancer)	Administer intravenously until disease progression or unacceptable toxicity*: – 840 mg every 2 weeks or – 1200 mg every 3 weeks or – 1680 mg every 4 weeks <i>*NSCLC and Urothelial Carcinoma (Bladder Cancer) adjuvant treatment may continue up to a maximum of 12 months in members without disease recurrence or unacceptable toxicity.</i>
HCC	Administer intravenously until disease progression or unacceptable toxicity: <u>First-line therapy:</u> – 840 mg every 2 weeks or – 1200 mg every 3 weeks or – 1680 mg every 4 weeks <u>Subsequent therapy:</u> – 1200 mg every 3 weeks
Cutaneous Melanoma	Administer intravenously until disease progression or unacceptable toxicity: – 840 mg every 2 weeks or – 1200 mg every 3 weeks or – 1680 mg every 4 weeks <i>*Prior to initiating atezolizumab, members should receive a 28-day treatment cycle of cobimetinib 60 mg orally once daily (21 days on and 7 days off) and vemurafenib 960 mg orally twice daily from Days 1-21 and vemurafenib 720 mg orally twice daily from Days 22-28.</i>
Mesotheliomas (peritoneal, pericardial, and tunica vaginalis testis)	Administer 1200 mg every 3 weeks intravenously until disease progression or unacceptable toxicity
ASPS	Administer intravenously until disease progression or unacceptable toxicity: <u>Adult members:</u> – 840 mg every 2 weeks or – 1200 mg every 3 weeks or – 1680 mg every 4 weeks <u>Pediatric members at least 2 years of age:</u> – 15 mg/kg (up to a maximum 1200 mg) every 3 weeks
Colon Cancer	Adjuvant Therapy: In combination with FOLFOX followed by single agent maintenance: – Administer 840 mg intravenously every 2 weeks for 12 cycles (6 months) then begin maintenance 840 mg intravenously every 2 weeks for 13 cycles (12 months total) In combination with CAPEOX followed by single agent maintenance: – Administer 1200 mg intravenously every 3 weeks for 8 cycles (6 months) then begin maintenance 1200 mg intravenously every 3 weeks for 8 cycles (12 months total)

	<u>Single Agent Therapy for locally unresectable, medically inoperable, advanced or metastatic disease:</u> Administer intravenously until disease progression or unacceptable toxicity: – 840 mg every 2 weeks or – 1200 mg every 3 weeks or – 1680 mg every 4 weeks
Small Bowel Adenocarcinoma	<u>Adjuvant Therapy:</u> In combination with FOLFOX followed by single agent maintenance: – Administer 840 mg intravenously every 2 weeks for 12 cycles (6 months) then begin maintenance 840 mg intravenously every 2 weeks for 13 cycles (12 months total) In combination with CAPEOX followed by single agent maintenance: – Administer 1200 mg intravenously every 3 weeks for 8 cycles (6 months) then begin maintenance 1200 mg intravenously every 3 weeks for 8 cycles (12 months total) <u>Single Agent Therapy for locally unresectable, medically inoperable, advanced or metastatic disease:</u> Administer intravenously until disease progression or unacceptable toxicity: – 840 mg every 2 weeks or – 1200 mg every 3 weeks or – 1680 mg every 4 weeks
Thymic Carcinoma	Administer intravenously for up to 24 months in absence of disease progression or unacceptable toxicity: – Administer 1200 mg every 3 weeks
CLL/SLL	Administer 1200 mg intravenously every 3 weeks for up to a total of 18 cycles. <i>Note: Administer atezolizumab on day 2 of cycle 1 (first infusion) and on day 1 of each subsequent 21-day cycle</i>
<u>Dosing should be calculated using actual body weight and not flat dosing (as applicable) based on the following:</u> ³⁰⁻³⁵ • 840 mg (15 mg/kg) in members receiving therapy every 21 days who weigh ≤ 61 kg • 1200 mg (20 mg/kg) in member receiving therapy every 28 days who weigh ≤ 66 kg <i>Note: This information is not meant to replace clinical decision making when initiating or modifying medication therapy and should only be used as a guide. Member-specific variables should be taken into account.</i>	

VI. Billing Code/Availability Information

HCPSC Code:

- J9022 – Injection, atezolizumab, 10 mg; 1 billable unit = 10 mg

NDC(s):

- Tecentriq 1200 mg/20 mL solution for injection single-dose vial: 50242-0917-xx
- Tecentriq 840 mg/14 mL solution for injection single-dose vial: 50242-0918-xx

VII. References (STANDARD)

1. Tecentriq [package insert]. South San Francisco, CA; Genentech, Inc; May 2026. Accessed June 2026.
2. Ventana Product Library, Roche Pharmaceuticals. VENTANA PD-L1 [SP142] Assay. <https://diagnostics.roche.com/global/en/products/lab/pd-l1-sp142-assay-ventana-rtd001231.html#productInfo> and product label https://www.accessdata.fda.gov/cdrh_docs/pdf16/P160006C.pdf.
3. U.S. Food and Drug Administrations (FDA). Division of Drug Information. Health Alert. FDA Alerts Health Care Professionals and Oncology Clinical Investigators about an Efficacy Issue Identified in Clinical Trials for Some Patients Taking Keytruda (pembrolizumab) or Tecentriq (atezolizumab) as Monotherapy to Treat Urothelial Cancer with Low Expression of PD-L1 | FDA. August 2018.
4. Balar AV, Galsky MD, Rosenberg JE, et al. Atezolizumab as first-line therapy in cisplatin-ineligible patients with locally advanced and metastatic urothelial carcinoma: a single-arm, multicentre, phase 2 trial. *Lancet*. 2017 January 07; 389(10064): 67–76. doi:10.1016/S0140-6736(16)32455-2.
5. Socinski MA, Jotte RM, Cappuzzo F, et. al. Atezolizumab for First-Line Treatment of Metastatic Nonsquamous NSCLC. *N Engl J Med* 2018; 378:2288-2301. DOI: 10.1056/NEJMoa1716948.
6. Referenced with permission from the NCCN Drugs and Biologics Compendium (NCCN Compendium®) atezolizumab. National Comprehensive Cancer Network, 2026. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed June 2026.
7. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Bladder Cancer. Version 2.2026. National Comprehensive Cancer Network, 2026. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed June 2026.

8. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Non-Small Cell Lung Cancer. Version 6.2026. National Comprehensive Cancer Network, 2026. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed June 2026.
9. Gupta S, Sonpavde G, Grivas P, et al. Defining “platinum-ineligible” patients with metastatic urothelial cancer (mUC). *J Clin Oncol*. 2019 Mar 1;37(7_suppl):451.
10. Rosenberg JE, Hoffman-Censits J, Powles T, et al. Atezolizumab in patients with locally advanced and metastatic urothelial carcinoma who have progressed following treatment with platinum-based chemotherapy: a single-arm, multicentre, phase 2 trial. *Lancet*. 2016 May 7;387(10031):1909-20. doi: 10.1016/S0140-6736(16)00561-4. Epub 2016 Mar 4.
11. West H, McCleod M, Hussein M, et al. Atezolizumab in combination with carboplatin plus nab-paclitaxel chemotherapy compared with chemotherapy alone as first-line treatment for metastatic non-squamous non-small-cell lung cancer (IMpower130): a multicentre, randomised, open-label, phase 3 trial. *Lancet Oncol*. 2019 Jul;20(7):924-937. doi: 10.1016/S1470-2045(19)30167-6. Epub 2019 May 20.
12. Rittmeyer A, Barlesi F, Waterkamp D, et al. Atezolizumab versus docetaxel in patients with previously treated non-small-cell lung cancer (OAK): a phase 3, open-label, multicentre randomised controlled trial. *Lancet*. 2017 Jan 21;389(10066):255-265. doi: 10.1016/S0140-6736(16)32517-X. Epub 2016 Dec 13.
13. Schmid P, Adams S, Rugo HS, et al. Atezolizumab and Nab-Paclitaxel in Advanced Triple-Negative Breast Cancer. *N Engl J Med*. 2018 Nov 29;379(22):2108-2121. doi: 10.1056/NEJMoa1809615. Epub 2018 Oct 20.
14. Horn L, Mansfield AS, Szczesna A, et al. First-Line Atezolizumab plus Chemotherapy in Extensive-Stage Small-Cell Lung Cancer. *N Engl J Med*. 2018 Dec 6;379(23):2220-2229. doi: 10.1056/NEJMoa1809064. Epub 2018 Sep 25.
15. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Hepatocellular Carcinoma. Version 1.2026. National Comprehensive Cancer Network, 2026. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed June 2026.
16. Pishvaian MJ, Lee MS, Ryoo B, et al. Updated safety and clinical activity results from a Phase Ib study of atezolizumab + bevacizumab in hepatocellular carcinoma (HCC). ESMO 2018 Congress. Munich, Germany; 2018.

17. De Marinis F, Jassem J, Spigel DR, et al. 480TiP IMpower110: Phase III study on 1L atezolizumab (atezo) in PD-L1–selected chemotherapy (chemo)-naive NSCLC patients (pts). *Annals of Oncology*. 2016 Dec 1;27(suppl_9).
18. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Small Cell Lung Cancer. Version 2.2026. National Comprehensive Cancer Network, 2026. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed June 2026.
19. Gutzmer R, Stroyakovskiy D, Gogas H, et al. Atezolizumab, vemurafenib, and cobimetinib as first-line treatment for unresectable advanced BRAFV600 mutation-positive melanoma (IMspire150): primary analysis of the randomised, double-blind, placebo-controlled, phase 3 trial. *Lancet*. 2020;395(10240):1835-1844. doi:10.1016/S0140-6736(20)30934-X.
20. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Melanoma: Cutaneous. Version 2.2026. National Comprehensive Cancer Network, 2026. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed June 2026.
21. Finn RS, Qin S, Ikeda M, et al; IMbrave150 Investigators. Atezolizumab plus Bevacizumab in Unresectable Hepatocellular Carcinoma. *N Engl J Med*. 2020 May 14;382(20):1894-1905.
22. Bellmunt, J, Valderrama BP (2026). Treatment of metastatic urothelial cancer of the bladder and urinary tract. In Lerner SP, Shah SM (Eds.), *UpToDate*. Last updated Feb 05, 2026. Accessed June 2026. Available from <https://www.uptodate.com/contents/treatment-of-metastatic-urothelial-carcinoma-of-the-bladder-and-urinary-tract>.
23. Felip E, Altorki N, Zhou C, et al. Adjuvant atezolizumab after adjuvant chemotherapy in resected stage IB-IIIA non-small-cell lung cancer (IMpower010): a randomised, multicentre, open-label, phase 3 trial. *Lancet*. 2021 Oct 9;398(10308):1344-1357. doi: 10.1016/S0140-6736(21)02098-5. Epub 2021 Sep 20.
24. Raghav KPS, Overman MJ, Liu S, et al. A phase II trial of atezolizumab and bevacizumab in patients with relapsed/refractory and unresectable malignant peritoneal mesothelioma. *Journal of Clinical Oncology* 2020 38:15_suppl, 9013-9013.
25. De Santis M, Bellmunt J, Mead G, et al. Randomized phase II/III trial assessing gemcitabine/ carboplatin and methotrexate/carboplatin/vinblastine in patients with advanced urothelial cancer "unfit" for cisplatin-based chemotherapy: phase II--results of EORTC study 30986. *J Clin Oncol*. 2009 Nov 20;27(33):5634-9. doi: 10.1200/JCO.2008.21.4924. Epub 2009 Sep 28.

26. Naqash AR, O'Sullivan Coyne GH, Moore N, et al. Phase II study of atezolizumab in advanced alveolar soft part sarcoma (ASPS). *Journal of Clinical Oncology* 2021 39:15_suppl, 11519-11519.
27. Raghav K, Liu S, Overman MJ, et al. Efficacy, Safety, and Biomarker Analysis of Combined PD-L1 (Atezolizumab) and VEGF (Bevacizumab) Blockade in Advanced Malignant Peritoneal Mesothelioma. *Cancer Discov.* 2021 Nov;11(11):2738-2747. doi: 10.1158/2159-8290.CD-21-0331.
28. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Atezolizumab + Bevacizumab: Hepatocellular Carcinoma Order Template, HEP32. National Comprehensive Cancer Network, 2026. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed June 2026.
29. Ascierto PA, Stroyakovskiy D, Gogas H, et al. Overall survival with first-line atezolizumab in combination with vemurafenib and cobimetinib in BRAFV600 mutation-positive advanced melanoma (IMspire150): second interim analysis of a multicentre, randomised, phase 3 study. *Lancet Oncol* 2023;24:33-44.
30. Kicken, M.P., Deenen, M.J., Moes, D.J.A.R. *et al.* An Evidence-Based Rationale for Dose De-escalation of Subcutaneous Atezolizumab. *Targ Oncol* 19, 779–787 (2024).
<https://doi.org/10.1007/s11523-024-01087-4>.
31. Morrissey KM, Marchand M, Patel H, *et al.* Alternative dosing regimens for atezolizumab: an example of model-informed drug development in the postmarketing setting. *Cancer Chemother Pharmacol.* 2019 Dec;84(6):1257-1267. doi: 10.1007/s00280-019-03954-8. Epub 2019 Sep 21. PMID: 31542806; PMCID: PMC6820606.
32. Liu SN, Marchand M, Liu X, et al. Extension of the Alternative Intravenous Dosing Regimens of Atezolizumab into Combination Settings through Modeling and Simulation. *The Journal of Clinical Pharmacology*, 10.1002, 62,11, (1393-1402), Nov. 2022.
33. Cody J. Peer, Keith T. Schmidt, Oluwatobi Arisa, William J. Richardson, Koosha Paydary, Daniel A. Goldstein, James L. Gulley, William D. Figg, Mark J. Ratain, In Silico Re-Optimization of Atezolizumab Dosing Using Population Pharmacokinetic Simulation and Exposure–Response Simulation, *The Journal of Clinical Pharmacology*, 10.1002/jcph.2203, 63, 6, (672-680), (2023).
34. Wesevich A, Goldstein DA, Paydary K, et al. Interventional pharmacoeconomics for immune checkpoint inhibitors through alternative dosing strategies. *British Journal of Cancer.* 2023 Oct;129(9):1389-1396. DOI: 10.1038/s41416-023-02367-y. PMID: 37542109; PMCID: PMC10628132.

35. Maritaz, C., Broutin, S., Chaput, N. *et al.* Immune checkpoint-targeted antibodies: a room for dose and schedule optimization?. *J Hematol Oncol* 15, 6 (2022).
<https://doi.org/10.1186/s13045-021-01182-3>
36. Oaknin A, Gladieff L, Martínez-García J, et al. ENGOT-Cx10–GEICO 68-C–JGOG1084–GOG-3030 Investigators. Atezolizumab plus bevacizumab and chemotherapy for metastatic, persistent, or recurrent cervical cancer (BEATcc): a randomised, open-label, phase 3 trial. *Lancet* 2024;403:31-43.
37. Sinicrope FA, Ou FS, Arnold D, et al. Randomized trial of standard chemotherapy alone or combined with atezolizumab as adjuvant therapy for patients with stage III deficient DNA mismatch repair colon cancer (Alliance A021502; ATOMIC). [abstract]. *J Clin Oncol* 2025; 43:LBA1
38. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Colon Cancer. Version 2.2026. National Comprehensive Cancer Network, 2026. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed June 2026.
39. Paz-Ares L, Borghaei H, Liu SV, et al. Efficacy and safety of first-line maintenance therapy with lurbinectedin plus atezolizumab in extensive-stage small-cell lung cancer (IMforte): a randomised, multicentre, open-label, phase 3 trial. *Lancet*. 2025;405(10495):2129-2143. doi:10.1016/S0140-6736(25)01011-6.
40. Shukuya T, Asao T, Goto Y, et al. Activity and safety of atezolizumab plus carboplatin and paclitaxel in patients with advanced or recurrent thymic carcinoma (MARBLE): a multicentre, single-arm, phase 2 trial. *Lancet Oncol* 2025;26:331-342.
41. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Thymomas and Thymic Carcinomas. Version 2.2026. National Comprehensive Cancer Network, 2026. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed June 2026.
42. Tedeschi A, Frustaci AM, Condoluci A, et al. Atezolizumab, venetoclax, and obinutuzumab combination in Richter transformation diffuse large B-cell lymphoma (MOLTO): a multicentre, single-arm, phase 2 trial. *Lancet Oncol* 2024;25:1298-1309.
43. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Chronic Lymphocytic Leukemia/Small Lymphocytic Lymphoma. Version 2.2026. National Comprehensive Cancer Network, 2026. The NCCN Compendium® is a derivative work

of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed June 2026.

44. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Small Bowel Adenocarcinoma. Version 2.2026. National Comprehensive Cancer Network, 2026. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed June 2026.
45. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Rectal Cancer. Version 2.2026. National Comprehensive Cancer Network, 2026. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed June 2026.
46. Powles T, Kann AG, Castellano D, et al; IMvigor011 Investigators. ctDNA-Guided Adjuvant Atezolizumab in Muscle-Invasive Bladder Cancer. *N Engl J Med*. 2025 Dec 18;393(24):2395-2408. doi: 10.1056/NEJMoa2511885. Epub 2025 Oct 20. PMID: 41124204.

VIII. References (ENHANCED)

- 1e. Balar AV, Castellano D, O'Donnell PH, et al. Pembrolizumab as first-line therapy in cisplatin-ineligible advanced urothelial cancer: Results from the total KEYNOTE-052 study population. *J Clin Oncol* 2017 Feb;35(6_suppl):284.
- 2e. De Santis M, Bellmunt J, Mead G, et al. Randomized phase II/III trial assessing gemcitabine/carboplatin and methotrexate/carboplatin/vinblastine in patients with advanced urothelial cancer who are unfit for cisplatin-based chemotherapy: EORTC study 30986. *J Clin Oncol*. 2012;30(2):191–199. doi:10.1200/JCO.2011.37.3571.
- 3e. Powles T, Durán I, van der Heijden MS, et al. Atezolizumab versus chemotherapy in patients with platinum-treated locally advanced or metastatic urothelial carcinoma (IMvigor211): a multicentre, open-label, phase 3 randomised controlled trial. *Lancet*. 2018 Feb 24;391(10122):748-757. doi: 10.1016/S0140-6736(17)33297-X.
- 4e. Bellmunt J, de Wit R, Vaughn DJ, et al. Pembrolizumab as Second-Line Therapy for Advanced Urothelial Carcinoma. *N Engl J Med*. 2017;376(11):1015–1026. doi:10.1056/NEJMoa1613683.

- 5e. Sharma P, Retz M, Siefker-Radtke A, et al. Nivolumab in metastatic urothelial carcinoma after platinum therapy (CheckMate 275): a multicentre, single-arm, phase 2 trial. *Lancet Oncol*. 2017 Mar;18(3):312-322. doi: 10.1016/S1470-2045(17)30065-7.
- 6e. Massard C, Gordon MS, Sharma S, et al. Safety and Efficacy of Durvalumab (MEDI4736), an Anti-Programmed Cell Death Ligand-1 Immune Checkpoint Inhibitor, in Patients With Advanced Urothelial Bladder Cancer. *J Clin Oncol*. 2016;34(26):3119–3125. doi:10.1200/JCO.2016.67.9761.
- 7e. Patel MR, Ellerton J, Infante JR, et al. Avelumab in metastatic urothelial carcinoma after platinum failure (JAVELIN Solid Tumor): pooled results from two expansion cohorts of an open-label, phase 1 trial. *Lancet Oncol*. 2018 Jan;19(1):51-64. doi: 10.1016/S1470-2045(17)30900-2.
- 8e. Siefker-Radtke AO, Necchi A, Park SH, et al. First results from the primary analysis population of the phase 2 study of erdafitinib (ERDA; JNJ-42756493) in patients (pts) with metastatic or unresectable urothelial carcinoma (mUC) and FGFR alterations (FGFRalt). *J Clin Oncol* 2018;36(15_suppl):4503.
- 9e. Reck M, Rodríguez-Abreu D, Robinson AG, et al. Pembrolizumab versus Chemotherapy for PD-L1–Positive Non–Small-Cell Lung Cancer. *N Engl J Med* 2016; 375:1823-1833.
- 10e. Gandhi L, Rodríguez-Abreu D, Gadgeel S, et al. Pembrolizumab plus Chemotherapy in Metastatic Non–Small-Cell Lung Cancer. *N Engl J Med* 2018; 378:2078-2092.
- 11e. Barlesi F, Park K, Ciardiello F, et al. Primary analysis from OAK, a randomized phase III study comparing atezolizumab with docetaxel in 2L/3L NSCLC. *Ann of Oncol* 2016 Oct;27(suppl_6):LBA44_PR.
- 12e. Borghaei H, Paz-Ares L, Horn L, et al. Nivolumab versus Docetaxel in Advanced Nonsquamous Non-Small-Cell Lung Cancer. *N Engl J Med*. 2015;373(17):1627–1639. doi:10.1056/NEJMoa1507643.
- 13e. Herbst RS, Baas P, Kim DW, et al. Pembrolizumab versus docetaxel for previously treated, PD-L1-positive, advanced non-small-cell lung cancer (KEYNOTE-010): a randomised controlled trial. *Lancet*. 2016 Apr 9;387(10027):1540-50. doi: 10.1016/S0140-6736(15)01281-7.
- 14e. Spigel D et al. IMpower110: Interim OS Analysis of a Phase III Study of Atezolizumab (atezo) vs Platinum-Based Chemotherapy (chemo) as 1L Treatment (tx) in PD-L1–selected NSCLC [ESMO 2019 Abstract LBA78].
- 15e. Paz-Ares L, Dvorkin M, Chen Y, et al. Durvalumab plus platinum-etoposide versus platinum-etoposide in first-line treatment of extensive-stage small-cell lung cancer (CASPIAN): a randomised, controlled, open-label, phase 3 trial. *Lancet*. 2019;394(10212):1929-1939. doi:10.1016/S0140-6736(19)32222-6.
- 16e. Zalcman G, Peters S, Mansfield AS, et al. Checkmate 743: A phase 3, randomized, open-label trial of nivolumab (nivo) plus ipilimumab (ipi) vs pemetrexed plus cisplatin or carboplatin as first-

line therapy in unresectable pleural mesothelioma. *Journal of Clinical Oncology* 2017 35:15_suppl, TPS8581-TPS8581.

- 17e. West H, McCleod M, Hussein M, et al. Atezolizumab in combination with carboplatin plus nab-paclitaxel chemotherapy compared with chemotherapy alone as first-line treatment for metastatic non-squamous non-small-cell lung cancer (IMpower130): a multicentre, randomised, open-label, phase 3 trial. *Lancet Oncol.* 2019 Jul;20(7):924-937. doi: 10.1016/S1470-2045(19)30167-6. Epub 2019 May 20.
- 18e. Powles T, Park SH, Voog E, et al. Maintenance avelumab + best supportive care (BSC) versus BSC alone after platinum-based first-line (1L) chemotherapy in advanced urothelial carcinoma (UC): JAVELIN Bladder 100 phase III interim analysis. *J Clin Oncol.* 2020;38(18_suppl):LBA1-LBA1.
- 19e. Cortes J, Cescon DW, Rugo HS, et al. KEYNOTE-355: Randomized, double-blind, phase III study of pembrolizumab + chemotherapy versus placebo + chemotherapy for previously untreated locally recurrent inoperable or metastatic triple-negative breast cancer. *Journal of Clinical Oncology* 38, no. 15_suppl(May 20, 2020)1000-1000.
- 20e. Sezer A, Kilickap S, Gümüş M, et al. Cemiplimab monotherapy for first-line treatment of advanced non-small-cell lung cancer with PD-L1 of at least 50%: a multicentre, open-label, global, phase 3, randomised, controlled trial. *Lancet.* 2021 Feb 13;397(10274):592-604.
- 21e. Miles D, Gligorov J, André F, et al. Primary results from IMpassion131, a double-blind, placebo-controlled, randomised phase III trial of first-line paclitaxel with or without atezolizumab for unresectable locally advanced/metastatic triple-negative breast cancer. *Ann Oncol.* 2021 Aug;32(8):994-1004. doi: 10.1016/j.annonc.2021.05.801. Epub 2021 Jul 1.
- 22e. Fennell DA, Ewings S, Ottensmeier C, et al. Nivolumab versus placebo in patients with relapsed malignant mesothelioma (CONFIRM): a multicentre, double-blind, randomised, phase 3 trial. *Lancet Oncol* 2021; 22:1530.
- 23e. Galsky MD, Ariba J, Bamias A, et al; IMvigor130 Study Group. Atezolizumab with or without chemotherapy in metastatic urothelial cancer (IMvigor130): a multicentre, randomised, placebo-controlled phase 3 trial. *Lancet.* 2020 May 16;395(10236):1547-1557. doi: 10.1016/S0140-6736(20)30230-0. PMID: 32416780.
- 24e. Johnson ML, Cho BC, Luft A, et al; POSEIDON investigators. Durvalumab With or Without Tremelimumab in Combination With Chemotherapy as First-Line Therapy for Metastatic Non-Small-Cell Lung Cancer: The Phase III POSEIDON Study. *J Clin Oncol.* 2022 Nov 3;JCO2200975. doi: 10.1200/JCO.22.00975.
- 25e. Gogishvili M, Melkadze T, Makharadze T, et al. LBA51 EMPOWER-Lung 3: Cemiplimab in combination with platinum doublet chemotherapy for first-line (1L) treatment of advanced non-small cell lung cancer (NSCLC). *Annals of Oncology*, ISSN: 0923-7534, Vol: 32, SUPPLEMENT 5, S1328, SEPTEMBER 01, 2021. DOI10.1016/j.annonc.2021.08.2130.

- 26e. Wilky BA, Trucco MM, Subhawong TK, et al. Axitinib plus pembrolizumab in patients with advanced sarcomas including alveolar soft-part sarcoma: a single-centre, single-arm, phase 2 trial. *Lancet Oncol*. 2019 Jun;20(6):837-848. doi: 10.1016/S1470-2045(19)30153-6. Epub 2019 May 8.
- 27e. D'Alessio A, Fulgenzi CAM, Nishida N, et al. Preliminary evidence of safety and tolerability of atezolizumab plus bevacizumab in patients with hepatocellular carcinoma and Child-Pugh A and B cirrhosis: A real-world study. *Hepatology*. 2022 Oct;76(4):1000-1012. doi: 10.1002/hep.32468.
- 28e. Stacchiotti S, Negri T, Zaffaroni N, et al. Sunitinib in advanced alveolar soft part sarcoma: evidence of a direct antitumor effect. *Ann Oncol*. 2011 Jul;22(7):1682-1690. doi: 10.1093/annonc/mdq644. Epub 2011 Jan 17.
- 29e. Stacchiotti S, Mir O, Le Cesne A, et al. Activity of Pazopanib and Trabectedin in Advanced Alveolar Soft Part Sarcoma. *Oncologist*. 2018 Jan;23(1):62-70. doi: 10.1634/theoncologist.2017-0161. Epub 2017 Jul 28.
- 30e. Kudo M. Durvalumab plus tremelimumab in unresectable hepatocellular carcinoma. *Hepatobiliary Surgery and Nutrition*. 2022;11(4):592-596. doi:https://doi.org/10.21037/hbsn-22-143
- 31e. Ascierto PA, Dréno B, Larkin J, et al. 5-Year Outcomes with Cobimetinib plus Vemurafenib in BRAFV600 Mutation–Positive Advanced Melanoma: Extended Follow-up of the coBRIM Study. *Clinical Cancer Research*. 2021;27(19):5225-5235. doi:https://doi.org/10.1158/1078-0432.ccr-21-0809
- 32e. Robert C, Karaszewska B, Schachter J, et al. Improved Overall Survival in Melanoma with Combined Dabrafenib and Trametinib. *New England Journal of Medicine*. 2015;372(1):30-39. doi:https://doi.org/10.1056/nejmoa1412690
- 33e. Dummer R, Ascierto PA, Gogas HJ, et al. Encorafenib plus binimetinib versus vemurafenib or encorafenib in patients with BRAF-mutant melanoma (COLUMBUS): a multicentre, open-label, randomised phase 3 trial. *The Lancet Oncology*. 2018;19(5):603-615. doi:https://doi.org/10.1016/S1470-2045(18)30142-6
- 34e. Ferrucci PF, Giacomo AMD, Vecchio MD, et al. KEYNOTE-022 part 3: a randomized, double-blind, phase 2 study of pembrolizumab, dabrafenib, and trametinib in BRAF-mutant melanoma. *Journal for ImmunoTherapy of Cancer*. 2020;8(2):e001806. doi:https://doi.org/10.1136/jitc-2020-001806
- 35e. Frumovitz M, Munsell MF, Burzawa JK, et al. Combination therapy with topotecan, paclitaxel, and bevacizumab improves progression-free survival in recurrent small cell neuroendocrine carcinoma of the cervix. *Gynecol Oncol*. 2017 Jan;144(1):46-50.
- 36e. Tempfer CB, Tischoff I, Dogan A, et al. Neuroendocrine carcinoma of the cervix: a systematic review of the literature. *BMC Cancer* 2018;18:530.

- 37e. Colombo N, Dubot C, Lorusso D, et al. Pembrolizumab for Persistent, Recurrent, or Metastatic Cervical Cancer. *New England Journal of Medicine*. 2021;385(20). doi:<https://doi.org/10.1056/nejmoa2112435>
- 38e. Lee SM, Schulz C, Prabhaskar K, et al. First-line atezolizumab monotherapy versus single-agent chemotherapy in patients with non-small-cell lung cancer ineligible for treatment with a platinum-containing regimen (IPSOS): a phase 3, global, multicentre, open-label, randomised controlled study. *Lancet*. 2023 Aug 5;402(10400):451-463. doi: 10.1016/S0140-6736(23)00774-2.
- 39e. Terashima T, Kido H, Takata N, et al. Phase II Study of Atezolizumab and Bevacizumab Combination Therapy for Patients with Advanced Hepatocellular Carcinoma Previously Treated with Lenvatinib. *Cancers*. 2025;17(2):278-278. doi:<https://doi.org/10.3390/cancers17020278>
- 40e. Mori N, Tamaki N, Takaki S, et al. Treatment response to durvalumab plus tremelimumab after progression with previous immune checkpoint inhibitor in unresectable hepatocellular carcinoma. *Investigational New Drugs*. 2024;42(5):559-565. doi:<https://doi.org/10.1007/s10637-024-01470-y>
- 41e. Kelley RK, Sangro B, Harris W, et al. Safety, Efficacy, and Pharmacodynamics of Tremelimumab Plus Durvalumab for Patients With Unresectable Hepatocellular Carcinoma: Randomized Expansion of a Phase I/II Study. *J Clin Oncol*. 2021 Sep 20;39(27):2991-3001.
- 42e. Lemma GL, Lee JW, Aisner SC et al. Phase II study of carboplatin and paclitaxel in advanced thymoma and thymic carcinoma. *J Clin Oncol*. 2011 May 20;29(15):2060-5. doi: 10.1200/JCO.2010.32.9607. Epub 2011 Apr 18. PMID: 21502559; PMCID: PMC3107762.
- 43e. Hirai F, Yamanaka T, Taguchi K, Daga H, et al. A multicenter phase II study of carboplatin and paclitaxel for advanced thymic carcinoma: WJOG4207L. *Ann Oncol*. 2015 Feb;26(2):363-8. doi: 10.1093/annonc/mdu541. Epub 2014 Nov 17. PMID: 25403584.
- 44e. Loehrer PJ Sr, Kim K, Aisner SC, et al. Cisplatin plus doxorubicin plus cyclophosphamide in metastatic or recurrent thymoma: final results of an intergroup trial. The Eastern Cooperative Oncology Group, Southwest Oncology Group, and Southeastern Cancer Study Group. *J Clin Oncol*. 1994 Jun;12(6):1164-8. doi: 10.1200/JCO.1994.12.6.1164. PMID: 8201378.
- 45e. Kim ES, Putnam JB, Komaki R, et al. Phase II study of a multidisciplinary approach with induction chemotherapy, followed by surgical resection, radiation therapy, and consolidation chemotherapy for unresectable malignant thymomas: final report. *Lung Cancer*. 2004 Jun;44(3):369-79. doi: 10.1016/j.lungcan.2003.12.010. PMID: 15140551.
- 46e. Fornasiero A, Daniele O, Ghiotto C, Piazza M, et al. Chemotherapy for invasive thymoma. A 13-year experience. *Cancer*. 1991 Jul 1;68(1):30-3. doi: 10.1002/1097-0142(19910701)68:1<30::aid-cnrc2820680106>3.0.co;2-4. PMID: 2049749.
- 47e. Giaccone G, Ardizzoni A, Kirkpatrick et al. Cisplatin and etoposide combination chemotherapy for locally advanced or metastatic thymoma. A phase II study of the European Organization for

- Research and Treatment of Cancer Lung Cancer Cooperative Group. *J Clin Oncol*. 1996 Mar;14(3):814-20. doi: 10.1200/JCO.1996.14.3.814. PMID: 8622029.
- 48e. Loehrer PJ Sr, Jiroutek M, et al. Combined etoposide, ifosfamide, and cisplatin in the treatment of patients with advanced thymoma and thymic carcinoma: an intergroup trial. *Cancer*. 2001 Jun 1;91(11):2010-5. PMID: 11391579.
- 49e. Özgüroğlu M, Kilickap S, et al. First-line cemiplimab monotherapy and continued cemiplimab beyond progression plus chemotherapy for advanced non-small-cell lung cancer with PD-L1 50% or more (EMPOWER-Lung 1): 35-month follow-up from a multicentre, open-label, randomised, phase 3 trial. *Lancet Oncol*. 2023 Sep;24(9):989-1001. doi: 10.1016/S1470-2045(23)00329-7. Epub 2023 Aug 14. Erratum in: *Lancet Oncol*. 2023 Oct;24(10):e405. doi: 10.1016/S1470-2045(23)00463-1. PMID: 37591293.
- 50e. Rocha Lima, Yothers G, George TJ, et al. Colorectal cancer metastatic dMMR immunotherapy (COMMIT) study: a randomized phase III study of atezolizumab (atezo) monotherapy versus mFOLFOX6/bevacizumab/atezo (FFX/bev) in the first-line treatment of patients (pts) with deficient DNA mismatch repair (dMMR) or microsatellite instability-high (MSI-H) metastatic colorectal cancer (mCRC)—NRG-GI004/SWOG-S1610. *J Clin Oncol*. 2026;44(Suppl 2):14. doi:10.1200/JCO.2026.44.2_suppl.14
- 51e. Friedman CF, Hainsworth JD, Kurzrock R, et al. Atezolizumab Treatment of Tumors with High Tumor Mutational Burden from MyPathway, a Multicenter, Open-Label, Phase IIa Multiple Basket Study. *Cancer Discov*. 2022 Mar 8;12(3):654–669. doi: 10.1158/2159-8290.CD-21-0450
- 52e. Oh CR, Kim JE, Hong YS, et al. Phase II study of durvalumab monotherapy in patients with previously treated microsatellite instability-high/mismatch repair-deficient or POLE-mutated metastatic or unresectable colorectal cancer. *Int J Cancer*. 2022 Jun 15;150(12):2038-2045. doi: 10.1002/ijc.33966. Epub 2022 Mar 1.
- 53e. Thomas DM, Kim, JE, Barlesi, et al. TAPISTRY: A Phase II Study of Atezolizumab in Patients with Tumor Mutational Burden-High Tumors. *Clin Cancer Res*. 2026 Mar 16;32(6):1078-1086. doi: 10.1158/1078-0432.CCR-25-3336
- 54e. Jassem J, Ramlau R, Santoro A, et al, Phase III Trial of Pemetrexed Plus Best Supportive Care Compared With Best Supportive Care in Previously Treated Patients With Advanced Malignant Pleural Mesothelioma. *J Clin Oncol*, 2008, 26(10):1698-704. [PubMed 18375898]
- 55e. Rocha Lima CMSP, Yothers G, George TJ, et al. Colorectal Cancer Metastatic dMMR Immunotherapy (COMMIT) study: A randomized phase III study of atezolizumab (atezo) monotherapy versus mFOLFOX6/bevacizumab/atezo (FFX/bev) in the first-line treatment of patients (pts) with deficient DNA mismatch repair (dMMR) or microsatellite instability-high (MSI-H) metastatic colorectal cancer (mCRC)—NRG-GI004/SWOG-S1610. *J Clin Oncol*. 2026;44(2_suppl):14. doi:10.1200/JCO.2026.44.2_suppl.14.
- 56e. Prime Therapeutics Management. Tecentriq IV/SQ Clinical Literature Review Analysis. Last updated June 2026. Accessed June 2026.

Appendix A – Non-Quantitative Treatment Limitations (NQTL) Factor Checklist

Non-quantitative treatment limitations (NQTLs) refer to the methods, guidelines, standards of evidence, or other conditions that can restrict how long or to what extent benefits are provided under a health plan. These may include things like utilization review or prior authorization. The utilization management NQTL applies comparably, and not more stringently, to mental health/substance use disorder (MH/SUD) Medical Benefit Prescription Drugs and medical/surgical (M/S) Medical Benefit Prescription Drugs. The table below lists the factors that were considered in designing and applying prior authorization to this drug/drug group, and a summary of the conclusions that Prime's assessment led to for each.

Factor	Conclusion
Indication	Yes: Consider for PA
Safety and efficacy	No: PA not a priority
Potential for misuse/abuse	No: PA not a priority
Cost of drug	Yes: Consider for PA

Appendix 1 – Covered Diagnosis Codes

ICD-10	ICD-10 Description
C17.0	Malignant neoplasm of duodenum
C17.1	Malignant neoplasm of jejunum
C17.2	Malignant neoplasm of ileum
C17.3	Meckel's diverticulum, malignant
C17.8	Malignant neoplasm of overlapping sites of small intestine
C17.9	Malignant neoplasm of small intestine, unspecified
C18.0	Malignant neoplasm of cecum
C18.2	Malignant neoplasm of ascending colon
C18.3	Malignant neoplasm of hepatic flexure
C18.4	Malignant neoplasm of transverse colon
C18.5	Malignant neoplasm of splenic flexure
C18.6	Malignant neoplasm of descending colon
C18.7	Malignant neoplasm of sigmoid colon
C18.8	Malignant neoplasm of overlapping sites of colon
C18.9	Malignant neoplasm of colon, unspecified
C19	Malignant neoplasm of rectosigmoid junction

ICD-10	ICD-10 Description
C20	Malignant neoplasm of rectum
C21.8	Malignant neoplasm of overlapping sites of rectum, anus and anal canal
C22.0	Liver cell carcinoma
C22.8	Malignant neoplasm of liver, primary, unspecified as to type
C22.9	Malignant neoplasm of liver, not specified as primary or secondary
C33	Malignant neoplasm of trachea
C34.00	Malignant neoplasm of unspecified main bronchus
C34.01	Malignant neoplasm of right main bronchus
C34.02	Malignant neoplasm of left main bronchus
C34.10	Malignant neoplasm of upper lobe, unspecified bronchus or lung
C34.11	Malignant neoplasm of upper lobe, right bronchus or lung
C34.12	Malignant neoplasm of upper lobe, left bronchus or lung
C34.2	Malignant neoplasm of middle lobe, bronchus or lung
C34.30	Malignant neoplasm of lower lobe, unspecified bronchus or lung
C34.31	Malignant neoplasm of lower lobe, right bronchus or lung
C34.32	Malignant neoplasm of lower lobe, left bronchus or lung
C34.80	Malignant neoplasm of overlapping sites of unspecified bronchus and lung
C34.81	Malignant neoplasm of overlapping sites of right bronchus and lung
C34.82	Malignant neoplasm of overlapping sites of left bronchus and lung
C34.90	Malignant neoplasm of unspecified part of unspecified bronchus or lung
C34.91	Malignant neoplasm of unspecified part of right bronchus or lung
C34.92	Malignant neoplasm of unspecified part of left bronchus or lung
C37	Malignant neoplasm of thymus
C43.0	Malignant melanoma of lip
C43.10	Malignant melanoma of unspecified eyelid, including canthus
C43.111	Malignant melanoma of right upper eyelid, including canthus
C43.112	Malignant melanoma of right lower eyelid, including canthus
C43.121	Malignant melanoma of left upper eyelid, including canthus
C43.122	Malignant melanoma of left lower eyelid, including canthus
C43.20	Malignant melanoma of unspecified ear and external auricular canal
C43.21	Malignant melanoma of right ear and external auricular canal

Medical Necessity Criteria

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

ICD-10	ICD-10 Description
C43.22	Malignant melanoma of left ear and external auricular canal
C43.30	Malignant melanoma of unspecified part of face
C43.31	Malignant melanoma of nose
C43.39	Malignant melanoma of other parts of face
C43.4	Malignant melanoma of scalp and neck
C43.51	Malignant melanoma of anal skin
C43.52	Malignant melanoma of skin of breast
C43.59	Malignant melanoma of other part of trunk
C43.60	Malignant melanoma of unspecified upper limb, including shoulder
C43.61	Malignant melanoma of right upper limb, including shoulder
C43.62	Malignant melanoma of left upper limb, including shoulder
C43.70	Malignant melanoma of unspecified lower limb, including hip
C43.71	Malignant melanoma of right lower limb, including hip
C43.72	Malignant melanoma of left lower limb, including hip
C43.8	Malignant melanoma of overlapping sites of skin
C43.9	Malignant melanoma of skin, unspecified
C45.1	Mesothelioma of peritoneum
C45.2	Mesothelioma of pericardium
C45.7	Mesothelioma of other sites
C45.9	Mesothelioma, unspecified
C49.0	Malignant neoplasm of connective and soft tissue of head, face and neck
C49.10	Malignant neoplasm of connective and soft tissue of unspecified upper limb, including shoulder
C49.11	Malignant neoplasm of connective and soft tissue of right upper limb including shoulder
C49.12	Malignant neoplasm of connective and soft tissue of left upper limb, including shoulder
C49.20	Malignant neoplasm of connective and soft tissue of unspecified lower limb, including hip
C49.21	Malignant neoplasm of connective and soft tissue of right lower limb, including hip
C49.22	Malignant neoplasm of connective and soft tissue of left lower limb, including hip
C49.3	Malignant neoplasm of connective and soft tissue of thorax
C49.4	Malignant neoplasm of connective and soft tissue of abdomen
C49.5	Malignant neoplasm of connective and soft tissue of pelvis
C49.6	Malignant neoplasm of connective and soft tissue of trunk, unspecified

Medical Necessity Criteria

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

ICD-10	ICD-10 Description
C49.8	Malignant neoplasm of overlapping sites of connective and soft tissue
C49.9	Malignant neoplasm of connective and soft tissue, unspecified
C53.0	Malignant neoplasm of endocervix
C53.1	Malignant neoplasm of exocervix
C53.8	Malignant neoplasm of overlapping sites of cervix uteri
C53.9	Malignant neoplasm of cervix uteri, unspecified
C67.0	Malignant neoplasm of trigone of bladder
C67.1	Malignant neoplasm of dome of bladder
C67.2	Malignant neoplasm of lateral wall of bladder
C67.3	Malignant neoplasm of anterior wall of bladder
C67.4	Malignant neoplasm of posterior wall of bladder
C67.5	Malignant neoplasm of bladder neck
C67.6	Malignant neoplasm of ureteric orifice
C67.7	Malignant neoplasm of urachus
C67.8	Malignant neoplasm of overlapping sites of bladder
C67.9	Malignant neoplasm of bladder, unspecified
C7A.1	Malignant poorly differentiated neuroendocrine tumors
C78.00	Secondary malignant neoplasm of unspecified lung
C78.01	Secondary malignant neoplasm of right lung
C78.02	Secondary malignant neoplasm of left lung
C78.6	Secondary malignant neoplasm of retroperitoneum and peritoneum
C78.7	Secondary malignant neoplasm of liver and intrahepatic bile duct
C83.00	Small cell B-cell lymphoma, unspecified site
C83.01	Small cell B-cell lymphoma, lymph nodes of head, face, and neck
C83.02	Small cell B-cell lymphoma, intrathoracic lymph nodes
C83.03	Small cell B-cell lymphoma, intra-abdominal lymph nodes
C83.04	Small cell B-cell lymphoma, lymph nodes of axilla and upper limb
C83.05	Small cell B-cell lymphoma, lymph nodes of inguinal region and lower limb
C83.06	Small cell B-cell lymphoma, intrapelvic lymph nodes
C83.07	Small cell B-cell lymphoma, spleen
C83.08	Small cell B-cell lymphoma, lymph nodes of multiple sites

Medical Necessity Criteria

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

ICD-10	ICD-10 Description
C83.09	Small cell B-cell lymphoma, extranodal and solid organ sites
C83.30	Diffuse large B-cell lymphoma, unspecified site
C83.31	Diffuse large B-cell lymphoma, lymph nodes of head, face, and neck
C83.32	Diffuse large B-cell lymphoma, intrathoracic lymph nodes
C83.33	Diffuse large B-cell lymphoma, intra-abdominal lymph nodes
C83.34	Diffuse large B-cell lymphoma, lymph nodes of axilla and upper limb
C83.35	Diffuse large B-cell lymphoma, lymph nodes of inguinal region and lower limb
C83.36	Diffuse large B-cell lymphoma, intrapelvic lymph nodes
C83.37	Diffuse large B-cell lymphoma, spleen
C83.38	Diffuse large B-cell lymphoma, lymph nodes of multiple sites
C83.39	Diffuse large B-cell lymphoma, extranodal and solid organ sites
C91.10	Chronic lymphocytic leukemia of B-cell type not having achieved remission
C91.12	Chronic lymphocytic leukemia of B-cell type in relapse
D09.0	Carcinoma in situ of bladder
D15.0	Benign neoplasm of thymus
D38.4	Neoplasm of uncertain behavior of thymus
Z85.038	Personal history of other malignant neoplasm of large intestine
Z85.068	Personal history of other malignant neoplasm of small intestine
Z85.118	Personal history of other malignant neoplasm of bronchus and lung
Z85.12	Personal history of malignant neoplasm of trachea
Z85.238	Personal history of other malignant neoplasm of thymus
Z85.51	Personal history of malignant neoplasm of bladder
Z85.831	Personal history of malignant neoplasm of soft tissue

Medical Necessity Criteria

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

Appendix 2 – Centers for Medicare and Medicaid Services (CMS)

The preceding information is intended for non-Medicare coverage determinations. Medicare coverage for outpatient (Part B) drugs is outlined in the Medicare Benefit Policy Manual (Pub. 100-2), Chapter 15, §50 Drugs and Biologicals. In addition, National Coverage Determinations (NCDs) and/or Local Coverage Determinations (LCDs) may exist and compliance with these policies is required where applicable. Local Coverage Articles (LCAs) may also exist for claims payment purposes or to clarify benefit eligibility under Part B for drugs which may be self-administered. The following link may be used to search for NCD, LCD, or LCA documents: <https://www.cms.gov/medicare-coverage-database/search.aspx>. Additional indications, including any preceding information, may be applied at the discretion of the health plan.

Medicare Part B Covered Diagnosis Codes (applicable to existing NCD/LCD/LCA): N/A

Medicare Part B Administrative Contractor (MAC) Jurisdictions		
Jurisdiction	Applicable State/US Territory	Contractor
E (1)	CA, HI, NV, AS, GU, CNMI	Noridian Healthcare Solutions, LLC
F (2 & 3)	AK, WA, OR, ID, ND, SD, MT, WY, UT, AZ	Noridian Healthcare Solutions, LLC
5	KS, NE, IA, MO	Wisconsin Physicians Service Insurance Corp (WPS)
6	MN, WI, IL	National Government Services, Inc. (NGS)
H (4 & 7)	LA, AR, MS, TX, OK, CO, NM	Novitas Solutions, Inc.
8	MI, IN	Wisconsin Physicians Service Insurance Corp (WPS)
N (9)	FL, PR, VI	First Coast Service Options, Inc.
J (10)	TN, GA, AL	Palmetto GBA
M (11)	NC, SC, WV, VA (excluding below)	Palmetto GBA
L (12)	DE, MD, PA, NJ, DC (includes Arlington & Fairfax counties and the city of Alexandria in VA)	Novitas Solutions, Inc.
K (13 & 14)	NY, CT, MA, RI, VT, ME, NH	National Government Services, Inc. (NGS)
15	KY, OH	CGS Administrators, LLC