

Trabectedin **Evdi; Yondelis®** (Intravenous)

-E-

Document Number: IC-0402

Date Reviewed: 06/2026**Date of Origin: 10/30/2018****Dates Approved: 10/2018, 01/2019, 04/2019, 07/2019, 10/2019, 01/2020, 04/2020, 07/2020, 10/2020, 01/2021, 04/2021, 07/2021, 10/2021, 02/2022, 05/2022, 07/2022, 10/2022, 01/2023, 04/2023, 07/2023, 10/2023, 01/2024, 05/2024, 06/2024, 10/02/2025, 07/01/2026**

I. Length of Authorization

- Initial: Prior authorization validity will be provided initially for 6 months (180 days).
- Renewal: Prior authorization validity may be renewed every 6 months (180 days) thereafter.

II. Dosing Limits

Max Units (per dose and over time) [HCPCS Unit]:**Yondelis**

- All Indications: 40 billable units every 21 days

Evdi

- All Indications: 4 mg every 21 days

III. Initial Approval Criteria ^{1,12}

Prior authorization validity is provided in the following conditions:

- Member is at least 18 years of age; **AND**

Universal Criteria ^{1,12}

- Left ventricular ejection fraction (LVEF) is within normal limits prior to initiating therapy and will be assessed at regular intervals (e.g., every 3 months) during treatment; **AND**

Soft Tissue Sarcoma (STS) ‡ ^{1-4,8-12,1e,3e,6e,7e,8e,20e,22e,23e}

- Used in combination with doxorubicin; **AND**
 - Member has leiomyosarcoma; **AND**
 - Used as first-line therapy; **AND**
 - Member has advanced, metastatic, unresectable, or recurrent disease of the extremity/body wall/head-neck; **OR**

- Member has advanced, metastatic, unresectable, or residual disease (R2 resection) of the retroperitoneal or intra-abdominal area; **OR**
- Used as single agent therapy; **AND**
 - Member has liposarcoma or leiomyosarcoma † Φ ; **AND**
 - Used for unresectable or metastatic disease after an anthracycline-containing regimen (e.g., doxorubicin, liposomal doxorubicin, epirubicin, etc.); **AND**

Liposarcoma ONLY:

- Member must also demonstrate an inadequate response to eribulin, unless there is a contraindication or intolerance, prior to approval of trabectedin; **OR**

- Used as palliative therapy for solitary fibrous tumor; **AND**
 - Member has not received prior cytotoxic chemotherapy; **OR**
- Member has dedifferentiated liposarcoma with or without concurrent well-differentiated liposarcoma; **AND**
 - Used for advanced disease after prior therapy with an anthracycline, unless contraindicated; **AND**

- Member must also demonstrate an inadequate response to eribulin, unless there is a contraindication or intolerance, prior to approval of trabectedin; **OR**

- Used as neoadjuvant therapy; **AND**
 - Member has myxoid liposarcoma; **AND**
 - Extremity/Body Wall, Head/Neck; **AND**
 - Used for stage III or stage IV (any T, N1, M0) resectable disease with acceptable functional outcomes; **AND**

- Use of trabectedin will be restricted to patients with a contraindication or intolerance to generic chemotherapy (e.g., AIM, etc. [see *NCCN Soft Tissue Sarcoma guidelines for complete list of alternative regimens*]); **OR**

- Retroperitoneal/Intra-Abdominal; **AND**
 - Used for resectable primary or recurrent localized disease at high risk of becoming metastatic OR if downstaging is needed to facilitate resection (*Note: Systemic therapy is not recommended for low-grade tumors*); **AND**

- ♦ Use of trabectedin will be restricted to patients with a contraindication or intolerance to generic chemotherapy (e.g., AIM, etc. [see *NCCN Soft Tissue Sarcoma guidelines for complete list of alternative regimens*]); **OR**

- Used for unfavorable tumor biology for stage IV disease with single organ and limited tumor bulk that are amenable to local therapy; **AND**

♦ Use of trabectedin will be restricted to patients with a contraindication or intolerance to generic chemotherapy (e.g., AIM, etc. [see *NCCN Soft Tissue Sarcoma guidelines for complete list of alternative regimens*]); **OR**

- Used as subsequent palliative therapy; **AND**
 - Retroperitoneal/Intra-Abdominal**; **AND**
 - Used for stage IV disease with disseminated metastases; **AND**

Liposarcoma ONLY:

- Patient must also demonstrate an inadequate response to eribulin, unless there is a contraindication or intolerance, prior to approval of trabectedin; **OR**

- Used as alternative systemic therapy for unresectable or progressive disease after receiving initial therapy for unresectable or localized disease; **AND**

Liposarcoma ONLY:

- Patient must also demonstrate an inadequate response to eribulin, unless there is a contraindication or intolerance, prior to approval of trabectedin; **OR**

- Extremity/Body Wall, Head/Neck*; **AND**
 - Used for advanced or metastatic disease with disseminated metastases; **AND**

Liposarcoma ONLY:

- Patient must also demonstrate an inadequate response to eribulin, unless there is a contraindication or intolerance, prior to approval of trabectedin

** For atypical lipomatous tumor/well-differentiated liposarcoma (ALT/WDLPS) of the extremity, abdominal wall, trunk that was initially diagnosed as ALT/WDLPS and shows evidence of de-differentiation, treat as other soft tissue sarcomas.*

*** For well-differentiated liposarcoma (WDLPS-retroperitoneum, paratesticular) with or without evidence of de-differentiation, treat as other soft tissue sarcomas.*

Uterine Sarcoma ‡^{2,5,8}

- Member has uterine leiomyosarcoma (uLMS); **AND**
- Member has advanced, recurrent/metastatic, or inoperable disease; **AND**
 - Used as subsequent therapy after an anthracycline-containing regimen; **AND**
 - Used as a single agent therapy; **OR**
 - Used as first-line therapy; **AND**
 - Used in combination with doxorubicin

Preferred therapies and recommendations are determined by review of clinical evidence. NCCN category of recommendation is taken into account as a component of this review. Regimens deemed equally efficacious (i.e., those having the same NCCN categorization) are considered to be therapeutically equivalent.

Enhanced Oncology Value (EOV) Program – Redacted indications

Uses not listed above have inadequate data to support efficacy and are excluded from prior authorization validity.

Other treatment options including, but are not limited to, the following may be appropriate: radiation therapy, surgery, traditional chemotherapy (e.g., platinum, taxane), compassionate use/expanded access programs, clinical trials, supportive care, integrative and complementary therapies.

† FDA Approved Indication(s); ‡ Compendia Recommended Indication(s); Ⓢ Orphan Drug

IV. Renewal Criteria ^{1,12}

Prior authorization validity may be renewed based upon the following criteria:

- Member continues to meet the universal and other indication-specific relevant criteria such as concomitant therapy requirements (not including prerequisite therapy), performance status, etc. identified in section III; **AND**
- Disease response with treatment as defined by stabilization of disease or decrease in size of tumor or tumor spread; **AND**
- Absence of unacceptable toxicity from the drug. Examples of unacceptable toxicity include: cardiomyopathy, rhabdomyolysis, hepatotoxicity and/or severe hepatic impairment, capillary leak syndrome (CLS), severe neutropenia/neutropenic sepsis, extravasation resulting in tissue necrosis, etc.; **AND**
- Left ventricular ejection fraction (LVEF) has not had an absolute decrease of $\geq 15\%$ from baseline OR is not below the lower limit of normal (LLN) with an absolute decrease of $\geq 5\%$ (LVEF results must be within the previous 3 months)

V. Dosage/Administration ^{1,6-8,12}

Indication	Dose
Soft Tissue Sarcoma	<u>Single agent therapy</u> Administer 1.5 mg/m² intravenously every 21 days, until disease progression or unacceptable toxicity <u>Single agent therapy (myxoid liposarcoma)</u>

	Administer 1.3 mg/m² intravenously every 21 days, until disease progression or unacceptable toxicity <u>In combination with doxorubicin (leiomyosarcoma ONLY as first-line therapy)</u> Administer 1.1 mg/m² intravenously, with doxorubicin, every 21 days for up to 6 cycles, followed by single agent maintenance treatment at a dose of 1.1 mg/m² every 21 days until disease progression or unacceptable toxicity
Uterine Sarcoma	<u>In combination with doxorubicin (first-line therapy)</u> Administer 1.1 mg/m² intravenously, with doxorubicin, every 21 days for up to 6 cycles, followed by single agent maintenance treatment at a dose of 1.1 mg/m² every 21 days until disease progression or unacceptable toxicity <u>Single agent therapy (subsequent therapy)</u> Administer 1.5 mg/m² intravenously every 21 days, until disease progression or unacceptable toxicity

VI. Billing Code/Availability Information

HCPCS Code(s):

- J9352 – Injection, trabectedin, 0.1 mg; 1 billable unit = 0.1 mg (*Yondelis Only*)
- J9999 – Not otherwise classified, antineoplastic drugs (*Evdi Only*)

NDC(s):

- Yondelis 1 mg single-dose vial for injection: 59676-0610-xx
- Evdi 1 mg/20 mL solution in a single-dose vial: 60505-6423-xx

VII. References (STANDARD)

1. Yondelis [package insert]. Horsham, PA; Janssen Products, LP; January 2026. Accessed June 2026.
2. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) trabectedin. National Comprehensive Cancer Network, 2026. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed June 2026.
3. Demetri GD, von Mehren M, Jones RL, et al. Efficacy and Safety of Trabectedin or Dacarbazine for Metastatic Liposarcoma or Leiomyosarcoma After Failure of Conventional Chemotherapy: Results of a Phase III Randomized Multicenter Clinical Trial. J Clin Oncol. 2016;34(8):786-793.

4. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) Soft Tissue Sarcoma Version 3.2026. National Comprehensive Cancer Network, 2026. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the NCCN Guidelines, go online to NCCN.org. Accessed June 2026.
5. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) Uterine Neoplasms Version 3.2026. National Comprehensive Cancer Network, 2026. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the NCCN Guidelines, go online to NCCN.org. Accessed June 2026.
6. Gronchi A, Ferrari S, Quagliuolo V, et al. Histotype-tailored neoadjuvant chemotherapy versus standard chemotherapy in patients with high-risk soft-tissue sarcomas (ISG-STs 1001): an international, open-label, randomised, controlled, phase 3, multicentre trial. *Lancet Oncol*. 2017 Jun;18(6):812-822. doi: 10.1016/S1470-2045(17)30334-0. Epub 2017 May 9.
7. Hensley ML, Patel SR, von Mehren M, et al. Efficacy and safety of trabectedin or dacarbazine in patients with advanced uterine leiomyosarcoma after failure of anthracycline-based chemotherapy: Subgroup analysis of a phase 3, randomized clinical trial. *Gynecol Oncol*. 2017 Sep;146(3):531-537. doi: 10.1016/j.ygyno.2017.06.018.
8. Pautier P, Italiano A, Neumann S, et al. Doxorubicin alone versus doxorubicin with trabectedin followed by trabectedin alone as first-line therapy for metastatic or unresectable leiomyosarcoma (LMS-04): a randomised, multicentre, open-label phase 3 trial. *Lancet Oncol*. 2022 Aug;23(8):1044-1054. doi: 10.1016/S1470-2045(22)00380-1. Epub 2022 Jul 11.
9. Pautier P, Floquet A, Chevreau C, et al. A single-arm multicentre phase II trial of doxorubicin in combination with trabectedin in the first-line treatment for leiomyosarcoma with long-term follow-up and impact of cytoreductive surgery. *ESMO Open*. 2021 Aug;6(4):100209. doi: 10.1016/j.esmoop.2021.100209. Epub 2021 Jul 26.
10. Kawai A, Araki N, Sugiura H, et al. Trabectedin monotherapy after standard chemotherapy versus best supportive care in patients with advanced, translocation-related sarcoma: a randomised, open-label, phase 2 study. *Lancet Oncol*. 2015 Apr;16(4):406-16. doi: 10.1016/S1470-2045(15)70098-7. Epub 2015 Mar 18. PMID: 25795406.
11. Samuels BL, Chawla S, Patel S, von Mehren M, Hamm J, Kaiser PE, Schuetze S, Li J, Aymes A, Demetri GD. Clinical outcomes and safety with trabectedin therapy in patients with advanced soft tissue sarcomas following failure of prior chemotherapy: results of a worldwide expanded access program study. *Ann Oncol*. 2013 Jun;24(6):1703-9. doi: 10.1093/annonc/mds659. Epub 2013 Feb 5. PMID: 23385197.
12. Evdi [package insert]. Weston, FL; Apotex Corp.; May 2026. Accessed June 2026.

VIII. References (ENHANCED)

- 1e. Schöffski P, Chawla S, Maki RG, et al. Eribulin versus dacarbazine in previously treated patients with advanced liposarcoma or leiomyosarcoma: a randomised, open-label, multicentre, phase 3 trial. *Lancet*. 2016 Apr 16;387(10028):1629-37. doi: 10.1016/S0140-6736(15)01283-0. Epub 2016 Feb 10.
- 2e. Demetri GD, Schöffski P, Grignani G, et al. Activity of Eribulin in Patients With Advanced Liposarcoma Demonstrated in a Subgroup Analysis From a Randomized Phase III Study of Eribulin Versus Dacarbazine. *J Clin Oncol*. 2017 Oct 20;35(30):3433-3439. doi: 10.1200/JCO.2016.71.6605. Epub 2017 Aug 30.
- 3e. Bui-Nguyen B, Butrynski JE, Penel N, et al. A phase IIb multicentre study comparing the efficacy of trabectedin to doxorubicin in patients with advanced or metastatic untreated soft tissue sarcoma: the TRUSTS trial. *Eur J Cancer*. 2015 Jul;51(10):1312-20.
- 4e. Seddon B, Strauss SJ, Whelan J, et al. Gemcitabine and docetaxel versus doxorubicin as first-line treatment in previously untreated advanced unresectable or metastatic soft-tissue sarcomas (GeDDiS): a randomised controlled phase 3 trial. *Lancet Oncol*. 2017 Oct;18(10):1397-1410.
- 5e. Le Cesne A, Blay J-Y, Cupissol D, Italiano A, Delcambre C, Penel N, et al. Results of a prospective randomized phase III T-SAR trial comparing trabectedin vs best supportive care (BSC) in patients with pretreated advanced soft tissue sarcoma (ASTS) *Ann Oncol*. 2016;27(suppl 6):1396O.
- 6e. Le Cesne A, Blay JY, Judson I, et al. Phase II study of ET-743 in advanced soft tissue sarcomas: a European Organisation for the Research and Treatment of Cancer (EORTC) soft tissue and bone sarcoma group trial. *J Clin Oncol*. 2005 Jan 20;23(3):576-84.
- 7e. Blay JY, Leahy MG, Nguyen BB, et al. Randomised phase III trial of trabectedin versus doxorubicin-based chemotherapy as first-line therapy in translocation-related sarcomas. *Eur J Cancer*. 2014 Apr;50(6):1137-47. doi: 10.1016/j.ejca.2014.01.012. Epub 2014 Feb 7.
- 8e. Baruchel S, Pappo A, Krailo M, et al. A phase 2 trial of trabectedin in children with recurrent rhabdomyosarcoma, Ewing sarcoma and non-rhabdomyosarcoma soft tissue sarcomas: a report from the Children's Oncology Group. *Eur J Cancer*. 2012 Mar;48(4):579-85. doi: 10.1016/j.ejca.2011.09.027. Epub 2011 Nov 14.
- 9e. Arndt CA, Stoner JA, Hawkins DS, et al. Vincristine, actinomycin, and cyclophosphamide compared with vincristine, actinomycin, and cyclophosphamide alternating with vincristine, topotecan, and cyclophosphamide for intermediate-risk rhabdomyosarcoma: children's oncology group study D9803. *J Clin Oncol*. 2009 Nov 1;27(31):5182-8. doi: 10.1200/JCO.2009.22.3768. Epub 2009 Sep 21.
- 10e. Penel N, Bui BN, Bay JO, et al. Phase II trial of weekly paclitaxel for unresectable angiosarcoma: the ANGIOTAX Study. *J Clin Oncol*. 2008 Nov 10;26(32):5269-74. doi: 10.1200/JCO.2008.17.3146. Epub 2008 Sep 22.

- 11e. Hensley ML, Patel SR, von Mehren M, et al. Efficacy and safety of trabectedin or dacarbazine in patients with advanced uterine leiomyosarcoma after failure of anthracycline-based chemotherapy: Subgroup analysis of a phase 3, randomized clinical trial. *Gynecol Oncol*. 2017 Sep;146(3):531-537. doi: 10.1016/j.ygyno.2017.06.018. Epub 2017 Jun 24.
- 12e. Hensley ML, Miller A2, O'Malley DM, et al. Randomized phase III trial of gemcitabine plus docetaxel plus bevacizumab or placebo as first-line treatment for metastatic uterine leiomyosarcoma: an NRG Oncology/Gynecologic Oncology Group study. *J Clin Oncol*. 2015 Apr 1;33(10):1180-5. doi: 10.1200/JCO.2014.58.3781. Epub 2015 Feb 23.
- 13e. Hensley ML, Blessing JA, Mannel R, Rose PG. Fixed-dose rate gemcitabine plus docetaxel as first-line therapy for metastatic uterine leiomyosarcoma: a Gynecologic Oncology Group phase II trial. *Gynecol Oncol*. 2008 Jun;109(3):329-34. doi: 10.1016/j.ygyno.2008.03.010.
- 14e. Agulnik M, Yarber JL, Okuno SH, et al. An open-label, multicenter, phase II study of bevacizumab for the treatment of angiosarcoma and epithelioid hemangioendotheliomas. *Ann Oncol*. 2013 Jan;24(1):257-63. doi: 10.1093/annonc/mds237. Epub 2012 Aug 21.
- 15e. Hawkins DS, Chi YY, Anderson JR, et al. Addition of Vincristine and Irinotecan to Vincristine, Dactinomycin, and Cyclophosphamide Does Not Improve Outcome for Intermediate-Risk Rhabdomyosarcoma: A Report From the Children's Oncology Group. *J Clin Oncol*. 2018 Sep 20;36(27):2770-2777. doi: 10.1200/JCO.2018.77.9694. Epub 2018 Aug 9.
- 16e. Muss HB, Bundy B, DiSaia PJ, et al. Treatment of recurrent or advanced uterine sarcoma. A randomized trial of doxorubicin versus doxorubicin and cyclophosphamide (a phase III trial of the Gynecologic Oncology Group). *Cancer*. 1985 Apr 15;55(8):1648-53.
- 17e. Omura GA, Major FJ, Blessing JA, Sedlacek TV, Thigpen JT, Creasman WT, Zaino RJ. A randomized study of adriamycin with and without dimethyl triazenoimidazole carboxamide in advanced uterine sarcomas. *Cancer*. 1983 Aug 15;52(4):626-32.
- 18e. Kawai A, Araki N, Naito Y, et al. Phase 2 study of eribulin in patients with previously treated advanced or metastatic soft tissue sarcoma. *Jpn J Clin Oncol*. 2017 Feb 1;47(2):137-144. doi: 10.1093/jjco/hyw175.
- 19e. Pautier P, Floquet A, Penel N, et al. Randomized multicenter and stratified phase II study of gemcitabine alone versus gemcitabine and docetaxel in patients with metastatic or relapsed leiomyosarcomas: a Federation Nationale des Centres de Lutte Contre le Cancer (FNCLCC) French Sarcoma Group Study (TAXOGEM study). *Oncologist*. 2012;17(9):1213-20. Epub 2012 Aug 20.
- 20e. Drilon A, Laetsch TW, Kummar S, et al. Efficacy of Larotrectinib in TRK Fusion-Positive Cancers in Adults and Children. *N Engl J Med*. 2018;378(8):731–739.
- 21e. Gronchi A, Ferrari S, Quagliuolo V, et al. Histotype-tailored neoadjuvant chemotherapy versus standard chemotherapy in patients with high-risk soft-tissue sarcomas (ISG-STs 1001): an international, open-label, randomised, controlled, phase 3, multicentre trial. *Lancet Oncol*. 2017 Jun;18(6):812-822. doi: 10.1016/S1470-2045(17)30334-0. Epub 2017 May 9. Erratum in: *Lancet Oncol*. 2017 Jun;18(6):e301.

- 22e. Constantinidou A, Jones RL, Olmos D, et al. Conventional anthracycline-based chemotherapy has limited efficacy in solitary fibrous tumour. *Acta Oncol.* 2012 Apr;51(4):550-4. doi: 10.3109/0284186X.2011.626450.
- 23e. van der Graaf WT, Blay JY, Chawla SP, et al. Pazopanib for metastatic soft-tissue sarcoma (PALETTE): a randomised, double-blind, placebo-controlled phase 3 trial. *Lancet.* 2012 May 19;379(9829):1879-86. doi: 10.1016/S0140-6736(12)60651-5.
- 24e. Talbot SM, Keohan ML, Hesdorffer M, et al. A phase II trial of temozolomide in patients with unresectable or metastatic soft tissue sarcoma. *Cancer.* 2003 Nov 1;98(9):1942-6. doi: 10.1002/cncr.11730.
- 25e. Sutton GP, Blessing JA, Barrett RJ, McGehee R. Phase II trial of ifosfamide and mesna in leiomyosarcoma of the uterus: a Gynecologic Oncology Group study. *Am J Obstet Gynecol.* 1992 Feb;166(2):556-9. doi: 10.1016/0002-9378(92)91671-v.
- 26e. Stacchiotti S, Grignani G, Vincenzi B et al. Trabectedin (T) versus adriamycin plus dacarbazine (A-DA) in advanced solitary fibrous tumor (SFT): Results from a phase II randomised clinical study (STRADA). *J Clin Oncol.* 2023; 41(16_suppl):11571. doi.org/10.1200/JCO.2023.41.16_suppl.1157
- 27e. Fabbroni C, Grignani G, Vincenzi B et al. TRAbectedin in adVanced rEtroperitoneal well differentiated/dedifferentiated Liposarcoma and Leiomyosarcoma (TRAVELL): results of a phase II study from the Italian Sarcoma Group. *ESMO Open.* 2024 Aug;9(8):103667. doi: 10.1016/j.esmoop.2024.103667. Epub 2024 Aug 8.
- 28e. Maki RG, Wathen JK, Patel SR, et al. Randomized phase II study of gemcitabine and docetaxel compared with gemcitabine alone in patients with metastatic soft tissue sarcomas: results of sarcoma alliance for research through collaboration study 002 [corrected]. *J Clin Oncol.* 2007 Jul 1;25(19):2755-63.
- 29e. Gronchi A, Bui BN, Bonvalot S, et al. Phase II clinical trial of neoadjuvant trabectedin in patients with advanced localized myxoid liposarcoma. *Ann Oncol.* 2012 Mar;23(3):771-776.
- 30e. Endo M, et al. Final results of a randomized phase II trial of second-line treatment for advanced soft tissue sarcoma comparing trabectedin, eribulin, and pazopanib: 2ND-STEP study (JCOG1802). *J Clin Oncol.* 2026;44(suppl 16):11508.
- 31e. Judson I, Verweij J, Gelderblom H, et al; European Organisation and Treatment of Cancer Soft Tissue and Bone Sarcoma Group. Doxorubicin alone versus intensified doxorubicin plus ifosfamide for first-line treatment of advanced or metastatic soft-tissue sarcoma: a randomised controlled phase 3 trial. *Lancet Oncol.* 2014 Apr;15(4):415-23.
- 32e. D'Ambrosio L, Touati N, Blay JY, et al; European Organization for Research and Treatment of Cancer Soft Tissue and Bone Sarcoma Group. Doxorubicin plus dacarbazine, doxorubicin plus ifosfamide, or doxorubicin alone as a first-line treatment for advanced leiomyosarcoma: A propensity score matching analysis from the European Organization for Research and Treatment of Cancer Soft Tissue and Bone Sarcoma Group. *Cancer.* 2020 Jun 1;126(11):2637-2647.

33e. Gronchi A, Palmerini E, Quagliuolo V, et al. Neoadjuvant Chemotherapy in High-Grade Myxoid Liposarcoma: Results of the Expanded Cohort of a Randomized Trial From Italian (ISG), Spanish (GEIS), French (FSG), and Polish Sarcoma Groups (PSG). J Clin Oncol. 2024 Mar 10;42(8):898-906.

34e. Prime Therapeutics Management. Trabectedin Clinical Literature Review Analysis. Last updated June 2026. Accessed June 2026.

Appendix A – Non-Quantitative Treatment Limitations (NQTL) Factor Checklist

Non-quantitative treatment limitations (NQTLs) refer to the methods, guidelines, standards of evidence, or other conditions that can restrict how long or to what extent benefits are provided under a health plan. These may include things like utilization review or prior authorization. The utilization management NQTL applies comparably, and not more stringently, to mental health/substance use disorder (MH/SUD) Medical Benefit Prescription Drugs and medical/surgical (M/S) Medical Benefit Prescription Drugs. The table below lists the factors that were considered in designing and applying prior authorization to this drug/drug group, and a summary of the conclusions that Prime’s assessment led to for each.

Factor	Conclusion
Indication	Yes: Consider for PA
Safety and efficacy	No: PA not a priority
Potential for misuse/abuse	No: PA not a priority
Cost of drug	Yes: Consider for PA

Appendix 1 – Covered Diagnosis Codes

ICD-10	ICD-10 Description
C47.0	Malignant neoplasm of peripheral nerves of head, face and neck
C47.10	Malignant neoplasm of peripheral nerves of unspecified upper limb, including shoulder
C47.11	Malignant neoplasm of peripheral nerves of right upper limb, including shoulder
C47.12	Malignant neoplasm of peripheral nerves of left upper limb, including shoulder
C47.20	Malignant neoplasm of peripheral nerves of unspecified lower limb, including hip
C47.21	Malignant neoplasm of peripheral nerves of right lower limb, including hip
C47.22	Malignant neoplasm of peripheral nerves of left lower limb, including hip
C47.3	Malignant neoplasm of peripheral nerves of thorax
C47.4	Malignant neoplasm of peripheral nerves of abdomen
C47.5	Malignant neoplasm of peripheral nerves of pelvis
C47.6	Malignant neoplasm of peripheral nerves of trunk, unspecified

ICD-10	ICD-10 Description
C47.8	Malignant neoplasm of overlapping sites of peripheral nerves and autonomic nervous system
C47.9	Malignant neoplasm of peripheral nerves and autonomic nervous system, unspecified
C48.0	Malignant neoplasm of retroperitoneum
C48.1	Malignant neoplasm of specified parts of peritoneum
C48.2	Malignant neoplasm of peritoneum, unspecified
C48.8	Malignant neoplasm of overlapping sites of retroperitoneum and peritoneum
C49.0	Malignant neoplasm of connective and soft tissue of head, face and neck
C49.10	Malignant neoplasm of connective and soft tissue of unspecified upper limb, including shoulder
C49.11	Malignant neoplasm of connective and soft tissue of right upper limb, including shoulder
C49.12	Malignant neoplasm of connective and soft tissue of left upper limb, including shoulder
C49.20	Malignant neoplasm of connective and soft tissue of unspecified lower limb, including hip
C49.21	Malignant neoplasm of connective and soft tissue of right lower limb, including hip
C49.22	Malignant neoplasm of connective and soft tissue of left lower limb, including hip
C49.3	Malignant neoplasm of connective and soft tissue of thorax
C49.4	Malignant neoplasm of connective and soft tissue of abdomen
C49.5	Malignant neoplasm of connective and soft tissue of pelvis
C49.6	Malignant neoplasm of connective and soft tissue of trunk, unspecified
C49.8	Malignant neoplasm of overlapping sites of connective and soft tissue
C49.9	Malignant neoplasm of connective and soft tissue, unspecified
C54.0	Malignant neoplasm of isthmus uteri
C54.1	Malignant neoplasm of endometrium
C54.2	Malignant neoplasm of myometrium
C54.3	Malignant neoplasm of fundus uteri
C54.8	Malignant neoplasm of overlapping sites of corpus uteri
C54.9	Malignant neoplasm of corpus uteri, unspecified
C55	Malignant neoplasm of uterus, part unspecified
D48.1	Neoplasm of uncertain behavior of connective and other soft tissue
Z85.42	Personal history of malignant neoplasm of other parts of uterus
Z85.831	Personal history of malignant neoplasm of soft tissue

Appendix 2 – Centers for Medicare and Medicaid Services (CMS)

The preceding information is intended for non-Medicare coverage determinations. Medicare coverage for outpatient (Part B) drugs is outlined in the Medicare Benefit Policy Manual (Pub. 100-2), Chapter 15,

§50 Drugs and Biologicals. In addition, National Coverage Determinations (NCDs) and/or Local Coverage Determinations (LCDs) may exist and compliance with these policies is required where applicable. Local Coverage Articles (LCAs) may also exist for claims payment purposes or to clarify benefit eligibility under Part B for drugs which may be self-administered. The following link may be used to search for NCD, LCD, or LCA documents: <https://www.cms.gov/medicare-coverage-database/search.aspx>. Additional indications, including any preceding information, may be applied at the discretion of the health plan.

Medicare Part B Covered Diagnosis Codes (applicable to existing NCD/LCD/LCA): N/A

Medicare Part B Administrative Contractor (MAC) Jurisdictions		
Jurisdiction	Applicable State/US Territory	Contractor
E (1)	CA, HI, NV, AS, GU, CNMI	Noridian Healthcare Solutions, LLC
F (2 & 3)	AK, WA, OR, ID, ND, SD, MT, WY, UT, AZ	Noridian Healthcare Solutions, LLC
5	KS, NE, IA, MO	Wisconsin Physicians Service Insurance Corp (WPS)
6	MN, WI, IL	National Government Services, Inc. (NGS)
H (4 & 7)	LA, AR, MS, TX, OK, CO, NM	Novitas Solutions, Inc.
8	MI, IN	Wisconsin Physicians Service Insurance Corp (WPS)
N (9)	FL, PR, VI	First Coast Service Options, Inc.
J (10)	TN, GA, AL	Palmetto GBA
M (11)	NC, SC, WV, VA (excluding below)	Palmetto GBA
L (12)	DE, MD, PA, NJ, DC (includes Arlington & Fairfax counties and the city of Alexandria in VA)	Novitas Solutions, Inc.
K (13 & 14)	NY, CT, MA, RI, VT, ME, NH	National Government Services, Inc. (NGS)
15	KY, OH	CGS Administrators, LLC